

CORRECTED VERSION

(19) World Intellectual Property
Organization
International Bureau



(43) International Publication Date
8 August 2002 (08.08.2002)

PCT

(10) International Publication Number
WO 2002/061113 A3

(51) International Patent Classification⁷: **A61K 39/00**,
C07H 21/04, C12N 15/00, 15/63, 15/85

(21) International Application Number:
PCT/US2002/002598

(22) International Filing Date: 1 February 2002 (01.02.2002)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/265,334 1 February 2001 (01.02.2001) US

(71) Applicant (for all designated States except US): **THE
JOHNS HOPKINS UNIVERSITY** [US/US]; 720 Rut-
land Avenue, Baltimore, MD 21205 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **WU, Tzyy-Chouu**
[—/US]; 11002 Nacirema Lane, Stevenson, MD 21153
(US). **HUNG, Chien-Fu** [—/US]; 6605 Sanzo Road,
Baltimore, MD 21209 (US).

(74) Agent: **LIVNAT, Shmuel**; Venable, Baetjer, Howard &
Civiletti, LLP, 1201 New York Avenue, NW, Suite 1000,
P.O. Box 34385, Washington, DC 20043-9998 (US).

(81) Designated States (national): AE, AG, AL, AM, AT, AU,
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU,

CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH,
GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC,
LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW,
MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG,
SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ,
VN, YU, ZA, ZM, ZW.

(84) Designated States (regional): ARIPO patent (GH, GM,
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW),
Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),
European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR,
GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent
(BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR,
NE, SN, TD, TG).

Published:
— with international search report

(88) Date of publication of the international search report:
12 December 2002

(48) Date of publication of this corrected version:
6 May 2004

(15) Information about Correction:
see PCT Gazette No. 19/2004 of 6 May 2004, Section II

For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.

(54) Title: NUCLEIC ACID DERIVED VACCINE THAT ENCODES AN ANTIGEN LINKED TO A POLYPEPTIDE THAT PRO-
MOTES ANTIGEN PRESENTATION

(57) Abstract: Improved molecular vaccines comprise nucleic acid vectors that encode a fusion polypeptide that includes polypep-
tide or peptide physically linked to an antigen. The linked polypeptide is one that (a) promotes processing of the expressed fusion
polypeptide via the MHC class I pathway and/or (b) promotes development or activity of antigen presenting cells, primarily den-
dritic cells. These vaccines employ one of several types of nucleic acid vectors, each with its own relative advantages: naked DNA
plasmids, self-replicating RNA replicons and suicidal DNA-based on viral RNA replicons. Administration of such a vaccine results
in enhance immune responses, primarily those mediated by CD8+ cytotoxic T lymphocytes, directed against the immunizing anti-
gen part of the fusion polypeptide. Such vaccines are useful against tumor antigens, viral antigens and antigens of other pathogenic
microorganisms and can be used in the prevention or treatment of diseases that include cancer and infections.



WO 2002/061113 A3

**SUPERIOR MOLECULAR VACCINE BASED ON SELF-REPLICATING RNA,
SUICIDAL DNA OR NAKED DNA VECTOR, THAT LINKS ANTIGEN WITH
POLYPEPTIDE THAT PROMOTES ANTIGEN PRESENTATION**

**STATEMENT OF RIGHTS TO INVENTIONS MADE UNDER
FEDERALLY SPONSORED RESEARCH**

This invention was made in part with federal government support under National Institutes of Health grants NIH 5 PO1 34582-01, U19 CA72108-02, RO1 CA72631-01, which provides to the United States government certain rights in this invention.

BACKGROUND OF THE INVENTION

Field of the Invention

The present invention in the fields of molecular biology, immunology and medicine relates to chimeric nucleic acids encoding fusion proteins and their use as vaccines to enhance immune responses, primarily cytotoxic T lymphocyte (CTL) responses to specific antigens such as tumor antigens. The fusion proteins comprise an antigenic polypeptide fused to a protein that promotes processing via the MHC class I pathway and/or promotes development or activity of antigen presenting cells (APCs), primarily dendritic cells (DCs). Preparation of the foregoing nucleic acid constructs as naked DNA plasmids, self-replicating RNA replicons and suicidal DNA-based viral RNA replicons confer various advantages on these molecular vaccines.

Description of the Background Art

Antigen-specific cancer immunotherapy has emerged as a promising approach because it is capable of engendering specific immunity against neoplastic cells while sparing normal cells. Increasing evidence suggests that professional antigen-presenting cells (APCs), particularly dendritic cells (DCs), are central players in this process. An effective vaccine strategy includes targeting the tumor antigen to professional APCs that in turn activate antigen-specific T cells (for review, see (Chen, CH *et al.*, *J Biomed Sci.* 5:231-52, 1998.).

Recently, DNA vaccines have become attractive as an approach for generating antigen-specific immunotherapy (for review, see (Robinson, HL *Vaccine* 15:785-778, 1997; Robinson, HL *et al.*, *Semin Immunol.* 9:271-83, 1997; Pardoll, DM *et al.*, *Immunity.* 3:165-9, 1995; Donnelly, JJ *et al.*, *Annu Rev Immunol.* 15: 617-48, 1997). The advantages of naked DNA include purity, ease of preparation and stability. In addition, DNA-based vaccines can be prepared inexpensively and rapidly in large-scale. Furthermore, multiple DNA vaccines can be administered simultaneously. However, naked DNA vaccines raise concerns such as potential integration into the host genome and cell transformation. Because they do not have the intrinsic ability to amplify *in vivo* as do viral vaccines, DNA vaccines may be more limited in their potency.

The present inventors conceived that a directing a DNA vaccine encoding an antigen (in the form of a fusion protein) to cells which activate immune responses, such as DCs, would enhance the vaccine's potency. Others demonstrated that linking DNA encoding the cytokine GM-CSF gene to DNA encoding an HIV or hepatitis C antigen enhanced the potency of DNA vaccines (Lee, AH *et al.*, *Vaccine* 17: 473-9, 1999; Lee, SW *et al.*, *J Virol.* 72: 8430-6, 1998). The chimeric GM-CSF/antigen is believed to act as an immunostimulatory signal to DCs, inducing their differentiation from an immature form (Banchereau, J *et al.*, *Nature* 392: 245-52, 1998). Since DCs and their precursors express high levels of GM-CSF receptors, the chimeric GM-CSF/antigen should target and concentrate the linked antigen to the DCs and further improve the vaccine's potency.

Use of self-replicating RNA vaccines (RNA replicons) has also been identified as an important strategy in nucleic acid vaccine development. RNA replicon vaccines may be derived from alphavirus vectors, such as Sindbis virus (Hariharan, MJ *et al.*, 1998. *J Virol* 72:950-8.), Semliki Forest virus (Berglund, PM *et al.*, 1997. *AIDS Res Hum Retroviruses* 13:1487-95; Ying, HT *et al.*, 1999. *Nat Med* 5:823-7.) or Venezuelan equine encephalitis virus (Pushko, PM *et al.*, 1997. *Virology* 239:389-401). These self-replicating and self-limiting vaccines may be administered as either (1) RNA or (2) DNA which is then transcribed into RNA replicons in cells transfected *in vitro* or *in vivo* (Berglund, PC *et al.*, 1998. *Nat Biotechnol* 16:562-5; Leitner, WW *et al.*, 2000. *Cancer Res* 60:51-5).

Self-replicating RNA infects a diverse range of cell types and allows the expression of a linked antigen of interest at high levels (Huang, HV 1996. *Curr Opin Biotechnol* 7:531-5) Because viral replication is toxic to infected host cells, such self-replicating RNA preparations eventually causes lysis of the transfected cells (Frolov, I *et al.*, 1996. *J Virol* 70:1182-90). These vectors cannot integrate into the host genome, and therefore do not raise concerns of associated with naked DNA vaccines. This is particularly important for vaccine development where target proteins are potentially oncogenic, such as human papillomavirus (HPV) E6 and E7 proteins.

The present inventors and their colleagues recently demonstrated that linkage of HPV-16 E7 antigen to *Mtb* heat shock protein 70 (Hsp70) leads to the enhancement of DNA vaccine potency (Chen, CH *et al.*, 2000. *Cancer Research* 60:1035-1042). (See also co-pending patent applications USSN 09/501,097, filed 09 February 2000; and USSN 099/421,608, filed 20 October 1999, all of which are incorporated by reference in their entirety.) Immunization with HSP complexes isolated from tumor or virus-infected cells induced potent anti-tumor immunity (Janetzki, S *et al.*, 1998. *J Immunother* 21:269-76) or antiviral immunity (Heikema, AE *et al.*, *Immunol Lett* 57:69-74). In addition, immunogenic HSP-peptide complexes could be reconstituted *in vitro* by mixing the peptides with HSPs (Ciupitu, AM *et al.*, 1998. *J Exp Med* 187:685-91). Furthermore, HSP-based protein vaccines have been created by fusing antigens to HSPs (Suzue, K *et al.*, 1996. *J Immunol* 156:873-9). The results of these investigations

point to HSPs a attractive candidates for use in immunotherapy. However, prior to the present inventors' work, HSP vaccines were all peptide/protein-based vaccines or, in more recent cases, were in the form of naked DNA. To date, there have been no reports of HSPs incorporated into self-replicating RNA vaccines.

Another molecule that stimulates growth of DC precursors and can help in generating large numbers of DCs *in vivo* is Flt3-ligand ("FL") (Maraskovsky, E *et al.*, *J Exp Med* 184: 1953-62, 1996, Shurin, MR *et al.*, *Cell Immunol.* 179: 174-84, 1997). FL has emerged as an important molecule in the development of tumor vaccines that augment numbers and action of DCs *in vivo*. Flt3, a murine tyrosine kinase receptor, first described in 1991 (Rosnet, O *et al.*, *Oncogene.* 6: 1641-50, 1991), was found to be a member of the type III receptor kinase family which includes -kit and c-fms (for review, see (Lyman, SD *Curr Opin Hematol.* 5:192-6, 1998). In hematopoietic tissues, the Flt3 expression is restricted to the CD34+ progenitor population. Flt3 has been used to identify and subsequently clone the corresponding ligand, Flt3-ligand or "FL" (Lyman, SD *et al.*, *Cell.* 75: 1157-67, 1993; Hannum, C *et al.*, *Nature.* 368: 643-8, 1994).

The predominant form of FL is synthesized as a transmembrane protein from which the soluble form is believed to be generated by proteolytic cleavage. The soluble form of FL (the extracellular domain or "ECD") is functionally similar to intact FL (Lyman, SD *et al.*, *Cell.* 75: 1157-67, 1993). These proteins function by binding to and activating unique tyrosine kinase receptors. Expression of the Flt3 receptor is primarily restricted, among hematopoietic cells, to the most primitive progenitor cells, including DC precursors. The soluble ECD of FL induced strong anti-tumor effects against several murine model tumors including fibrosarcoma (Lynch, DH *et al.*, *Nat Med.* 3: 625-31, 1997), breast cancer (Chen, K *et al* *Cancer Res.* 57: 3511-6, 1997; Braun, SE *et al.*, *Hum Gene Ther.* 10: 2141-51, 1999), liver cancer (Peron, JM *et al.*, *J Immunol.* 161: 6164-70, 1998), lung cancer (Chakravarty, PK *et al.*, *Cancer Res.* 59: 6028-32, 1999), melanoma and lymphoma (Esche, C *et al.*, *Cancer Res.* 58: 380-3, 1998).

There is a need in the art for improved molecular vaccines, such as nucleic acid vaccines, that combine potency and safety. The present invention helps meet this need by its design of novel fusion or chimeric polypeptides and nucleic acids coding therefor, that link the antigen with specialized polypeptides that promote antigen presentation by various mechanisms and that exploit delivery of these constructs by various nucleic acid vectors.

Partial List of Abbreviations used

APC, antigen presenting cell; BHK, baby hamster kidney; CMV, cytomegalovirus; CTL, cytotoxic T lymphocyte; DC, dendritic cell; ECD, extracellular domain; ELISA, enzyme-linked immunosorbent assay; FL, Flt3 ligand; GFP, green fluorescent protein; HPV, human papillomavirus; HSP, heat shock

protein; Hsp70, mycobacterial heat shock protein 70; IFN- γ , interferon- γ ; i.m., intramuscular(ly); i.v., intravenous(ly); MHC, major histocompatibility complex; PBS, phosphate-buffered saline; PCR, polymerase chain reaction; β -gal, β -galactosidase

SUMMARY OF THE INVENTION

5 Self-replicating RNA vaccines (RNA replicons) have emerged as an important strategy for nucleic acid vaccine development. The present inventors evaluated the effect of linking HPV type 16 (HPV-16) E7 as a model antigen to *Mycobacterium tuberculosis* (*Mtb*) heat shock protein 70 (Hsp70) on the potency of antigen-specific immunity generated by a Sindbis virus self-replicating RNA vector, SINrep5. The results indicated that this RNA replicon vaccine containing E7/Hsp70 fusion genes generated
10 significantly greater E7-specific T cell-mediated immunity than vaccines comprising wild type E7 DNA.

HPV-16 E7 was selected as a model antigen for vaccine development because HPVs, particularly HPV-16, are associated with most cervical cancers. HPV oncogenic proteins, E6 and E7, are co-expressed in most HPV-containing cervical cancers and are important in the induction and maintenance of cell transformation. Therefore, vaccines targeting E6 or E7 provide an opportunity to
15 prevent and treat HPV-associated cervical malignancies. HPV-16 E7 is a well-characterized cytoplasmic/nuclear protein that is more conserved than E6 in HPV-associated cancer cells; E7 has been tested in a variety of HPV vaccines.

Furthermore, *in vitro* studies demonstrated that E7 antigen from apoptotic cells that have been transfected with E7/Hsp70 RNA replicons is taken up by bone marrow-derived dendritic cells (DC's)
20 and presented more efficiently through the MHC class I pathway compared to antigen from than apoptotic cells transfected by wild-type E7 RNA replicons.

Importantly, the fusion of Hsp70 to E7 converted a less effective vaccine into one with significant potency against E7-expressing tumors. This antitumor effect involved NK cells and CD8⁺ T cells. Thus, fusion of a nucleic acid sequence encoding Hsp70 to nucleic acid encoding an antigen of interest
25 in the form of a self-replicating RNA vaccine greatly enhances the potency of this vaccine.

Naked DNA vaccines represent an attractive approach for generating antigen-specific immunity because of their stability and simplicity of delivery. Concerns with DNA vaccines include potential integration into the host genome, cell transformation, and limited potency. The use of DNA-based alphaviral RNA replicons ("suicidal DNA vectors"), as disclosed herein, may alleviate concerns
30 surrounding DNA integration or cell transformation since suicidal DNA vectors eventually cause lysis of the cells they transfect.

To further improve the potency of suicidal DNA vaccines, the present inventors linked Hsp70 to E7 (as a model antigen) using DNA-based Semliki Forest virus (SFV) RNA vector, pSCA1. This suicidal DNA vaccine containing E7/Hsp70 fusion DNA produced a significantly greater E7-specific T cell-

mediated immune response in mice than did vaccines containing the wild type E7 DNA alone. Importantly, this fusion converted a less effective vaccine into one with significant therapeutic potency against established E7-expressing metastatic tumors. The antitumor effect was dependent upon CD8+ T cells. Thus, linkage of Hsp70 to an antigen enhances the potency of a suicidal DNA vaccine.

Flt3 (fms-like tyrosine kinase 3)-ligand is an important cytokine in the development and differentiation of professional APCs, particularly DCs. A recombinant chimeric or fusion polypeptide molecule comprising the extracellular domain (ECD) of Flt3-ligand (FL) linked to an antigen targets the antigen to DCs and their precursors. Using HPV-16 E7 as a model antigen, the present inventors linked FL to E7 and caused stimulation of an antigen-specific immune response by a naked DNA vaccine administered intradermally via gene gun. Vaccines that included DNA encoding a chimeric FL-E7 fusion polypeptide dramatically increased the frequency of E7-specific CD8⁺ T cells when compared to vaccines of only E7 DNA. Cells transfected *in vitro* with FL-E7 DNA presented E7 via the MHC class I pathway more efficiently than did cells transfected with wild-type E7 DNA. Furthermore, bone marrow-derived DCs pulsed with lysates of cells that had been transfected to express an FL-E7 fusion protein presented E7 (via the MHC class I pathway) more efficiently than did DCs pulsed with lysates of cells expressing (after transfection) E7 protein alone. More importantly, this fusion construct rendered a less effective vaccine highly potent in inducing a therapeutic response against established E7-expressing metastatic tumors. The FL-E7 fusion vaccines mainly targeted CD8⁺ T cells as anti-tumor effects were completely independent of CD4⁺ T cells. Thus, fusion of DNA encoding the ECD of FL to DNA encoding an antigen markedly enhances the potency of a DNA vaccine acting via CD8-dependent pathways.

In one embodiment, the antigen (*e.g.*, the MHC class I-binding peptide epitope) is derived from a pathogen, *e.g.*, it comprises a peptide expressed by a pathogen. The pathogen can be a virus, such as, *e.g.*, a papilloma virus, a herpesvirus, a retrovirus (*e.g.*, an immunodeficiency virus, such as HIV-1), an adenovirus, and the like. The papilloma virus can be a human papilloma virus; for example, the antigen (*e.g.*, the Class I-binding peptide) can be derived from an HPV-16 E7 polypeptide. In one embodiment, the HPV-16 E7 polypeptide is substantially non-oncogenic, *i.e.*, it does not bind retinoblastoma polypeptide (pRB) or binds pRB with such low affinity that the HPV-16 E7 polypeptide is effectively non-oncogenic when expressed or delivered *in vivo*.

In alternative embodiments, the pathogen is a bacteria, such as *Bordetella pertussis*; *Ehrlichia chaffeensis*; *Staphylococcus aureus*; *Toxoplasma gondii*; *Legionella pneumophila*; *Brucella suis*; *Salmonella enterica*; *Mycobacterium avium*; *Mycobacterium tuberculosis*; *Listeria monocytogenes*; *Chlamydia trachomatis*; *Chlamydia pneumoniae*; *Rickettsia rickettsii*; or, a fungi, such as, *e.g.*, *Paracoccidioides brasiliensis*; or other pathogen, *e.g.*, *Plasmodium falciparum*.

In another embodiment, the MHC class I-binding peptide epitope is derived from a tumor cell. The tumor cell-derived peptide epitope can comprise a tumor associated antigen, *e.g.*, a tumor specific antigen, such as, *e.g.*, a HER-2/neu antigen.

In one embodiment, the isolated or recombinant nucleic acid molecule is operatively linked to a promoter, such as, *e.g.*, a constitutive, an inducible or a tissue-specific promoter. The promoter can be expressed in any cell, including cells of the immune system, including, *e.g.*, antigen presenting cells (APCs), *e.g.*, in a constitutive, an inducible or a tissue-specific manner.

In alternative embodiments, the APCs are dendritic cells, keratinocytes, astrocytes, monocytes, macrophages, B lymphocytes, a microglial cell, or activated endothelial cells, and the like.

The details of one or more embodiments of the invention are set forth in the accompanying drawings and the description below. Other features, objects, and advantages of the invention will be apparent from the description and drawings, and from the claims.

All publications, patents, patent applications, GenBank sequences and ATCC deposits, cited herein are hereby expressly incorporated by reference for all purposes.

BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1 is a schematic diagram showing the SINrep5 self-replicating RNA transcripts. A methylated M⁷G, "cap" is located at the 5' end of the mRNA, followed by a sequence responsible for the self-replication (replicase), the gene of interest (*i.e.* E7, Hsp70, E7/GFP or E7/Hsp70), and a polyadenylated tail (AAAA).

Figure 2. Antigen ppecific CD8⁺ T cell cytotoxic activity. Mice (5 per group) were immunized with various RNA vaccines via intramuscular (*i.m.*) injection. Splenocytes from each group (5 mice per group) were pooled 14 days after vaccination. To perform the cytotoxicity assay, pooled splenocytes from the various self-replicating RNA vaccines were cultured with E7 peptide (aa 49-57, RAHYNIVTF, SEQ ID NO:22, which includes a MHC class I epitope) for 6 days and used as effector cells. TC-1 tumor target cells were mixed with splenocytes at various effector/target (E/T) ratios. Cytolysis was determined by quantitative measurement of LDH release. The self-replicating RNA E7/Hsp70 vaccine generated significantly higher lysis than the other RNA vaccines (*p*<0.001). Error bars reflect 3 samples for each group. CTL assays shown are from one representative experiment of three performed.

Figure 3. IFN- γ secretion by E7-specific CD8⁺ T cells. Mice were immunized *i.m.* with various self-replicating RNA vaccines. Splenocytes were collected 14 days after vaccination. Splenocytes from various self-replicating RNA vaccines were cultured *in vitro* with E7 peptide RAHYNIVTF, with or without peptide for 6 days. The culture supernatants were collected for measurement of IFN- γ concentration by ELISA. The CD8⁺ T cells were induced by the MHC class I epitope of E7.

Splenocytes from the self-replicating E7/Hsp70 RNA group stimulated with the E7 RAHYNIVTF secreted the most IFN- γ compared to the other RNA vaccines ($p < 0.001$, one-way ANOVA). Results from the ELISA are from one representative experiment of three performed.

Figure 4. IFN- γ secretion by E7-specific CD4⁺ T cells. Splenocytes from mice vaccinated with various self-replicating RNA vaccines were cultured *in vitro* with E7 peptide containing the MHC class II epitope (aa 30-67, DSSEEEDEIDGPAGQAEPDRAHYNIVTFCKCDSTLRL, (SEQ ID NO:23), or no peptide (control). The culture supernatants were collected for measurement of IFN- γ concentrations by ELISA. There was no significant increase in secretion of IFN- γ by splenocytes from the self-replicating E7/Hsp70 RNA group stimulated with the above peptide compared to the other RNA vaccines (one-way ANOVA). Results from the ELISA are from one representative experiment of three performed.

Figure 5. Various SINrep5 self-replicating RNA vaccines induce tumor protection. Mice (5 per group) were immunized i.m. with the RNA vaccines. Two weeks later, mice were challenged with TC-1 tumor cells i.v. (tail vein) at 10^4 cells/mouse. Mice were monitored twice weekly and sacrificed on day 21 after challenge. Lungs were dissected from the mice 35 days after vaccination with SINrep5, SINrep5-Hsp70, SINrep5-E7, SINrep5-E7 mixed with SINrep5-Hsp70, SINrep5-E7/GFP or SINrep5-E7/Hsp70 RNA. The mean number of tumor nodules on the lung surface of the vaccinated mice were counted, where a decrease was an indication of vaccine effectiveness at controlling growth of a tumor expressing HPV-16 E7. There were fewer mean pulmonary nodules in mice vaccinated with self-replicating E7/Hsp70 RNA vaccines (0.1 μ g, 1 μ g, and 10 μ g) compared to mice vaccinated with the other RNA vaccines (10 μ g) ($p < 0.001$, one-way ANOVA). Self-replicating SINrep5-E7/Hsp70 RNA vaccines protect mice from intravenous tumor challenge even at the low dose of 0.1 μ g whereas mice vaccinated with 10 μ g of all the other vaccines developed numerous lung nodules. These tumor protection experiments were repeated three times with similar results.

Figure 6 shows representative photographs of lung tumors in each group. C57BL/6 mice were vaccinated i.m. with various RNA replicon-based vaccines (10 μ g/mouse) and challenged with TC-1 tumor i.v., as above. Mice were sacrificed 35 days after vaccination. There are multiple grossly visible lung tumors in unvaccinated control mice and mice vaccinated with SINrep5 or SINrep5-E7 RNA vaccines. Lung tumors in SINrep5-E7/Hsp70 RNA vaccinated group are not evident at the magnification used.

Figure 7 shows the effect of lymphocyte subset depletion on the potency of self-replicating SINrep5-E7/Hsp70 RNA vaccine. Mice were immunized with 1 μ g of this vaccine i.m.. Two weeks after vaccination, mice were challenged with 1×10^4 TC-1 cells/ mouse i.v. Depletions were initiated one week prior to tumor challenge and were carried out for 28 days. Three weeks after tumor challenge,

mice were sacrificed. The mean number of pulmonary nodules in vaccinated mice was determined as above. Depletion of CD8⁺ T cells or of NK1.1 cells resulted in a higher number of pulmonary nodules vs. animals receiving control IgG_{2a} isotype antibody. The mean number of lung nodules in mice depleted of CD4⁺ T cells were similar to those in mice receiving control antibody, indicating that CD4⁺ T cells
 5 were not critical for the antitumor effect. Depletion of NK1.1+ cells had a greater impact on antitumor activity than did loss of CD8⁺ T cells.

Figure 8 shows flow cytometric analysis of NK cells in mice immunized with various self-replicating SINrep5 RNA vaccines. Splenocytes were stained for CD3 and NK1.1 immediately, without stimulation. Fig. 8A shows numbers of NK cells in mice immunized with various self-replicating RNA
 10 vaccines. The percentage of NK cells among the spleen cells is indicated in the upper left corner. Fig. 8B is a histogram demonstrating the percentages of NK cells in vaccinated mice. The percentage of NK cells in mice immunized with self-replicating RNA vaccines was higher than in unimmunized controls. There was no significant difference between the percentage of NK cells among mice given various self-replicating RNA vaccines. These results are from one representative experiment of two
 15 performed.

Figure 9 shows apoptotic death of host cells induced by self-replicating RNA vaccines. To determine if self-replicating RNA vaccines killed host cells, RNA transcribed *in vitro* from various SINrep5 plasmids was transfected into BHK21 cells. BHK 21 cells that were electroporated in the absence of RNA or unhandled BHK21 cells served as controls. The percentages of apoptotic and
 20 necrotic BHK21 cells were determined by staining with annexin V-FITC and propidium iodide (PI) and flow cytometric analysis. Transfection with SINrep5 RNA vaccines caused a decline in the percentages of apoptotic cells 24 hr to 72 hr after electroporation (representative with SIN5-E7/Hsp70 70.3±3.6% for 24 hr, 49.3±4.2% for 48 hr, 18.0±3.1% for 72 hr, p<0.001, one-way ANOVA). Thus no statistically significant difference were observed when comparing the percentage of apoptotic cells transfected with
 25 various SINrep5 RNA vaccines. This experiment was repeated twice with similar results.

Figure 10. Enhanced MHC class I presentation of E7 to in bone marrow derived DCs pulsed with BHK21 cells transfected by SINrep5-E7/Hsp70 RNA as measured in CTL assays. BHK21 cells were electroporated with various self-replicating RNA constructs and co-cultured with BM-derived DCs. The DCs were then used as target cells for E7-specific CD8⁺ T effector cells (28). Cytolysis was
 30 determined by quantitative measurements of LDH as described herein. Self-replicating E7/Hsp70 RNA vaccines generated significantly higher lysis (at 3:1 and 9:1 E/T ratios) compared to the other RNA vaccines (p<0.001). CTL assays shown here are from one representative experiment of two performed.

Figure 11 is a schematic diagram of pSCA1-E7, pSCA1-Hsp70 and pSCA1-E7/Hsp70 constructs. The DNA-based SFV replicon vector, pSCA1, encodes the alphaviral replicon from Semliki

Forest virus. E7, Hsp70, and E7/Hsp70 DNA were cloned into the BamHI/SmaI sites of pSCA1. The HCMV IE promoter with subgenomic promoter are indicated with arrows.

Figures 12A and 12B show measurement of E7-specific CD8⁺ T cell precursors ("Tp") by intracellular cytokine staining and flow cytometric analysis. C57BL/6 mice were immunized with DNA-based self-replicating pSCA1-E7, pSCA1-Hsp70, pSCA1-E7/Hsp70, or pSCA1 without insert, using a gene gun and were boosted with the same regimen one week later. Fig. 12A): splenocytes from vaccinated mice were cultured with E7 peptide RAHYNIVTF (see above) overnight, stained for both CD8 and intracellular IFN- γ , and analyzed by flow cytometry. Mice vaccinated with E7/Hsp70 DNA generated the highest number of IFN γ -secreting CD8⁺ "double positive" T cells compared to other groups. (Fig. 12B). Flow cytometry was performed in the presence (solid columns) and absence (open columns) of RAHYNIVTF peptide. Results are expressed as the mean number of IFN- γ -secreting CD8⁺ T cells/ 3×10^5 splenocytes (\pm SEM). Results shown here are from one representative experiment of two performed.

Figures 13A and 13B show flow cytometry analysis of IFN- γ secreting or interleukin-4 (IL-4)-secreting E7-specific CD4⁺ T cells in mice vaccinated with various suicidal DNA vaccines. Mice were immunized as described in the description of Figure 12 (Fig. 13). Splenocytes from vaccinated mice were stimulated *in vitro* with the T-helper E7 peptide DSSEEEDEIDGPAGQAEPDRAHYNIVTFCKCDSTLRL (see above) overnight and were stained for both CD4 and intracellular IFN- γ and subjected to flow cytometry. No significant differences in the frequency of E7-specific IFN- γ -secreting CD4⁺ cells was observed in mice immunized with various recombinant DNA vaccines vs control. Fig. 13B: Splenocytes from vaccinated mice were cultured *in vitro* with the above E7 peptide overnight and stained for both CD4 and intracellular IL-4. The number of IL-4 secreting CD4⁺ T cells was analyzed by flow cytometry. No significant difference in the frequency of IL-4 secreting E7-specific CD4⁺ cells was observed among mice immunized with various recombinant DNA vaccines. This figure represents the mean value of three experiments \pm SEM.

Figure 14 shows E7-specific antibody responses in mice immunized with various pSCA1 suicide DNA vaccines. E7-specific antibodies were measured with ELISA using a serial dilution of serum. The results from the 1:100 dilution are presented, showing mean absorbance (OD₄₅₀) (\pm SEM). The results are from one representative experiment of two performed.

Figure 15 shows *in vivo* tumor protection against the growth of TC-1 tumors. Mice were immunized with various suicidal DNA vaccines and inoculated with tumors as described below. 80% of mice vaccinated with the pSCA1-E7/Hsp70 suicidal DNA vaccine remained tumor-free 70 days after TC-1 challenge. The results shown here are from one representative experiment of two performed.

Figure 16A and 16B show results of *in vivo* tumor therapy against pre-existing metastatic TC-1 tumor cells. Mice were first inoculated i.v. with tumors and then treated with various suicidal DNA vaccines. Fig. 16A shows that the pSCA1-E7/Hsp70 group had the lowest number of pulmonary metastatic nodules of all the groups (ANOVA, $P < 0.001$). Results are from one representative experiment of two performed. Fig. 16B shows representative lung tumors in each vaccinated group. Multiple grossly visible lung tumors were observed in unvaccinated control mice and mice vaccinated with Hsp70, E7, or vector DNA alone. No lung tumors were observed at this magnification in the pSCA1-E7/Hsp70 vaccinated group.

Figure 17. *In vivo* antibody depletion experiments determined the requirement for cells of certain lymphocyte subsets on the potency of the pSCA1-E7/Hsp70 suicidal DNA vaccine. Mice were inoculated i.v. with tumors and treated with the pSCA1-E7/Hsp70 suicidal DNA vaccine as described herein. CD4, CD8 or NK1.1 depletion were initiated one week prior to tumor challenge and lasted 40 days after tumor challenge. All naïve mice and mice depleted of CD8⁺ T cells grew tumors within 10 days

Figure 18A and 18B show schematic domain structure of the Flt3-ligand protein and FL-E7 fusion peptide (Fig. 18A) and the sequence of the FL-E7 construct, comprising the ECD of FL (Fig. 18B; SEQ ID NO:11 and 12). Residues 1-189 are FL-derived, residues 191-287 are E7-derived. The remaining residues (e.g., 288-302) are from the vector DNA.

Figure 19A and 19B. Intracellular cytokine staining with flow cytometry analysis to determine E7-specific CD8⁺ T cell precursors in C57BL/6 mice. Mice were immunized with FL DNA, E7 DNA (E7), FL-E7 DNA or FL mixed with E7 DNA (FL+E7) via gene gun, or received no vaccination. For vaccinated mice, 2 µg DNA /mouse was administered twice. Splenocytes were harvested 7 days after the last DNA vaccination. E7-specific CD8⁺ T cells. (Fig 19A) Splenocytes from vaccinated mice were cultured *in vitro* with E7 peptide (aa 49-57) overnight and were stained for both CD8 and intracellular IFN-γ. The number of IFN-γ secreting CD8⁺ T cell precursors in mice immunized with various recombinant DNA vaccines was analyzed by flow cytometry. Mice vaccinated with FL-E7 DNA generated the highest IFN-γ⁺ CD8⁺ double positive T cells compared to other groups. (Fig. 19B) The number of IFN-γ-producing E7-specific CD8⁺ T cells was determined using flow cytometry in the presence (solid columns) and absence (open columns) of E7 peptide (aa 49-57). Data are expressed as mean number of CD8⁺ IFN-γ⁺ cells/ 3×10^5 splenocytes ± SEM. The data from intracellular cytokine staining shown here are from one representative experiment of two performed.

Figure 20A and 20B. Flow cytometry analysis of IFN-γ secreting and IL-4-secreting E7-specific CD4⁺ cells in mice vaccinated with various recombinant DNA vaccines. Mice were immunized as

described in Figure Legend 2. (Fig. 20A) Splenocytes from vaccinated mice were cultured *in vitro* with E7 peptide (aa 30-67) overnight and were stained for both CD4 and intracellular IFN- γ . The number of IFN- γ -secreting CD4⁺ T cells was analyzed using flow cytometry. No significant difference in the frequency of E7-specific IFN- γ -secreting CD4⁺ cells was observed in mice immunized with various recombinant DNA vaccines. (Fig. 20B) Splenocytes from vaccinated mice were cultured *in vitro* with E7 peptide (aa 30-67) overnight and stained for both CD4 and intracellular IL-4. The percentage of IL-4 secreting CD4⁺ T cells was analyzed by flow cytometry. The IL-4 secreting activated mouse splenocytes (MiCK-2) from PharMingen were used as positive controls to assure the success of intracytoplasmic IL-4 staining for this study. The specificity of IL-4 staining was demonstrated by the absence of CD4⁺ IL-4⁺ T cells when the IL-4 antibody was omitted. No significant difference in the frequency of IL-4 secreting E7-specific CD4⁺ cells was observed in mice immunized with various recombinant DNA vaccines. The intracellular cytokine staining shown here are from one representative experiment of two performed.

Figure 21. *In vivo* tumor protection experiments against the growth of TC-1 tumors. Mice were immunized with FL DNA, E7 DNA, FL-E7 DNA or FL mixed with E7 DNA (FL+E7) via gene gun and boosted with the same regimen one week later. One week after the last vaccination, mice were challenged with 1×10^4 TC-1 cells/ mouse subcutaneously. Mice were monitored for evidence of tumor growth by palpation and inspection twice a week. 100% of mice receiving FL-E7 DNA vaccination remained tumor-free 60 days after TC-1 challenge. The data collected from the *in vivo* tumor protection experiments shown here are from one representative experiment of two performed.

Figure 22A and 22B. *In vivo* tumor treatment experiments against pre-existing metastatic TC-1 tumor cells. The mice were intravenously challenged with 1×10^4 cells/mouse TC-1 tumor cells in the tail vein on day 0. Three days after challenge with TC-1 tumor cells, mice received 2 μ g of FL DNA, E7 DNA, FL-E7 DNA, FL mixed with E7 (FL+E7), via gene gun or unvaccinated. One week later, these mice were boosted with the same regimen as the first vaccination. The mice were sacrificed on day 25. The FL-E7 group has (Fig. 22A) the least number of pulmonary metastatic nodules and (Fig. 22B) the lowest lung weight as compared with the other vaccinated groups (one-way ANOVA, $P < 0.001$). The data obtained from these *in vivo* treatment experiments are from one representative experiment of two performed.

Figure 23. Representative gross pictures of the lung tumors in each vaccinated group. Following *in vivo* tumor treatment experiments against pre-existing metastatic TC-1 tumor cells, there are multiple grossly visible lung tumors in unvaccinated control mice and mice vaccinated with FL, wild-type E7 DNA or FL mixed with E7 DNA. The lung tumors in FL-E7 vaccinated group cannot be seen at the magnification provided in this figure.

Figure 24. *In vivo* antibody depletion experiments to determine the effect of lymphocyte subset on the potency of FL-E7 DNA vaccine. Mice were immunized with 2 µg FL-E7 DNA via gene gun and boosted with 2 µg FL-E7 DNA one week later. One week after the last vaccination, mice were challenged with 1×10^4 TC-1 cells/ mouse subcutaneously. CD4, CD8 and NK1.1 depletions were initiated one week prior to tumor challenge and lasted 40 days after tumor challenge. Note: all of the unvaccinated mice and all of the mice depleted of CD8⁺ T cells grew tumors within 14 days after tumor challenge. The data of antibody depletion experiments shown here are from one representative experiment of two performed.

Figure 25. CTL assays to demonstrate enhanced presentation of E7 through the MHC class I pathway in cells transfected with FL-E7 DNA. 293 D^bK^b cells were transfected with various DNA vaccines with lipofectamine and collected 40-44 hr after transfection. Transfected 293 D^bK^b cells were used as target cells while D^b-restricted E7-specific CD8⁺ T cells were used as effector cells. CTL assays with various E/T ratios were performed. Note: The 293 D^bK^b cells transfected with FL-E7 DNA generated significantly higher percentages of specific lysis as compared to 293 D^bK^b cells transfected with other DNA vaccines. CTL assays shown here are from one representative experiment of two performed.

Figure 26. CTL assays to demonstrate enhanced MHC class I presentation of E7 in bone marrow derived dendritic cells pulsed with cell lysates containing chimeric FL-E7 protein. Bone marrow-derived DCs were pulsed with cell lysates from 293 D^bK^b cells transfected with various DNA vaccines in different concentration (50 µg/ml, 10 µg/ml, 2 µg/ml, and 0.4 µg/ml) for 16-24 hrs. D^b-restricted E7-specific CD8⁺ T cells were used as effector cells. CTL assays was performed at fixed E/T (9/1) ratio with 9×10^4 of E7-specific T cells mixed with 1×10^4 of prepared DCs in a final volume of 200 µl. Results of CTL assays were assessed using by quantitative measurements of LDH as described in the Materials and Methods. Note: DCs pulsed with lysates from cells transfected with FL-E7 DNA generated significantly higher percentages of specific lysis compared to DCs pulsed with lysates from cells transfected with other DNA vaccines. CTL assays shown here are from one representative experiment of two performed.

DESCRIPTION OF THE PREFERRED EMBODIMENTS

The invention provides compositions and methods for enhancing the immune responses, particularly cytotoxic T cell immune responses, induced by *ex vivo* or *in vivo* administration of chimeric polypeptides or, preferably, nucleic acid vaccines that encode these chimeric polypeptides. The preferred chimeric or fusion polypeptide comprises(1) at least one first polypeptide or peptide that, upon

introduction to cells of the host immune system, *in vitro* or *in vivo*, promotes (a) processing via the MHC class I pathway and/or (b) development or activity of APCs, primarily DCs, and (2) at least one second polypeptide or peptide that is an antigenic polypeptide or peptide in the host. As noted, in a preferred embodiment, the chimeric or fusion polypeptides are “indirectly” administered by administration of a nucleic acid that encodes the chimeric molecule; the nucleic acid construct, and thus the fusion protein, is expressed *in vivo*. The chimeric nucleic acids are administered in the form of DNA vaccines, either naked DNA or suicidal DNA, or a self-replicating RNA replicons.

The fusion protein comprises at least two domains or repeats thereof. The first domain again comprises a polypeptide that promotes (a) processing via the MHC class I pathway and/or (b) development or activity of APCs, and the second domain comprises a peptide or polypeptide, that includes one or several epitopes, derived from an antigen against which it is desired to induce an immune response.

For convenience, a polypeptide or peptide that promotes processing via the MHC class I pathway is abbreviated herein as “MHC-I-PP.” A polypeptide or peptide that promotes development or activity of APCs, preferably DC’s, is abbreviated DC-PP.

The exemplary MHC-I-PP protein described herein is Hsp70. However, it is understood that any protein, or functional fragment or variant thereof, that has this activity can be used in the invention. A preferred fragment is a C-terminal domain (“CD”) of Hsp70, which is designated “Hsp70_{CD}”. One Hsp70_{CD} spans from about residue 312 to the C terminus of Hsp70 (SEQ ID NO:4). A preferred shorter polypeptide spans from about residue 517 to the C-terminus of SEQ ID NO:4. Shorter peptides from that sequence that have the ability to promote protein processing via the MHC-I class I pathway are also included, and may be defined by routine experimentation.

The second type of domain of the chimeric molecule comprises an antigenic peptide, which can be derived from a pathogen, a cancer cell, or any source to which induction, enhancement or suppression of an immune response is desired. In a preferred embodiment, the peptide comprises at least one MHC class I-binding peptide epitope that helps stimulate CD8+ CTLs and is recognized by such cells and their precursors.

The order in which the two (or more) component polypeptides of the fusion protein are arranged, and therefore, the order of the encoding nucleic acid fragments in the nucleic acid vector, can be altered without affecting immunogenicity of the fusion polypeptides proteins and the utility of the composition. For example, the Hsp70-encoding (or FL -encoding) DNA sequences may be located 5’ or 3’ to the target antigen-encoding sequences. In one embodiment, these polypeptide-encoding nucleic acid domains are in-frame so that the DNA construct encodes a recombinant fusion polypeptide in which the antigen is located N- terminal to the Hsp70 or FL derived polypeptide.

The vaccines of the present invention include, the antigenic epitope itself and an MHC-I-PP such as Hsp70 or its active domain (CD), or DC-PP intercellular spreading protein such as FL. In addition to the specific antigens and vectors employed in the Examples, the present invention is intended to encompass a vector such as naked DNA, naked RNA, self replicating RNA replicons and viruses including vaccinia, adenoviruses, adeno-associated virus (AAV), lentiviruses and RNA alphaviruses.

In addition to the MHC-I-PP and/or DC-PP, the vaccine construct of the present invention optionally, may also include

- (a) an additional antigen targeting or processing signal such as proteins that promote intercellular transport, *e.g.*, VP22 protein from herpes simplex virus and related herpes viruses; an endoplasmic reticulum chaperone polypeptide such as calreticulin, ER60, GRP94 or gp96, well-characterized ER chaperone polypeptide that representatives of the HSP90 family of stress-induced proteins (Argon (1999) *Semin. Cell Dev. Biol.* 10:495-505; Sastry (1999) *J. Biol. Chem.* 274:12023-12035; Nicchitta (1998) *Curr. Opin. Immunol.* 10:103-109; U.S. Patent 5,981,706); cytoplasmic translocation polypeptide domains of pathogen toxins, such as domain II of *Pseudomonas* exotoxin ETA (ETAdII) or of similar toxins from *Diphtheria*, *Clostridium*, *Botulinum*, *Bacillus*, *Yersinia*, *Vibrio cholerae*, or *Bordetella pertussis*; or active fragments or domains of any of the foregoing polypeptides.
- (b) an immunostimulatory cytokine, preferably those that target APCs, preferably DC's, such as granulocyte macrophage colony stimulating factor (GM-CSF), or active fragments or domains thereof; and
- (c) a costimulatory signal, such as a B7 family protein, including B7-DC (see commonly assigned U.S. patent application Serial No. 09/794,210), B7.1, B7.2, soluble CD40, *etc.*).

(For description of some of the foregoing, see, for example, commonly owned International patent applications PCT/US01/23966, PCT/US01/24134, PCTUS/00/41422))

In the methods of the invention, the chimeric polypeptide or nucleic acid that encodes it are employed to induce or enhance immune responses. In one embodiment, the compositions of the invention synergistically enhance immune responses and antitumor effects through both immunological and anti-angiogenic mechanisms.

The experiments described herein demonstrate that the methods of the invention can enhance a cellular immune response, particularly, tumor-destructive CTL reactivity, induced by a DNA vaccine encoding an epitope of a human pathogen. Human HPV-16 E7 was used as a model antigen for vaccine development because human papillomaviruses (HPVs), particularly HPV-16, are associated with most human cervical cancers. The oncogenic HPV protein E7 is important in the induction and maintenance of cellular transformation and co-expressed in most HPV-containing cervical cancers and their precursor

lesions. Therefore, cancer vaccines, such as the compositions of the invention, that target E7 can be used to control of HPV-associated neoplasms (Wu (1994) *Curr. Opin. Immunol.* 6:746-754).

Unless defined otherwise, all technical and scientific terms used herein have the meaning commonly understood by a person skilled in the art of this invention. As used herein, the following terms have the meanings ascribed to them unless specified otherwise.

The term "antigen" or "immunogen" as used herein refers to a compound or composition comprising a peptide, polypeptide or protein which is "antigenic" or "immunogenic" when administered (or expressed *in vivo* by an administered nucleic acid, *e.g.*, a DNA vaccine) in an appropriate amount (an "immunogenically effective amount"), *i.e.*, capable of inducing, eliciting, augmenting or boosting a cellular and/or humoral immune response either alone or in combination or linked or fused to another substance (which can be administered at once or over several intervals). An immunogenic composition can comprise an antigenic peptide of at least about 5 amino acids, a peptide of 10 amino acids in length, a polypeptide fragment of 15 amino acids in length, 20 amino acids in length or longer. Smaller immunogens may require presence of a "carrier" polypeptide *e.g.*, as a fusion protein, aggregate, conjugate or mixture, preferably linked (chemically or otherwise) to the immunogen. The immunogen can be recombinantly expressed from a vaccine vector, which can be naked DNA comprising the immunogen's coding sequence operably linked to a promoter, *e.g.*, an expression cassette as described herein. The immunogen includes one or more antigenic determinants or epitopes which may vary in size from about 3 to about 15 amino acids.

The term "epitope" as used herein refers to an antigenic determinant or antigenic site that interacts with an antibody or a T cell receptor (TCR), *e.g.*, the MHC class I-binding peptide compositions (or expressed products of the nucleic acid compositions of the invention) used in the methods of the invention. An "antigen" is a molecule or chemical structure that either induces an immune response or is specifically recognized or bound by the product or mediator of an immune response, such as an antibody or a CTL. The specific conformational or stereochemical "domain" to which an antibody or a TCR bind is an "antigenic determinant" or "epitope." TCRs bind to peptide epitopes which are physically associated with a third molecule, a major histocompatibility complex (MHC) class I or class II protein.

The term "recombinant" refers to (1) a nucleic acid or polynucleotide synthesized or otherwise manipulated *in vitro*, (2) methods of using recombinant DNA technology to produce gene products in cells or other biological systems, or (3) a polypeptide encoded by a recombinant nucleic acid. For example, the FL-encoding nucleic acid or polypeptide, the nucleic acid encoding an MHC class I-binding peptide epitope (antigen) or the peptide itself can be recombinant. "Recombinant means" includes ligation of nucleic acids having various coding regions or domains or promoter sequences from

different sources into a single unit in the form of an expression cassette or vector for expression of the coding sequences in the vectors resulting in production of the encoded polypeptide.

The term "self-replicating RNA replicon" refers to a construct based on an RNA viruses, such as alphavirus genome RNAs (*e.g.*, Sindbis virus, Semliki Forest virus, *etc.*), that have been engineered to allow expression of heterologous RNAs and proteins. These recombinant vectors are self-replicating ("replicons") which can be introduced into cells as naked RNA or DNA, as described in detail, below. In a preferred embodiment, the self-replicating RNA replicon comprises a Sindbis virus self-replicating RNA vector SINrep5, which is described in detail in U.S. Patent No. 5,217,879.

SEQUENCES OF POLYPEPTIDES AND NUCLEIC ACIDS

The section that follows lists the sequences of the MHC-I-PP and DC-PP polypeptides alone or in fusion with E7 antigen, the nucleic acids encoding some of these peptides and nucleic acids of the vectors into which the sequences encoding these polypeptides are cloned.

HPV-E7 (nucleic acid is SEQ ID NO:1; amino acids are SEQ ID NO:2)

```

1/1          31/11
atg cat gga gat aca cct aca ttg cat gaa tat atg tta gat ttg caa cca gag aca act
Met his gly asp thr pro thr leu his glu tyr met leu asp leu gln pro glu thr thr
61/21
gat ctc tac tgt tat gag caa tta aat gac agc tca gag gag gag gat gaa ata gat ggt
asp leu tyr cys tyr glu gln leu asn asp ser ser glu glu glu asp glu ile asp gly
121/41
cca gct gga caa gca gaa ccg gac aga gcc cat tac aat att gta acc ttt tgt tgc aag
pro ala gly gln ala glu pro asp arg ala his tyr asn ile val thr phe cys cys lys
181/61
tgt gac tct acg ctt cgg ttg tgc gta caa agc aca cac gta gac att cgt act ttg gaa
cys asp ser thr leu arg leu cys val gln ser thr his val asp ile arg thr leu glu
241/81
gac ctg tta atg ggc aca cta gga att gtg tgc ccc atc tgt tct cag gat aag ctt
asp leu leu met gly thr leu gly ile val cys pro ile cys ser gln asp lys leu

```

GENBANK Accession No. AAD33353 ("native" sequence). After cloning the last 9 nucleotides (3 amino acids were modified, as shown above underscored and/or bold).

The original GENBANK sequence has a Lys-Pro after the Gln in position 96 (encoded by aaa/cca/taa, rather than Asp-Lys-Leu (encoded by gat/aag/ctt)).

Hsp70 from *M. tuberculosis*

(nucleic acid is SEQ ID NO:3; amino acids are SEQ ID NO:4)

```

1/1          31/11
atg gct cgt gcg gtc ggg atc gac ctc ggg acc acc aac tcc gtc gtc tcg gtt ctg gaa
Met ala arg ala val gly ile asp leu gly thr thr asn ser val val ser val leu glu
61/21
ggg ggc gac ccg gtc gtc gtc gcc aac tcc gag ggc tcc agg acc acc ccg tca att gtc
gly gly asp pro val val val ala asn ser glu gly ser arg thr thr pro ser ile val
121/41
gcg ttc gcc cgc aac ggt gag gtg ctg gtc gcc cag ccc gcc aag aac cag gca gtg acc
ala phe ala arg asn gly glu val leu val gly gln pro ala lys asn gln ala val thr

```

	181/61		211/71	
	aac gtc gat cgc acc gtg cgc tcg gtc aag	cga cac atg ggc agc gac tgg tcc ata gag		
	asn val asp arg thr val arg ser val lys	arg his met gly ser asp trp ser ile glu		
5	241/81		271/91	
	att gac ggc aag aaa tac acc gcg ccg gag	atc agc gcc cgc att ctg atg aag ctg aag		
	ile asp gly lys lys tyr thr ala pro glu	ile ser ala arg ile leu met lys leu lys		
	301/101		331/111	
	cgc gac gcc gag gcc tac ctc ggt gag gac	att acc gac gcg gtt atc acg acg ccc gcc		
10	361/121		391/131	
	arg asp ala glu ala tyr leu gly glu asp	ile thr asp ala val ile thr thr pro ala		
	tac ttc aat gac gcc cag cgt cag gcc acc	aag gac gcc ggc cag atc gcc ggc ctc aac		
	tyr phe asn asp ala gln arg gln ala thr	lys asp ala gly gln ile ala gly leu asn		
	421/141		451/151	
15	481/161		511/171	
	gtg ctg cgg atc gtc aac gag ccg acc gcg	gcc gcg ctg gcc tac ggc ctc gac aag ggc		
	val leu arg ile val asn glu pro thr ala	ala ala leu ala tyr gly leu asp lys gly		
	gag aag gag cag cga atc ctg gtc ttc gac	ttg ggt ggt ggc act ttc gac gtt tcc ctg		
	glu lys glu gln arg ile leu val phe asp	leu gly gly gly thr phe asp val ser leu		
	541/181		571/191	
20	601/201		631/211	
	ctg gag atc ggc gag ggt gtg gtt gag gtc	cgt gcc act tcg ggt gac aac cac ctc ggc		
	leu glu ile gly glu gly val val glu val	arg ala thr ser gly asp asn his leu gly		
	ggc gac gac tgg gac cag ccg gtc gtc gat	tgg ctg gtc gac aag ttc aag ggc acc agc		
25	661/221		691/231	
	gly asp asp trp asp gln arg val val asp	trp leu val asp lys phe lys gly thr ser		
	ggc atc gat ctg acc aag gac aag atg gcg	atg cag cgg ctg cgg gaa gcc gcc gag aag		
	gly ile asp leu thr lys asp lys met ala	met gln arg leu arg glu ala ala glu lys		
	721/241		751/251	
30	781/261		811/271	
	gca aag atc gag ctg agt tcg agt cag tcc	acc tcg atc aac ctg ccc tac atc acc gtc		
	ala lys ile glu leu ser ser ser gln ser	thr ser ile asn leu pro tyr ile thr val		
	gac gcc gac aag aac ccg ttg ttc tta gac	gag cag ctg acc cgc gcg gag ttc caa cgg		
	asp ala asp lys asn pro leu phe leu asp	glu gln leu thr arg ala glu phe gln arg		
	841/281		871/291	
35	901/301		931/311	
	atc act cag gac ctg ctg gac cgc act cgc	aag ccg ttc cag tcg gtg atc gct gac acc		
	ile thr gln asp leu leu asp arg thr arg	lys pro phe gln ser val ile ala asp thr		
	ggc att tcg gtg tcg gag atc gat cac gtt	gtg ctc gtg ggt ggt tcg acc cgg atg ccc		
40	961/321		991/331	
	gly ile ser val ser glu ile asp his val	val leu val gly gly ser thr arg met pro		
	gcg gtg acc gat ctg gtc aag gaa ctc acc	ggc ggc aag gaa ccc aac aag ggc gtc aac		
	ala val thr asp leu val lys glu leu thr	gly gly lys glu pro asn lys gly val asn		
	1021/341		1051/351	
45	1081/361		1111/371	
	ccc gat gag gtt gtc gcg gtg gga gcc gct	ctg cag gcc ggc gtc ctc aag ggc gag gtg		
	pro asp glu val val ala val gly ala ala	leu gln ala gly val leu lys gly glu val		
	aaa gac gtt ctg ctg ctt gat gtt acc ccg	ctg agc ctg ggt atc gag acc aag ggc ggc		
	lys asp val leu leu leu asp val thr pro	leu ser leu gly ile glu thr lys gly gly		
	1141/381		1171/391	
50	1201/401		1231/411	
	gtg atg acc agg ctc atc gag cgc aac acc	acg atc ccc acc aag ccg tcg gag act ttc		
	val met thr arg leu ile glu arg asn thr	thr ile pro thr lys arg ser glu thr phe		
	acc acc gcc gac gac aac caa ccg tcg gtg	cag atc cag gtc tat cag ggc gag cgt gag		
55	1261/421		1291/431	
	thr thr ala asp asp asn gln pro ser val	gln ile gln val tyr gln gly glu arg glu		
	atc gcc gcg cac aac aag ttg ctc ggg tcc	ttc gag ctg acc ggc atc ccg ccg gcg ccg		
	ile ala ala his asn lys leu leu gly ser	phe glu leu thr gly ile pro pro ala pro		
	1321/441		1351/451	
60	1381/461		1411/471	
	cgg ggg att ccg cag atc gag gtc act ttc	gac atc gac gcc aac ggc att gtg cac gtc		
	arg gly ile pro gln ile glu val thr phe	asp ile asp ala asn gly ile val his val		
	acc gcc aag gac aag ggc acc ggc aag gag	aac acg atc cga atc cag gaa ggc tcg ggc		
	thr ala lys asp lys gly thr gly lys glu	asn thr ile arg ile gln glu gly ser gly		
	1441/481		1471/491	
65	1501/501		1531/511	
	ctg tcc aag gaa gac att gac cgc atg atc	aag gac gcc gaa gcg cac gcc gag gag gat		
	leu ser lys glu asp ile asp arg met ile	lys asp ala glu ala his ala glu glu asp		
	cgc aag cgt cgc gag gag gcc gat gtt cgt	aat caa gcc gag aca ttg gtc tac cag acg		
70	1561/521		1591/531	
	arg lys arg arg glu glu ala asp val arg	asn gln ala glu thr leu val tyr gln thr		
	gag aag ttc gtc aaa gaa cag cgt gag gcc	gag ggt ggt tcg aag gta cct gaa gac acg		
	glu lys phe val lys glu gln arg glu ala	glu gly gly ser lys val pro glu asp thr		
	1621/541		1651/551	

540

ctg aac aag gtt gat gcc gcg gtg gcg gaa gcg aag gcg gca ctt ggc gga tcg gat att
 leu asn lys val asp ala ala val ala glu ala lys ala ala leu gly gly ser asp ile 560
 1681/561 1711/571
 5 tcg gcc atc aag tcg gcg atg gag aag ctg ggc cag gag tcg cag gct ctg ggg caa gcg
 ser ala ile lys ser ala met glu lys leu gly gln glu ser gln ala leu gly gln ala 580
 1741/581 1771/591
 atc tac gaa gca gct cag gct gcg tca cag gcc act ggc gct gcc cac ccc ggc tcg gct
 ile tyr glu ala ala gln ala ala ser gln ala thr gly ala ala his pro gly ser ala
 1801/601
 10 gat gaa AGC
 asp glu ser

GENBANK Z95324 AL123456; encoded by nucleotides 10633-12510 of *Mycobacterium tuberculosis*
 genome). As a result of cloning, this has been modified from the original GENBANK sequence which
 15 had at its 3' end:

ggc gag ccg ggc ggt gcc cac ccc ggc tcg gct gat gac gtt gtg gac gcg gag gtg
 gtc gac gac ggc cgg gag gcc aag (SEQ ID NO:5)

which was replaced in the cloned version used herein, by tcg gct gat gaa agc (SEQ ID NO:6)
 which is bold and underlined above.

E7-Hsp70 Fusion

(nucleic acid is SEQ ID NO:7; amino acids are SEQ ID NO:8)

E7 coding sequence is capitalized and underscored.

1/1 31/11
 25 ATG CAT GGA GAT ACA CCT ACA TTG CAT GAA TAT ATG TTA GAT TTG CAA CCA GAG ACA ACT
 Met his gly asp thr pro thr leu his glu tyr met leu asp leu gln pro glu thr thr
 61/21 91/31
GAT CTC TAC TGT TAT GAG CAA TTA AAT GAC AGC TCA GAG GAG GAG GAT GAA ATA GAT GGT
 asp leu tyr cys tyr glu gln leu asn asp ser ser glu glu glu asp glu ile asp gly
 121/41 151/51
 30 CCA GCT GGA CAA GCA GAA CCG GAC AGA GCC CAT TAC AAT ATT GTA ACC TTT TGT TGC AAG
 pro ala gly gln ala glu pro asp arg ala his tyr asn ile val thr phe cys cys lys
 181/61 211/71
TGT GAC TCT ACG CTT CGG TTG TGC GTA CAA AGC ACA CAC GTA GAC ATT CGT ACT TTG GAA
 35 cys asp ser thr leu arg leu cys val gln ser thr his val asp ile arg thr leu glu
 241/81 271/91
GAC CTG TTA ATG GGC ACA CTA GGA ATT GTG TGC CCC ATC TGT TCT CAA GGA TCC atg gct
 asp leu leu met gly thr leu gly ile val cys pro ile cys ser gln gly ser met ala
 301/101 331/111
 40 cgt gcg gtc ggg atc gac ctc ggg acc acc tac tcc gtc gtc tcg gtt ctg gaa ggt ggc
 arg ala val gly ile asp leu gly thr thr asn ser val val ser val leu glu gly gly
 361/121 391/131
gac ccg gtc gtc gtc gcc aac tcc gag ggc tcc agg acc acc ccg tca att gtc gcg ttc
 asp pro val val val ala asn ser glu gly ser arg thr thr pro ser ile val ala phe
 421/141 451/151
 45 gcc cgc aac ggt gag gtg ctg gtc ggc cag ccc gcc aag aac cag gca gtg acc aac gtc
 ala arg asn gly glu val leu val gly gln pro ala lys asn gln ala val thr asn val
 481/161 511/171
gat cgc acc gtg cgc tcg gtc aag cga cac atg gcc agc gac tgg tcc ata gag att gac
 50 asp arg thr val arg ser val lys arg his met gly ser asp trp ser ile glu ile asp
 541/181 571/191
ggc aag aaa tac acc gcg ccg gag atc agc gcc cgc att ctg atg aag ctg aag cgc gac
 gly lys lys tyr thr ala pro glu ile ser ala arg ile leu met lys leu lys arg asp
 601/201 631/211
 55 gcc gag gcc tac ctc ggt gag gac att acc gac gcg gtt atc acg acg ccc gcc tac ttc
 ala glu ala tyr leu gly glu asp ile thr asp ala val ile thr thr pro ala tyr phe
 661/221 691/231
aat gac gcc cag cgt cag gcc acc aag gac gcc gcc cag atc gcc ggc ctc aac gtg ctg
 721/241 751/251
 60 cgg atc gtc aac gag ccg acc gcg gcc gcg ctg gcc tac ggc ctc gac aag ggc gag aag
 arg ile val asn glu pro thr ala ala ala leu ala tyr gly leu asp lys gly glu lys
 781/261 811/271

gag cag cga atc ctg gtc ttc gac ttg ggt ggt ggc act ttc gac gtt tcc ctg ctg gag
 glu gln arg ile leu val phe asp leu gly gly thr phe asp val ser leu leu glu
 841/281 871/291
 5 atc ggc gag ggt gtg gtt gag gtc cgt gcc act tcg ggt gac aac cac ctc ggc ggc gac
 ile gly glu gly val val glu val arg ala thr ser gly asp asn his leu gly gly asp
 901/301 931/311
 gag tgg gac cag cgg gtc gtc gat tgg ctg gtg gac aag ttc aag ggc acc agc ggc atc
 asp trp asp gln arg val val asp trp leu val asp lys phe lys gly thr ser gly ile
 961/321 991/331
 10 gat ctg acc aag gac aag atg gcg atg cag cgg ctg cgg gaa gcc gcc gag aag gca aag
 asp leu thr lys asp lys met ala met gln arg leu arg glu ala ala glu lys ala lys
 1021/341 1051/351
 15 atc gag ctg agt tcg agt cag tcc acc tcg atc aac ctg ccc tac atc acc gtc gac gcc
 ile glu leu ser ser ser gln ser thr ser ile asn leu pro tyr ile thr val asp ala
 1081/361 1111/371
 gag aag aac ccg ttg ttc tta gac gag cag ctg acc cgc gcg gag ttc caa cgg atc act
 asp lys asn pro leu phe leu asp glu gln leu thr arg ala glu phe gln arg ile thr
 1141/381 1171/391
 20 cag gac ctg ctg gac cgc act cgc aag ccg ttc cag tcg gtg atc gct gac acc ggc att
 gln asp leu leu asp arg thr arg lys pro phe gln ser val ile ala asp thr gly ile
 1201/401 1231/411
 tcg gtg tcg gag atc gat cac gtt gtg ctc gtg ggt ggt tcg acc cgg atg ccc gcg gtg
 ser val ser glu ile asp his val val leu val gly gly ser thr arg met pro ala val
 1261/421 1291/431
 25 acc gat ctg gtc aag gaa ctc acc ggc ggc aag gaa ccc aac aag ggc gtc aac ccc gat
 thr asp leu val lys glu leu thr gly gly lys glu pro asn lys gly val asn pro asp
 1321/441 1351/451
 gag gtt gtc gcg gtg gga gcc gct ctg cag gcc ggc gtc ctc aag ggc gag gtg aaa gac
 glu val val ala val gly ala ala leu gln ala gly val leu lys gly glu val lys asp
 1381/461 1411/471
 30 gtt ctg ctg ctt gat gtt acc ccg ctg agc ctg ggt atc gag acc aag ggc ggc gtg atg
 val leu leu leu asp val thr pro leu ser leu gly ile glu thr lys gly gly val met
 1441/481 1471/491
 35 acc agg ctc atc gag cgc aac acc acg atc ccc acc aag cgg tcg gag act ttc acc acc
 thr arg leu ile glu arg asn thr thr ile pro thr lys arg ser glu thr phe thr thr
 1501/501 1531/511
 gcc gac gac aac caa ccg tcg gtg cag atc cag gtc tat cag ggg gag cgt gag atc gcc
 ala asp asp asn gln pro ser val gln ile gln val tyr gln gly glu arg glu ile ala
 1561/521 1591/531
 40 gcg cac aac aag ttg ctc ggg tcc ttc gag ctg acc ggc atc ccg ccg gcg ccg cgg ggg
 ala his asn lys leu leu gly ser phe glu leu thr gly ile pro pro ala pro arg gly
 1621/541 1651/551
 att ccg cag atc gag gtc act ttc gac atc gac gcc aac ggc att gtg cac gtc acc gcc
 ile pro gln ile glu val thr phe asp ile asp ala asn gly ile val his val thr ala
 1681/561 1711/571
 45 aag gac aag ggc acc ggc aag gag aac acg atc cga atc cag gaa ggc tcg ggc ctg tcc
 lys asp lys gly thr gly lys glu asn thr ile arg ile gln glu gly ser gly leu ser
 1741/581 1771/591
 50 aag gaa gac att gac cgc atg atc aag gac gcc gaa gcg cac gcc gag gag gat cgc aag
 lys glu asp ile asp arg met ile lys asp ala glu ala his ala glu glu asp arg lys
 1801/601 1831/611
 cgt cgc gag gag gcc gat gtt cgt aat caa gcc gag aca ttg gtc tac cag acg gag aag
 arg arg glu glu ala asp val arg asn gln ala glu thr leu val tyr gln thr glu lys
 1861/621 1891/631
 55 ttc gtc aaa gaa cag cgt gag gcc gag ggt ggt tcg aag gta cct gaa gac acg ctg aac
 phe val lys glu gln arg glu ala glu gly gly ser lys val pro glu asp thr leu asn
 1921/641 1951/651
 60 aag gtt gat gcc gcg gtg gcg gaa gcg aag cgc gca ctt ggc gga tcg gat att tcg gcc
 lys val asp ala ala val ala glu ala lys ala ala leu gly gly ser asp ile ser ala
 1981/661 2011/671
 atc aag tcg gcg atg gag aag ctg ggc cag gag tcg cag gct ctg ggg caa gcg atc tac
 ile lys ser ala met glu lys leu gly gln glu ser gln ala leu gly gln ala ile tyr
 2041/681 2071/691
 65 gaa gca gct cag gct gcg tca cag gcc act ggc gct gcc cac ccc ggc tcg gct gat gaA
 glu ala ala gln ala ala ser gln ala thr gly ala ala his pro gly ser ala asp glu
 2101/701
 AGC a
 ser

Flt3 Ligand (FL) extracellular domain

(nucleic acid is SEQ ID NO:9; amino acids are SEQ ID NO:10)

```

1/1          31/11
5  atg aca gtg ctg gcg cca gcc tgg agc cca aat tcc tcc ctg ttg ctg ctg ttg ctg ctg
   Met thr val leu ala pro ala trp ser pro asn ser ser leu leu leu leu leu leu
61/21      91/31
   ctg agt cct tgc ctg cgg ggg aca cct gac tgt tac ttc agc cac agt ccc atc tcc tcc
   leu ser pro cys leu arg gly thr pro asp cys tyr phe ser his ser pro ile ser ser
121/41     151/51
10  aac ttc aaa gtg aag ttt aga gag ttg act gac cac ctg ctt aaa gat tac cca gtc act
   asn phe lys val lys phe arg glu leu thr asp his leu leu lys asp tyr pro val thr
181/61     211/71
   gtg gcc gtc aat ctt cag gac gag aag cac tgc aag gcc ttg tgg agc ctc ttc cta gcc
   val ala val asn leu gln asp glu lys his cys lys ala leu trp ser leu phe leu ala
15 241/81     271/91
   cag cgc tgg ata gag caa ctg aag act gtg gca ggg tct aag atg caa acg ctt ctg gag
   gln arg trp ile glu gln leu lys thr val ala gly ser lys met gln thr leu leu glu
301/101    331/111
20  gac gtc aac acc gag ata cat ttt gtc acc tca tgt acc ttc cag ccc cta cca gaa tgt
   asp val asn thr glu ile his phe val thr ser cys thr phe gln pro leu pro glu cys
361/121    391/131
   ctg cga ttc gtc cag acc aac atc tcc cac ctg ctg aag gac acc tgc aca cag ctg ctt
   leu arg phe val gln thr asn ile ser his leu leu lys asp thr cys thr gln leu leu
421/141    451/151
25  gct ctg aag ccc tgt atc ggg aag gcc tgc cag aat ttc tct cgg tgc ctg gag gtg cag
   ala leu lys pro cys ile gly lys ala cys gln asn phe ser arg cys leu glu val gln
481/161    511/171
   tgc cag ccg gac tcc tcc acc ctg ctg ccc cca agg agt ccc ata gcc cta gaa gcc acg
   cys gln pro asp ser ser thr leu leu pro pro arg ser pro ile ala leu glu ala thr
30 541/181
   gag ctc cca gag cct cgg ccc agg cag
   glu leu pro glu pro arg pro arg gln

```

FL-E7 Fusion Polypeptide

35 (nucleic acid is SEQ ID NO:11; amino acids are SEQ ID NO:12)

N-terminal sequence is FL, followed by E7 (underscored, cap)

```

1/1          31/11
40  atg aca gtg ctg gcg cca gcc tgg agc cca aat tcc tcc ctg ttg ctg ctg ttg ctg ctg
   Met thr val leu ala pro ala trp ser pro asn ser ser leu leu leu leu leu leu
61/21      91/31
   ctg agt cct tgc ctg cgg ggg aca cct gac tgt tac ttc agc cac agt ccc atc tcc tcc
   leu ser pro cys leu arg gly thr pro asp cys tyr phe ser his ser pro ile ser ser
121/41     151/51
45  aac ttc aaa gtg aag ttt aga gag ttg act gac cac ctg ctt aaa gat tac cca gtc act
   asn phe lys val lys phe arg glu leu thr asp his leu leu lys asp tyr pro val thr
181/61     211/71
   gtg gcc gtc aat ctt cag gac gag aag cac tgc aag gcc ttg tgg agc ctc ttc cta gcc
   val ala val asn leu gln asp glu lys his cys lys ala leu trp ser leu phe leu ala
50 241/81     271/91
   cag cgc tgg ata gag caa ctg aag act gtg gca ggg tct aag atg caa acg ctt ctg gag
   gln arg trp ile glu gln leu lys thr val ala gly ser lys met gln thr leu leu glu
301/101    331/111
55  gac gtc aac acc gag ata cat ttt gtc acc tca tgt acc ttc cag ccc cta cca gaa tgt
   asp val asn thr glu ile his phe val thr ser cys thr phe gln pro leu pro glu cys
361/121    391/131
   ctg cga ttc gtc cag acc aac atc tcc cac ctg ctg aag gac acc tgc aca cag ctg ctt
   leu arg phe val gln thr asn ile ser his leu leu lys asp thr cys thr gln leu leu
421/141    451/151
60  gct ctg aag ccc tgt atc ggg aag gcc tgc cag aat ttc tct cgg tgc ctg gag gtg cag
   ala leu lys pro cys ile gly lys ala cys gln asn phe ser arg cys leu glu val gln
481/161    511/171
   tgc cag ccg gac tcc tcc acc ctg ctg ccc cca agg agt ccc ata gcc cta gaa gcc acg
   cys gln pro asp ser ser thr leu leu pro pro arg ser pro ile ala leu glu ala thr
541/181    571/191
65  gag ctc cca gag cct cgg ccc agg cag gaa ttc ATG CAT GGA GAT ACA CCT ACA TTG CAT
   glu leu pro glu pro arg pro arg gln glu phe met his gly asp thr pro thr leu his

```

601/201 631/211
 GAA TAT ATG TTA GAT TTG CAA CCA GAG ACA ACT GAT CTC TAC TGT TAT GAG CAA TTA AAT
 glu tyr met leu asp leu gln pro glu thr thr asp leu tyr cys tyr glu gln leu asn
 661/221 691/231
 5 GAC AGC TCA GAG GAG GAG GAT GAA ATA GAT GGT CCA GCT GGA CAA GCA GAA CCG GAC AGA
 asp ser ser glu glu glu asp glu ile asp gly pro ala gly gln ala glu pro asp arg
 721/241 751/251
 10 GCC CAT TAC AAT ATT GTA ACC TTT TGT TGC AAG TGT GAC TCT ACG CTT CGG TTG TGC GTA
 ala his tyr asn ile val thr phe cys cys lys cys asp ser thr leu arg leu cys val
 781/261 811/271
 CAA AGC ACA CAC GTA GAC ATT CGT ACT TTG GAA GAC CTG TTA ATG GGC ACA CTA GGA ATT
 gln ser thr his val asp ile arg thr leu glu asp leu leu met gly thr leu gly ile
 841/281
 15 GTG TGC CCC ATC TGT TCT CAA GGA TCC
 val cys pro ile cys ser gln gly ser

FL-E7-GFP Fusion Polypeptide

(nucleic acid is SEQ ID NO:13; amino acids are SEQ ID NO:14)

N-terminal sequence is FL, followed by E7 (underscored, cap) followed by GFP (italic)

20 1/1 31/11
 atg aca gtg ctg gcg cca gcc tgg agc cca aat tcc tcc ctg ttg ctg ctg ttg ctg ctg
 Met thr val leu ala pro ala trp ser pro asn ser ser leu leu leu leu leu leu
 61/21 91/31
 25 ctg agt cct tgc ctg cgg ggg aca cct gac tgt tac ttc agc cac agt ccc atc tcc tcc
 leu ser pro cys leu arg gly thr pro asp cys tyr phe ser his ser pro ile ser ser
 121/41 151/51
 30 aac ttc aaa gtg aag ttt aga gag ttg act gac cac ctg ctt aaa gat tac cca gtc act
 asn phe lys val lys phe arg glu leu thr asp his leu leu lys asp tyr pro val thr
 181/61 211/71
 gtg gcc gtc aat ctt cag gac gag aag cac tgc aag gcc ttg tgg agc ctc ttc cta gcc
 val ala val asn leu gln asp glu lys his cys lys ala leu trp ser leu phe leu ala
 241/81 271/91
 35 cag cgc tgg ata gag caa ctg aag act gtg gca ggg tct aag atg caa acg ctt ctg gag
 gln arg trp ile glu gln leu lys thr val ala gly ser lys met gln thr leu leu glu
 301/101 331/111
 gac gtc aac acc gag ata cat ttt gtc acc tca tgt acc ttc cag ccc cta cca gaa tgt
 asp val asn thr glu ile his phe val thr ser cys thr phe gln pro leu pro glu cys
 361/121 391/131
 40 ctg cga ttc gtc cag acc aac atc tcc cac ctg ctg aag gac acc tgc aca cag ctg ctt
 leu arg phe val gln thr asn ile ser his leu leu lys asp thr cys thr gln leu leu
 421/141 451/151
 45 gct ctg aag ccc tgt atc ggg aag gcc tgc cag aat ttc tct cgg tgc ctg gag gtg cag
 ala leu lys pro cys ile gly lys ala cys gln asn phe ser arg cys leu glu val gln
 481/161 511/171
 tgc cag ccg gac tcc tcc acc ctg ctg ccc cca agg agt ccc ata gcc cta gaa gcc acg
 cys gln pro asp ser ser thr leu leu pro pro arg ser pro ile ala leu glu ala thr
 541/181 571/191
 50 gag ctg cca gag cct cgg ccc agg cag gaa ttc ATG CAT GGA GAT ACA CCT ACA TTG CAT
 glu leu pro glu pro arg pro arg gln glu phe met his gly asp thr pro thr leu his
 601/201 631/211
 GAA TAT ATG TTA GAT TTG CAA CCA GAG ACA ACT GAT CTC TAC TGT TAT GAG CAA TTA AAT
 glu tyr met leu asp leu gln pro glu thr thr asp leu tyr cys tyr glu gln leu asn
 661/221 691/231
 55 GAC AGC TCA GAG GAG GAG GAT GAA ATA GAT GGT CCA GCT GGA CAA GCA GAA CCG GAC AGA
 asp ser ser glu glu glu asp glu ile asp gly pro ala gly gln ala glu pro asp arg
 721/241 751/251
 60 GCC CAT TAC AAT ATT GTA ACC TTT TGT TGC AAG TGT GAC TCT ACG CTT CGG TTG TGC GTA
 ala his tyr asn ile val thr phe cys cys lys cys asp ser thr leu arg leu cys val
 781/261 811/271
 CAA AGC ACA CAC GTA GAC ATT CGT ACT TTG GAA GAC CTG TTA ATG GGC ACA CTA GGA ATT
 gln ser thr his val asp ile arg thr leu glu asp leu leu met gly thr leu gly ile
 841/281 871/291
 65 GTG TGC CCC ATC TGT TCT CAA GGA TCC atg gtg agc aag ggc gag gag ctg ttc acc ggg
 val cys pro ile cys ser gln gly ser met val ser lys gly glu glu leu phe thr gly
 901/301 931/311
 gtg gtg ccc atc ctg gtc gag ctg gac ggc gac gta aac ggc cac aag ttc agc gtg tcc
 val val pro ile leu val glu leu asp gly asp val asn gly his lys phe ser val ser
 961/321 991/331

ggc gag ggc gag ggc gat gcc acc tac ggc aag ctg acc ctg aag ttc atc tgc acc acc
 gly glu gly glu gly asp ala thr tyr gly lys leu thr leu lys phe ile cys thr thr
 1021/341 1051/351
 5 ggc aag ctg ccc gtg ccc tgg ccc acc ctc gtg acc acc ttc acc tac ggc gtg cag tgc
 gly lys leu pro val pro trp pro thr leu val thr thr phe thr tyr gly val gln cys
 1081/361 1111/371
 ttc agc cgc tac ccc gac cac atg aag cag cac gac ttc ttc aag tcc gcc atg ccc gaa
 phe ser arg tyr pro asp his met lys gln his asp phe phe lys ser ala met pro glu
 1141/381 1171/391
 10 ggc tac gtc cag gag cgc acc atc ttc ttc aag gac gac ggc aac tac aag acc cgc gcc
 gly tyr val gln glu arg thr ile phe phe lys asp asp gly asn tyr lys thr arg ala
 1201/401 1231/411
 gag gtg aag ttc gag ggc gac acc ctg gtg aac cgc atc gag ctg aag ggc atc gac ttc
 glu val lys phe glu gly asp thr leu val asn arg ile glu leu lys gly ile asp phe
 1261/421 1291/431
 15 aag gag gac ggc aac atc ctg ggc cac aag ctg gag tac aac tac aac agc cac aac gtc
 lys glu asp gly asn ile leu gly his lys leu glu tyr asn tyr asn ser his asn val
 1321/441 1351/451
 20 tat atc atg gcc gac aag cag aag aac ggc atc aag gtg aac ttc aag atc cgc cac aac
 tyr ile met ala asp lys gln lys asn gly ile lys val asn phe lys ile arg his asn
 1381/461 1411/471
 atc gag gac ggc agc gtg cag ctc gcc gac cac tac cag cag aac acc ccc atc ggc gac
 ile glu asp gly ser val gln leu ala asp his tyr gln gln asn thr pro ile gly asp
 1441/481 1471/491
 25 ggc ccc gtg ctg ctg ccc gac aac cac tac ctg agc acc cag tcc gcc ctg agc aaa gac
 gly pro val leu leu pro asp asn his tyr leu ser thr gln ser ala leu ser lys asp
 1501/501 1531/511
 30 ccc aac gag aag cgc gat cac atg gtc ctg ctg gag ttc gtg acc gcc gcc ggc atc act
 pro asn glu lys arg asp his met val leu leu glu phe val thr ala ala gly ile thr
 1561/521
 ctc ggc atg gac gag ctg tac aag
 leu gly met asp glu leu tyr lys

pcDNA3 plasmid vector:

(SEQ ID NO:15)

35 GACGGATCGG GAGATCTCCC GATCCCCTAT GGTGCGACTCT CAGTACAATC TGCTCTGATG CCGCATAGTT
 AAGCCAGTAT CTGCTCCCTG CTGTGTGTGT GGAGGTGCGT GAGTAGTGCG CGAGCAAAAT TTAAGCTACA
 ACAAGGCAAG GCTTGACCGA CAATTGCATG AAGAATCTGC TTAGGGTTAG GCGTTTTGCG CTGCTTCGCG
 ATGTACGGGC CAGATATACG CGTTGACATT GATTATTGAC TAGTTATTAA TAGTAATCAA TTACGGGGTC
 40 ATGATTTTCA AGCCCATATA TGGAGTCCG CGTTACATAA CTTACGGTAA ATGGCCCGCC TGGCTGACCG
 CCAACGACC CCCGCCATT GACGTCAATA ATGACGTATG TTCCCATAGT AACGCCAATA GGGACTTTCC
 ATTGACGTCA ATGGGTGGAG TATTTACGGT AAATGCCCCA CTTGGCAGTA CATCAAGTGT ATCATATGCC
 AAGTACGCCC CTTATTGACG TCAATGACGG TAAATGGCCC GCCTGGCATT ATGCCAGTA CATGACCTTA
 TGGGACTTTC CTACTTGGCA GTACATCTAC GTCTATATAA TCGCTATTAC CATGGTGATG CCGTTTTGGC
 45 AGTACATCAA TGGGCGTGGG TAGCGGTTTG ACTCACGGGG ATTTCCAAGT CTCCACCCCA TTGACGTCAA
 TGGGAGTTTG TTTTGGCACC AAAATCAACG GGACTTTCCA AAATGTCGTA ACAACTCCGC CCCATTGACG
 CAAATGGGCG GTAGGCGTGT ACGGTGGGAG GTCTATATAA GCAGAGCTCT CTGGCTAACT AGAGAACCCA
 CTGCTTACTG GCTTATCGAA ATTAATACGA CTCACTATAG GGAGACCCAA GCTGGCTAGC GTTTAAACGG
 GCCCTCTAGA CTCGAGCGGC CGCCACTGTG CTGGATATCT GCAGAAATCC ACCACACTGG ACTAGTGGAT
 50 CCGAGCTCGG TACCAAGCTT AAGTTTAAAC CGCTGATCAG TGACCCTGGA AGGTGCCACT CCCACTGTCC
 CTGTTGTTTG CCCCTCCCCC GTGCCTTCCT TGACCCTGGA AGGTGCCACT CCCACTGTCC TTTCTTAATA
 AAATGAGGAA ATTGCATCGC ATTGTCTGAG TAGGTGTCAT TCTATTCTGG GGGGTGGGGT GGGGCAGGAC
 AGCAAGGGG AGGATTGGGA AGACAATAGC AGGCATGCTG GGGATGCGGT GGGCTCTATG GCTTCTGAGG
 CGGAAAGAAC CAGCTGGGGC TCTAGGGGGT ATCCCCAGC GCCCTGTAGC GGCGCATTAA GCGCGGCGGG
 55 TGTGGTGGTT ACGCGCAGCG TGACCGCTAC ACTTGCCAGC GCCCTAGCGC CCGCTCCTTT CGCTTTCTTC
 CCTTCCTTTC TCGCCACGTT CGCCGGCTTT CCCCCTCAAG CTCTAAATCG GGGCATCCCT TTAGGGTTCC
 GATTTAGTGC TTTACGGCAC CTCGACCCCA AAAAATCTGA TTAGGGTGAT GGTTACGTA GTGGGCCATC
 GCCCTGATAG ACGGTTTTTC GCCCTTTGAC GTTGGAGTCC ACCTTCTTTA ATATTATAAGG GATTTTGGGG ATTTCCGCCT
 ACTGGAACAA CACTCAACCC TATCTCGGTC TATTCTTTTG GAATTAATTC TGTTGAATGT GTGTCAGTTA
 60 ATTGGTTAAA AAATGAGCTG ATTTAACAAA AATTTAACCG GAATTAATTC TGTTGAATGT GTGTCAGTTA
 GGGTGTGGAA AGTCCCCAGG CTCGCCAGGC AGGCAGAAAGT ATGCAAAAGCA TGCACTCTCA TTAGTCAGCA
 ACCAGGTGTG GAAAGTCCCC AGGCTCCCCA GCAGGCAGAA GTATGCAAAG CATGCATCTC AATTAGTCAG
 CAACCATAGT CCCGCCCTA ACTCCGCCCA TCCCCCCCCT AACTCCGCCC AGTTCCGCCC ATTCTCCGCC
 CCATGGCTGA CTAATTTTTT TTATTTATGC AGAGGCCGAG GCCGCTCTG CCTCTGAGCT ATTCAGAAAG
 65 TAGTGAGGAG GCTTTTTTGG AGGCCTAGGC TTTTGCAAAA AGCTCCCGGG AGCTTGATA TCCATTTTCG
 GATCTGATCA AGAGACAGGA TGAGGATCGT TTCGATGATG TGAACAAGAT GGATTGCACG CAGGTTCTCC
 GGCCGCTTGG GTGGAGAGGC TATTGCGCTA TGATCGGCA CAACAGACAA TCGGCTGCTC TGATGCCGCC
 GTGTTCCGGC TGTCAGCGCA GGGGCGCCCG GTTCTTTTTG TCAAGACCGA CCTGTCCGGT GCCCTGAATG
 AACTGCAGGA CGAGGCAGCG CGGCTATCGT GGCTGGCCAC GACGGGCGTT CCTTGCGCAG CTGTGCTCGA
 70 CGTTGTCACT GAAGCGGGA GGGACTGGCT GCTATTGGGG GAAGTGCCGG GGCAGGATCT CCTGTCACT
 CACCTTGCTC CTGCCGAGAA AGTATCCATC ATGGCTGATG CAATGCGGCG GCTGCATACG CTTGATCCGG
 CTACCTGCCC ATTCGACCAC CAAGCGAAAC ATCGCATCGA GCGAGCACGT ACTCGGATGG AAGCCGGTCT

5 TGTCGATCAG GATGATCTGG ACGAAGAGCA TCAGGGGGCTC GCGCCAGCCG AACTGTTTCGC CAGGCTCAAG
 GCGCGCATGC CCGACGGCGA GGATCTCGTC GTGACCCATG GCGATGCCTG CTTGCCGAAT ATCATGGTGG
 AAAATGGCCG CTTTTCTGGA TTCATCGACT GTGGCCGGCT GGGGTGGGCG GACCGCTATC AGGACATAGC
 GTTGGCTACC CGTGATATTG CTGAAGAGCT TGGCGGCGAA TGGGCTGACC GCTTCCTCGT GCTTTACGGT
 ATCGCCGCTC CCGATTTCGA GCGCATCGCC TTCTATCGCC TTCTTGACGA GTTCTTCTGA GCGGGACTCT
 GGGGTTTCGA ATGACCGACC AAGCGACGCC CAACCTGCCA TCACGAGATT TCGATTCCAC CGCCGCCTTC
 TATGAAAGGT TGGGCTTCGG AATCGTTTTT CCGGACGCCG GCTGGATGAT CCTCCAGCGC GGGGATCTCA
 TGCTGGAGTT CTTCGCCAC CCAACTTGT TTATTGCAGC TTATAATGGT TACAAATAAA GCAATAGCAT
 10 CACAAATTTT ACAAATAAAG CATTTTTTTC ACTGCATTCT AGTTGTGGTT TGTCAAAACAT CATCAAGTGA
 TCTTATCATG TCTGTATACC GTCCGACCTCT AGCTAGAGCT TGCGTAATC ATGGTCATAG CTGTTTCTG
 TGTGAAATTG TTATCCGCTC ACAATTCAC ACAACATACG AGCCGGAAGC ATAAAGTGTA AAGCCTGGGG
 TGCCTAATGA GTGAGCTAAT TCACATTAAT TGCGTTGCGC TCACTGCCCG CTTTCCAGTC GGGAAACCTG
 TCGTGCCAGC TGCATTAATG AATCGGCCAA CGCGCGGGGA TCGTTGCGCT GCGTATTGGG ACTCAAAGGC
 15 CTTCTCGCT CACTGACTCG CTGCGCTCGG TCGTTGCGCT TAACGCAGGA AAGAACATGT GAGCAAAAGG CCAGCAAAAG
 GGTAAATACCG TTATCCACAG AATCAGGGGA CGCGTTGCTG GCGTTTTTCC ATAGGCTCCG CCCCCTGAC GAGCAATACA
 GCCAGGAAGC GTAAAAAGG AGGTGGCGAA ACCCGACAGG ACTATAAAGA TACCAGCGT TTCCCTGG
 AAAATCGACG CTCAGTCTC CTGTTCCGAC CCTGCCGCTT ACCGGATACC TGTCCGCCTT TCTCCCTTCG
 AAGCTCCCTC GGAAGCGTGG CGCTTTCTCA ATGCTCACG TGTAGGTATC TCAAGTTCGGT GTAGGTCGTT CGCTCCAAGC
 20 TGGGCTGTGT GCACGAACCC CCCGCTGACG CCGACCGCTG GGCAGCAGCC ACTGGTAACA GGTAACATC GTCTTGAGTC
 CAACCCGGTA AGACACGACT TATCGCCACT CTTGAAGTGG TGGCCTAACT ACGGCTACAC TAGAAGGACA GTATTTGGTA
 GTAGGCGGTG CTACAGAGTT GTTACCTTCG TTGTTTGCAA GTGAGCAGATT ACAGCGAGAA AAAAAGGATC TCAAGAAGAT
 TCTGCGCTCT GCTGAAGCCA GTTACCTTCG GTCTGACGCT CAGTGGAACG AAAACTCACG TTAAGGGATT TTGGTCATGA
 25 CGCTGGTAGC GGTGGTTTTT TTCTACGGG GTCTGACGCT CAGTGAACG AAAACTCACG TTAAGGGATT TTGGTCATGA
 CCTTTGATCT TTTCTACGGG GTCTGACGCT CAGTGAACG AAAACTCACG TTAAGGGATT TTGGTCATGA
 GATTATCAAA AAGGATCTTC ACCTAGATCC TTTTAAATTA AAAATGAAGT TTTAAATCAA TCTAAAGTAT
 ATATGAGTAA ACTTGGTCTG ACAGTTACCA ATGCTTAAAT AGTGAGGCAC CTATCTCAGC GATCTCTCTA
 TTTGCTTCAT CCATAGTTGC CTGACTCCCC GTCGTGTAGA TAACTACGAT ACGGGAGGGC TTACCATCTG
 30 GCCCCAGTGC TGCAATGATA CCGCGAGACC CACGCTCACC GGCTCCAGAT TTATCAGCAA TAAACCAGCC
 AGCCGGAAGG GCCGAGCGCA GAAAGTGGTC TGCAATTTGA TCCGCCTCCA TCCAGTCTAT TAATTGTTGC
 CGGGAAGCTA GAGTAAGTAG TTCGCCAGTT AATAGTTTTG AATAGTTTTG GCAACGTTGT TGCCATTGCT ACAGAGGATC
 TGGTGTACAG CTCGTGTTTT GGTATGGCTT CATTACGCTC CGGTTCCCAA CGATCAAGGC GAGTTACATG
 ATCCCCATG TGTGCAAAA AAGCGGTTAG CTCCTTCGGT CTTCTACTGT CATGCCATCC CATGAGGATC
 35 GCAGTGTAT TACTGCAAGT TATGGCAGTA TATGGCAGTA CTTCTACTGT CATGCCATCC CATGAGGATC
 TTTCTGTGAC TGGTGAGTAC TCAACCAAGT CATTCTGAGA ATAGTGTATG ATAGTGTATG GCGCGACCGA GTTGCTCTTG
 CCCGGGTCA ATACGGGATA ATACCGCGCC ACATAGCAGA ACTTTAAAGG TGCTCATCAT TGGAAAACGT
 TCTTCGGGGC GAAAACCTC AAGGATCTTA CCGTGTGGA GATCCAGTTC GATGTAACCC ACTCGTGCAC
 CCAACTGATC TTCAGCATCT TTTACTTTCA CCAGCGTTTC TGGGTGAGCA AAAACAGGAA GGCAGGATGC
 40 CGAAAAAAG GGAATAAGGG CGACACGGAA ATGTTGAATA CTCATACTCT TGAATGTAT TTAGAAAAAT AAACAAATAG
 AGCATTTATC AGGGTTATTG TCTCATGAGC GGATACATAT CACCTGACGT C

SINrep5 self replicating replicon**(SEQ ID NO:16)**

(includes cloning sites)

45 ATTGACGGCG TAGTACACAC TATTGAATCA AACAGCCGAC CAATTGCACT ACCATCACAA TGGAGAAGCC
 AGTAGTAAAC GTAGACGTAG ACCCCCAGAG TCCGTTTGTG GTGCAACTGC AAAAAAGCTT CCCGCAATTT
 GAGGTAGTAG CACAGCAGGT CACTCCAAAT GACCATGCTA ATGCCAGAGC ATTTTCGCAT CTGGCCAGTA
 AACTAATCGA GCTGGAGGTT CCTACCACAG CGACGATCTT GGACATAGGC AGCGCACCGG CTCGTAGAAT
 50 GTTTTCCGAG CACCAATATC ATTGTGTCTG CCCATGCGT AGTCCAGAAG ACCCGGACCG CATGATGAAA
 TACGCCAGTA AACTGGCGGA AAAAGCGTGC AAGATTACAA ACAAGAACTT GCATGAGAAG ATTAAGGATC
 TCCGGACCGT ACTTGATACG CCGGATGCTG AAACACCATC CGTGTATATC AACGCTCCCG GAACTATCTA TCAATCAGGCT
 CATGCGTGCC GAATATTCCG TCATGCAGGA GTACTGGATT GCTTTCGACA CCACCCAGTT CATGTTCTCG CATATGGCAG
 55 ATGAAAGGCG TCGGACCCCT GTACTGGATT GCTTTCGACA CCACCCAGTT CATGTTCTCG CATATGGCAG
 GTTCGTACCC TGCGTACAAC ACCAACTGGG GTAGGACAGG AAAATTGTGCG ATAATGAGGA AGAAGGAGTT GAAGCCCGGG
 CAGCACAAAG CTGAGTGAAG AGGATCGACA CTTTATCCAG AACACAGAGC TCGGCTGCTT GATGAGGATC
 TCGCGGTTT ATTTCTCCGT AATGGAAGC AGTTCGTACAC TTGCCGCTGT TCGGCTGCTT GATGAGGATC
 TTCCATCGGT GTTCCACTTG TCACTATCAG TCCCGGGATC CAGTAAAAGG AGAACGGGTA TATCACTGA
 60 AGGCTACGTA GTGAAGAAAA GCTATGCAAA GCTATGACA GTTACTGACA TGGTATAATG GCCACGGATA
 CACAATAGCG AGGGCTTCTT ACCATATGCG ATCAGATGAC CCAGCGAATT GTATTAAACG GTAGGACTAA CAGGAACACC
 CGATGCACAA AAACCTCTGG TTGGGCTCAA ATAGACAAG ATAGACAAG GTTTCAGCAA ATGGGCTAAG GAGCGCAAGG
 AACACCATGC AAAATTACCT TCTGCCGATC CTAGAGAACG CAAGCTTACG TATGGCTGCT TGTGGCGTT
 ATGATCTTGA TAACGAGAAA CGTGTGGGTA CGTTTATCG CCCACCTGGA ACAGCAGACCT GCGTAAAAGT CCCAGCCTCT
 65 TCGCACTAAG AAAGTACATT GTCGGTATGG GATGCTCAGG AGGGAAGCCAG AGCGGAGAAG GTGAGGGGCT
 TTTAGCGCTT TTCCCATGTC GTCCGATGAG GATGCTCAGG AGGGAAGCCAG AGCGGAGAAG GTGAGGGGCT
 TGGCATTGCA ACCAAAAGAG GATGCTCAGG AGGGAAGCCAG AGCGGAGAAG GTGAGGGGCT
 TGCTTTTGAG GATGCTCAGG AGGGAAGCCAG AGCGGAGAAG GTGAGGGGCT TCAAGCAAAAT
 70 GACAAAGGCA TCGAGGAGC CGCAGAAAGT GGATAATACC GAATATCCAA CTCGACACAG CCGACCCGCT
 CATTAGTTGA AACCCCGCGC GGTACAGTAA GGTACATCCAA CTCGACACAG CCGACCCGCT
 TATCGTTGTC TCGCCAAACT CTGTGCTGAA

GTTAAGATCA TAACACACTC CGGAAGATCA GGAAGGTACG CGGTCGAACC ATACGACGCT AAAGTACTGA
 TGCCAGCAGG AGGTGCCGTA CCATGGCCAG AATTCTTAGC ACTGAGTGAG AGCGCCACGT TAGTGTACAA
 CGAAAGAGAG TTTGTGAACC GCAAACTATA CCACATTGCC ATGCATGGCC CCGCCAAGAA TACAGAAGAG
 5 GAGCAGTACA AGGTTACAAA GGCAGAGCTT GCAGAAACAG AGTACGTGTT TGACGTGGAC AAGAAGCGTT
 GCGTTAAGAA GGAAGAAGCC TCAGGTCTGG TCCTCTCGGG AGAACTGACC AACCTCCCT ATCATGAGCT
 AGCTCTGGAG GGAAGTGAAGA CCCGACCTGC GGTCCCGTAC TCAACTGTCA TCTTGTTACC AGCGGAAAGA
 CCGGGGTCCG GCAAGTCAGC TATTATCAAG TCAACTGTCA TCAACTGTCA TCAACTGTCA TCAACTGTCA
 AAGAAAAATTG TCGCGAAATT GAGGCCGACG TGCTAAGACT GTGCTGTACG TTGACGAAGC GTTCGCGTGC
 10 AGATTGCGTT ATGCTCAACG GATGCCACAA AGCCGTAGAA ATCGTCAGGC CCCGCAAGAA GGTAGTACTA TCGGAGACC
 CACGCAGGAG CACTACTTGC CTTGATTGCT AACTAAAGGT ACATTTCAAT CACCTGAAA CAGCTATTGT ATCGACACTG
 CCATGCAATG CGGATTCTTC AACATGATGC GCGTTGCACA CAGCCAGTTA AGAATTGA AATCGATATT ACAGGGGCCA
 CACCAAGACA TTCTACAAGT ATATCTCCCG CCGTGCACA CAGCCAGTTA AGAATTGA AATCGATATT ACAGGGGCCA
 CATTACGATG GAAAGATGAA AACCACGAAC CCGTGCACA CAGCCAGTTA AGAATTGA AATCGATATT ACAGGGGCCA
 15 CAAAGCCGAA GCCAGGGGAT ATCATCCTGA CATGTTTTCCG CCGGTGGGTT AAGCAATTGC AAATCGACTA
 TCCCGGACAT GAAGTAATGA CAGCCGCGGC CTCACAAGGG CTAACCAGAA AAGGAGTGTA TGCCGTCCGG
 CAAAAAGTCA ATGAAAACCC ACTGTACGCG ATCACATCAG AGCATGTGAA CGTGTTGCTC ACCCGCACTG
 AGGACAGGCT AGTGTGAAAA ACCTTGCAGG GCGACCCATG GATTAAGCAG CCCACTAACA TACCTAAAGG
 AAATTTTCAG GCTACTATAG AGGACTGGGA AGCTGAACAC AAGGGAATAA TTGCTGCAAT AAACAGCCCC
 20 ACTCCCCGTG CCAATCCGTT CAGCTGCAAG ACCAACGTTT GCTGGGCGAA AGCATTGGAA CCGATACTAG
 CCACGCCCGG TATCGTACTT ACCGTTTGCC ACTGTTGCCA TAGTGGTACG ATTCAGCGAG TGCCGCCGAA CTCTCCCGTA
 ACATTGCGCC ATTTACGCCT TAGACGTAAT TGGCATTAAAG TTTTTCGGCA TGGACTTGAC AAGCGGACTG
 TTTTCTAAAC AGAGCATCCC ACTAACGTAC CATCCCGCGG ATTCAGCGAG TGCCGCCGAA CTCTCCCGTA
 25 ACAGCCCAGG AACCCGCAAG TATGGGTACG ATACGCCAT TGCCGCCGAA CTCTCCCGTA CTCTGCACTG
 GTTCCAGCTA GCTGGGAAGG GCACACAAC CCAGCAATCTT CCTCACGCCT TAGTCCCGGA GTACAAGGAG AAGCAACCCG
 CATAACCTGG TCCCGGTGAA AAAATTCTTG AACCAGTTCA AACACCACTC CCGGATTTGGC ATAGCCGTTG TGAATAAGAA
 GCGCCGTCAT AAAAGTCTTG AAGAGAATCG AATGGATCGC GACCTGGTGT TCATCAACAT TGGAATAAAA TACAGAAACC
 30 AGCTCCCCGT TCCCGCCGCA TCCCGCCGCA GACCTGGTGT TCATCAACAT TGGAATAAAA TACAGAAACC
 GCTTTGCGGT ACCACTTTCA GCAGTGCAGG GACCATGCGG TCGTGGCTAC CTATGGCTAC GCAGCGAGAC CAGATTGTGT
 TAACCCAGGA GGCACCTCG TGGTGAAGT GCAGCGAGAC CAGATTGTGT CTCAAGCAAT CAGCAAAATG
 GCTCTTGCCA GAAAGTTTGT CAGGGTGTCT GACAAACAGC GTACACGGCA ATTCACCCCG CACCATCTGA ATTGCGTGAT
 35 ACCTGATTTT CCGACAACCTA GTCAAGAGGA AGCAGTTGTC TGGCCGACCA GCGCGTCCG CCGCGTCCG CCTGATTTCC
 TTCGTCGTTG TATGAGGTA CAAGAGATGG AGCAGTTGTC TGGCCGACCA GCGCGTCCG CCGCGTCCG CCTGATTTCC
 ATTGCTGACT GTCAAGAGGA AGCAGTTGTC TGGCCGACCA GCGCGTCCG CCGCGTCCG CCTGATTTCC
 GCCGTGCCAT CTATAAACGT TGGCCGACCA GCGCGTCCG CCGCGTCCG CCTGATTTCC
 40 GACTGTGTGC CTAGGAAAGA AAGTGATCCA CGCTACCAT ACAGGCATTT GAGCTAGACA GAACTGACGC
 GCCTTGAATG TACTAGGTA ACTGCTATCT ACAGGCATTT GAGCTAGACA GAACTGACGC
 TCGCCATTCC ACTGCTATCT ACAGGCATTT GAGCTAGACA GAACTGACGC
 CTTGACAACC GCGCTAGACA GAACTGACGC GCGCACTCCA ACTTAAGAGG CCAGACAGTT GCACCAAAATT
 45 AGAATCGACG ATGGATTAT TACTTGAAG GCACCAAAATT TAATGAACAA CATAAAGAA TCCACAGACT
 ATGAGTTAGT ATGGATTAT TACTTGAAG GCACCAAAATT TAATGAACAA CATAAAGAA TCCACAGACT
 ATTGTATTCC TACTTGAAG GCACCAAAATT TAATGAACAA CATAAAGAA TCCACAGACT
 TCCCTAATG ACCAGGAAAG CCGGTCGAC CCAGAAAGGG CTAAGCACAA AATTAAGAA TCCACAGACT
 50 GCGAAAGTG CCGGTCGAC CCAGAAAGGG CTAAGCACAA AATTAAGAA TCCACAGACT
 TGCCATGACG CTAAGCACAA AATTAAGAA TCCACAGACT CCGCACTCC CGCCTACCG CAGCCTACCG CTGCTCTGAT
 ACCCCCTTC CTAAGCACAA AATTAAGAA TCCACAGACT CCGCACTCC CGCCTACCG CAGCCTACCG CTGCTCTGAT
 CGCACACTCC CGCCTACCG CAGCCTACCG CTGCTCTGAT AGCGGATCGG
 55 ACAGGCCGAG GAGGCCCCCG AAGTTGTAGC TATGGATGAC GACAGTTGTT ATGCCGTC CAGCGGCAAG
 GTACAGACA TCTACTGGA TACTAGTATG GCTGACGTTT CAGCGGCAAG AAAAGAGCCC TCCCTCGGAT
 ACAACTCTAT TACTAGTATG GCTGACGTTT CAGCGGCAAG AAAAGAGCCC TCCCTCGGAT
 GGTGGTGGTG GCTGACGTTT CAGCGGCAAG AAAAGAGCCC TCCCTCGGAT
 60 GCGGCTGGG GGTATCCATG CCGATGAGCC TAGTGGGTA GCTTACAGAA GAGGAACAAC AAGTAGAAAA
 ACCCCTGGCA ACAGGCCCA GAGTAAGTGA GTCCGAACCC GTATCTTTTC TAGTGGGTA GCTTACAGAA
 TTATATCGTC CCGATCAGCC TAGTGGGTA GCTTACAGAA GAGGAACAAC AAGTAGAAAA
 TGAATACTGA CTAACCGGGG TAGTGGGTA GCTTACAGAA GAGGAACAAC AAGTAGAAAA
 AAGTCCGTTT TGCAGAACCA CACGTCGAAA AGTCTCGTA AGTCTCGTA AGTCTCGTA AGTCTCGTA
 65 CGGTGCTCGA CAGTCTCGTA AGTCTCGTA AGTCTCGTA AGTCTCGTA AGTCTCGTA AGTCTCGTA
 AAGTAGTAC ACTCTGCCA GTAGCGTACC GGCAGTACC ATCAGATTAC AACTCTGCTT TGGATACTGC
 CGACTGTATA ACTCTGCCA GTAGCGTACC GGCAGTACC ATCAGATTAC AACTCTGCTT TGGATACTGC
 GTAGCGTACC GGCAGTACC ATCAGATTAC AACTCTGCTT TGGATACTGC
 70 CTATCCGACA TTAGCATCTT TGGATACTGC TTAGCATCTT TGGATACTGC TTAGCATCTT TGGATACTGC
 GTCCGCTGCC TTAGCATCTT TGGATACTGC TTAGCATCTT TGGATACTGC TTAGCATCTT TGGATACTGC
 GAGCCCCGAA TATCCGAGT AACTGCAACG GAAAAATGAC ATGTAATGAC TATGTAAGTCA AAGAAGTGCC
 AACTAAAAAGA AATTGCAACG GAAAAATGAC ATGTAATGAC TATGTAAGTCA AAGAAGTGCC
 GAATGCTTTC TGTACCCGCA GTCCCATGTC GCACGAAACA CACAGAAAGG AATTACCCGG CCGCGGAGGA
 CCACTGAGTT TGTACCCGCA GTCCCATGTC GCACGAAACA CACAGAAAGG AATTACCCGG CCGCGGAGGA
 GTATAATTTG GTCCCATGTC GCACGAAACA CACAGAAAGG AATTACCCGG CCGCGGAGGA
 GTTACACCG CACTGCTTA TTTGACATGT AGACGGATAT CGCATCATTC

5 TGATCTTGGA GGACCTGGGT GTGGATCAAC CACTACTCGA CTTGATCGAG TGCGCCTTTG GAGAAATATC
 ATCCACCCAT CTACCTACGG GTACTCGTTT TAAATTCGGG GCGATGATGA AATCCGGAAT GTTCCTCACA
 CTTTTTGTCA ACACAGTTTT GAATGTCTGT AGTCCAGCA GAGTACTAGA AGAGCGGCTT AAAACGTCCA
 GATGTGCAGC GTTCATTGGC GACGACAACA TCATACATGG AGTAGTATCT GACAAAGAAA TGGGTGAGAG
 GTGCGCCACC TGGCTCAACA TGGAGGTTAA GATCATCGAC GCAGTCATCG GTGAGAGACC ACCTTACTTC
 TGCGGCGGAT TTATCTTGCA AGATTCGGTT ACTTCCACAG CGTGCCGCGT GGCGGATCCC CTGAAAAAGG
 TGTTTAAAGT GGGTAAACCG CTCCCAGCCG ACGACGAGCA AGACGAAGAC AGAAGACGCG CTCTGCTAGA
 TGAACCAAAG GCGTGGTTTA GAGTAGGTAT AACAGGCACT TTAGCAGTGG CCGTGACGAC CCGGTATGAG
 10 GTAGACAATA TTACACCTGT CCTACTGGCA TTGAGAACTT TTGCCAGAG CAAAAGAGCA TTCCAAGCCA
 TCAGAGGGGA AATAAAGCAT CTCTACGGTG TGAAGACGAA GTCAGCATAG TACATTTTCT CTGACTAATA
 CTACAACACC ACCACCTCTA GACGCGTAGA TCTCACGTGA GCATGCAGGC CTTGGGCCCA ATGATCCGAC
 CAGCAAAACT CGATGTACTT CCGAGGAACT CTGATGTAC CTGGTACATT AGATCCCCCG
 TTACCCGCGG CAATATAGCA ACACTAAAAA CTCGATGTAC TTCCGAGGAA GCGCAGTGCA TAATGCTGCG
 15 CAGTGTTGCC ACATAACCAC TATATTAACC ATTTATCTAG CCGACGCCAA AAACCTCAATG TATTTCTGAG
 GAAGCGTGGT GCATAATGCC ACGCAGCGTC TGCATAACTT TTATTATTTT TTTTATTAAT CAACAAAATT
 TTGTTTTTAA CATTTCAAAA AAAAAAAAAA AAAAAAAAAA AAAAAAAAAA AAAGGGAATT CCTCGATTAA
 TTAAGCGGCC GCTCGAGGGG AATTAATTCT TTTCTCACC CTTCCTGTTT AGGGCCAGGT GCGCAATTTT
 GCGCGGAACC CCTATTTGTT TATTTTCTA AATACATTCA AATATGTATC CGCTCATGAG ACAATAACCC
 20 TGATAAATGC TTCAATAATA TTGAAAAAGG AAGAGTATGA GTATTCAACA TTTCCGTGTC GCCCTTATTC
 CCTTTTTTGC GATCTTTTGC TTTCTGCTTT TTTCTGCGC AGAAACGCTG GTGAAAGTAA AAGATGCTGA
 AGATCAGTTG GGTGCACGAG TGGGTTACAT CGAACTGGAT CTCAACAGCG GTAAGATCCT TGAGAGTTTT
 CGCCCCGAAG AACGTTTTCC AATGATGAGC ACTTTTAAAG TTCTGCTATG TGGCGCGGTA TTATCCCGTG
 TTGACGCGCG GCAAGAGCAA CTCGGTCGCC GCATACATA TTCTCAGAAT GACTTGGTTG AGTACTCACC
 25 AGTCACAGAA AAGCATCTTA CCGATGGCAT GACAGTAAGA GAATTATGCA GTGCTGCCAT AACCATGAGT
 GATAACACTG CCGCCAACCT ACTTCTGACA ACGATCGGAG GACCGAAGGA GCTAACCGCT TTTTTCACA
 ACATGGGGGA TCATGTAACG CGCCTTGATC GTTGGGAACC GGAGCTGAAT GAAGCCATAC CAAACGACGA
 GCGTGACACC ACGATGCTG TAGCAATGGC AACAACGTTG CGCAAACCTAT TAAGTGGCGA ACTACTTACT
 CTAGCTTCCC GGCAACAATT AATAGACTGG ATGGAGGCGG ATAAAGTTGC AGGACCACTT CTGCGCTCGG
 30 CCCTTCCGGG TGCTGGTTT ATTGCTGATA AATCTGGAGC CCGTGAGCGT GGGTCTCGCG GTATCATTGC
 AGCACTGGGG CCAGATGGTA AGCCCTCCCG TATCGTAGTT ATCTACACGA CGGGGAGTCA GGCAACTATG
 GATGAACGAA ATAGACAGAT CGCTGAGATA GGTGCCTCAC TGATTAAGCA TTGGTAACTG TCAGACCAAG
 TTTACTCATA TATACTTTAG ATTGATTTAA AACTTCATTT TTAATTTAAA AGGATCTAGG TGAAGATCCT
 TTTTGATAAT CTCATGACCA AAATCCCTTA ACGTGAGTTT TCGTTCCTCT GAGCGTCAGA CCCCGTAGAA
 35 AAGATCAAAG GATCTTTTGG AGATCTTTT CTCTGCGCGG TAATCTGCTG TAATCTGCTG CTTGCAAACA AAAAAACCAC
 CGCTACCAGC GGTGGTTTGT TTGCCGGATC AAGAGTACC AACTCTTTTT CCGAAGGTAA CTGGCTTCAG
 CAGAGCGCAG ATACCAAATA CTGTCTTCT AGTGAGCCG TAGTTAGGCC ACCACTTCAA GAACTCTGTA
 GCACCGCCTA CACATCTCGC TCTGCTAATC CTGTTACCAG TGGCTGCTGC CAGTGGCGAT AAGTCTGTC
 40 TTACCGGGTT GGAATCAAGA CGATAGTTAC CGGATAAGGC GCAGCGGTGC GGCTGAACGG AAGTCTGTC
 CACACAGCCC AGCTTGGAGC GAACGACCTA CACCGAAGTG AGATACCTAC AGCGTGAGCA TGGAGAAAGC
 GCCACGCTTC CCGAAGGGAG AAAGGCGGAG AAGGTATCCG TAAGCGGCAG GGTGGAACA GGAGAGCGCA
 CGAGGGAGCT TCCAGGGGGA AACGCCTGGT ATCTTTATAG TCTGTGCGGG TTTGCGCCAC TGTGACTTGA
 GCGTCGATTT TTGTGATGCT CGTCAGGGGG GCGGAGCCTA TGGAAAAACG CCAGCAACGC GAGCTCGTAT
 45 GGACATATTG TCGTTAGAAC GCGGCTACAA TTAATACATA ACCTTATGTA TCATACACAT ACGATTTAGG
 GGACACTATA G

pSCA1 suicide DNA vector**SEQ ID NO:17:**

(includes cloning sites)

50 ATGGCGGATG TGTGACATAC ACGACGCCAA AAGATTTTGT TCCAGCTCCT GCCACCTCCG CTACGCGAGA
 GATTAACCAC CCACGATGGC CGCCAAAGTG CATGTTGATA TTGAGGCTGA CAGCCCATTC ATCAAGTCTT
 TGCAGAAAGC ATTTCCGTCG TTCGAGGTGG AGTCATTGCA GGTCAACCA AATGACCATG CAAATGCCAG
 AGCATTTTTC CACCTGGCTA CCAAATTGAT CGAGCAGGAG ACTGACAAAG ACACACTCAT CTTGGATATC
 GGCAGTGCGC CTTCCAGGAG AATGATGTCT ACGCACAAT ACCACTGCGT ATGCCCTATG CGCAGCGCAG
 55 AAGACCCCGA AAGGCTCGAT AGCTACGCAA AGAACTGGC AGCGGCCTCC GGGAAGGTGC TGGATAGAGA
 GATCGCAGGA AAAATCACCG ACCTGCAGAC CGTCATGGCT ACGCCAGACG CTGAATCTCC TACCTTTTGC
 CTGCATACAG ACGTCACGTG TCGTACGGCA GCCGAAGTGG CCGTATACCA GGACGTGTAT GCTGTACATG
 CACCAACATC GCTGTACCAT CAGGCGATGA AAGGTGTCAG AACGCGGTAT TGGATTGGGT TTGACACCAC
 CCCGTTTTAT TTTGACGCGC TAGCAGGCGC GTATCCAACC TACGCCCAA ACTGGGCCGA CGAGCAGGTG
 60 TTACAGGCCA GGAACATAGG ACTGTGTGCA CTGAGGGGAA CTGAGGGGAA ACTCGGCAAA CTGTCCATT
 TCCGCAAGAA GCAATTGAAA CCTTGCGACA CAGTCATGTT CTCGGTAGGA TCTACATTGT AACTGAGAG
 CAGAAAGCTA CTGAGGAGCT GGCACCTTACC CTCCGTATTC CACCTGAAAAG GTAAACAATC CTTTACCTGT
 AGGTGCGATA CCATCGTATC ATGTGAAGGG TACGTATTGA TACGTGAAGG AGAAAAATC TATGTGCCCC
 65 GTAAAACGGT AGGGTACGCC GTGACGTATC ACGCGGAGGG ATTCTAGTG TGCAAGACCA CAGACACTGT
 CAAAGGAGAA AGAGTCTCAT TCCCTGTATG CACCTACGTC CCCTCAACCA TCTGTGATCA AATGACTGGC
 ATACTAGCGA CCGACGTAC ACGGAGGAG GCACAGAAG TGTTAGTGGG ATTGAATCAG AGGATAGTTG
 TGAACGGAAG AACACAGCGA AACACTAACA CGATCAGAAG CTATCTGCTT CCGATTGTGG CCGTCGATT
 70 TAGCAAGTGG GCGAGGGAAT ACAAGGCAGA CCTTGATGAT GAAAAACCTC TGGGTGTCCG AGAGAGGTCA
 CTTACTTGCT GCTGCTTGTG GGCATTTAAA ACGAGGAAGA TGCACACCAT GTACAAGAAA CCAGACACCC
 AGACAATAGT GAAGGTGCGT TCAGAGTTTA ACCTGTTCTC CTATGGTCTA CAGGCTCGC
 AATCCAGTCA AGATCACGCA TTAAGATGCT TTTGGCCAAG AAGACCAAGC GAGAGTTAAT ACCTGTTCTC
 GACGCGTCGT CAGCCAGGGA TGCTGAACAA GAGGAGAAGG AGAGGTTGGA GGCCGAGCTG ACTAGAGAAG
 CCTTACCACC CCTCGTCCCC ATCGCGCCGG CGGAGACGGG AGTCGTCGAC GTCGACGTTG AAGAAGTAGA

5 GTATCACGCA GGTGCAGGGG TCGTGGAAC ACCTCGCAGC GCGTTGAAAG TCACCGCACA GCCGAACGAC
 GTACTACTAG GAAATTACGT AGTTCTGTCC CCGCAGACCG TGCTCAAGAG CTCCAAGTTG GCCCCCCGTGC
 ACCCTCTAGC AGAGCAGGTG AAAATAATAA CACATAACCG GAGGCGCCGC GGTTACCAGG TCGACGGATA
 TGACCGCAGG GTCCTACTAC CATGTGGATC GGCCATTCCG GTCCTGAGT TTCAAGCTTT GAGCGAGAGC
 10 GCCACTATGG TGTACAACGA AAGGGAGTTC GTCAACAGGA AACTATACCA TATTGCCGTT CACGGACCGT
 CGCTGAACAC CGACGAGGAG AACTACGAGA AAGTCAGAGC TGAAAGAACT GACGCCGAGT ACGTGTTCGA
 CGTAGATAAA AAATGCTGCG TCAAGAGAGA GGAAGCGTCG GGTTCGTTGT TGGTGGGAGA GCTAACCAAC
 CCCCCGTTCC ATGAATTCGC CTACGAAGGG CTGAAGATCA GGCCGTCGGC ACCATATAAG ACTACAGTAG
 TAGGAGTCTT TGGGGTTCG GGATCAGGCA AGTCTGCTAT TATTAAGAGC CTCGTGACCA AACACGATCT
 15 GGTCACCAGC GGCAAGAAGG AGAACTGCCA CACATGCGA AAGTATGATT AACGACGTGA AGAAGCACCG CGGAAGGGG
 ACAAGTAGGG AAAACAGTGA CTCCATCCTG TCCGCTACTC TGGTCTGCTT TTAAGGTGAA CTTCAACCAC
 ACGAGGCTTT CGCTAGCCAT AGCAATGCGG TCATAAAAGT ATATCCAGAC GTTGACGCG TCCAGTCCGT
 GGTGTTATGC GGAGACCCCA AGCAATGCGG TCATAAAAGT ATATCCAGAC GTTGACGCG TCCAGTCCGT
 AACATCTGCA CTGAAGTATG CTACGGAGGC AAGATGCGCA CGACCAACCC GTGCAACAAA CCCATAATCA
 20 CTACGTTGCA CTACGGAGGC AAGCCCAAGC CAGGAGACAT CGTGTTAACA TGCTTCCGAG GCTGGGCAAA
 AGGACAGACC GTGGACACGA AGTCAAGCAT CTCAGGCGCT CACCCGCAAA CACCGTGAATG TACTGCTGAC
 TTGGACTACC GTGGACACGA AGTCAAGCAT CTCAGGCGCT CACCCGCAAA CACCGTGAATG TACTGCTGAC
 CCGTAAGGCA GAAGGTGAAT GAAAATCCCT TGTATGCCCC GATCCCTGGA TTAAGGTCTT ATCAAACATT
 GCGCACTGAG GATAGGCTGG TGTGAAAAAC GCTGGCCGCG GAATGTGCAAG AAGAACACGA CAAAATAATG
 25 CCACAGGGTA ACTTTACGGC CACATGCGA AAGTATGATT TGTATTACGA GAACGTGTGT TGGGCGAAAA
 AAGGACCGGC TGCGCCTGTG GACGCGTTC TCAAGATTGAC AATGAAATTT TGGAGCACCA TAATTACAGC
 GTGCTGGAG ACTGCCGGA GGTGGCCTTG AATGAAATTT TGTATTACGA GAACGTGTGT TGGGCGAAAA
 GACAGAGCTT ACTCTCAGT GGTGGCCTTG AATGAAATTT TGTATTACGA GAACGTGTGT TGGGCGAAAA
 GTGGCCTGTT TTCTGCCCCG AAGGTGTCCC TGTATTACGA TGCCAGGCTG GAACTAGAC ATACCTTCCT
 30 AAGGATGTAT GGATTCAATG CCGCAACAGC AGTTATCGCA GAAAGAAAAA TCCAACCGCT TTCTGTGCTG
 TGGCATAACG GCAAGCAGGC CCGCAGCGCC CCGCAGCGCC GTTAAAGACG GTTAAAGGCA GTTAAAGGCA
 TTCTATCAA CCGCAGCTG AAGGATGTAT GGATTCAATG CCGCAACAGC AGTTATCGCA GAAAGAAAAA
 GTGGCTGGTC AATAAAGTAA GAGGGTACCA CCGCAGCGCC CCGCAGCGCC GTTAAAGACG GTTAAAGGCA
 35 CGCAGGGTCA CTGGTTGTC ACCGCTGAAT TCGACTTGG CCATGAAGCT TTACGGATAC GCGGATAAAA
 TGCCGGCTGA CGCCGGCAGG TCGACTTGG CCATGAAGCT TTACGGATAC GCGGATAAAA
 CCAGCAGTGT GTCGACCAG GTGTTGCGCC CCGGATAAAA GCGGATAAAA CACCAGCAAT
 GCGCGCATCT TGATGAGAGC GTGTTGCGCC CCGGATAAAA GCGGATAAAA CACCAGCAAT
 GAAAGTTCTC GTCTGCAAGA GACAACGGAA AGAGACCCCT TACGTTACAC CAGATGAATA
 40 CTCCAACCTT GACTGAGAGC CCATGCACAC GGCCGGGTGT AACGCAAGTA ACGCCCGTGG
 GCCGGAGAAG GCTGTGGTT GACTGAGAGC CCATGCACAC GGCCGGGTGT AACGCAAGTA ACGCCCGTGG
 GCACAGAAGC GGCTGTGGTT GACTGAGAGC CCATGCACAC GGCCGGGTGT AACGCAAGTA ACGCCCGTGG
 GGCAGAAGAA TGCCCGTCAG CCGTCAATCA CGCTGTAGCG CCGTCAATCA CGCTGTAGCG
 45 GGCTCGTACC CCGTCAATCA CGCTGTAGCG CCGTCAATCA CGCTGTAGCG CCGTCAATCA CGCTGTAGCG
 AATTGGCCCG TGCTACCGG ACAGGAGTGT CCACGGACGC CATGAGGACG GATGAGGACG
 GCTGCTGTC CCACGGACGC CATGAGGACG GATGAGGACG GATGAGGACG GATGAGGACG
 GCAATGGACG CCACGGACGC CATGAGGACG GATGAGGACG GATGAGGACG GATGAGGACG
 50 AAGCCATTGA GACAGCAGC CGAAATTCAG CGAAGCAGATA TCCGATTCTAT
 TTTGAAGGTA CGAAATTCAG CGAAGCAGATA TCCGATTCTAT TCCGATTCTAT
 AAGAGGCAAA CGAAGCAGATA TCCGATTCTAT TCCGATTCTAT TCCGATTCTAT
 GGTGAACGAT TCCGATTCTAT TCCGATTCTAT TCCGATTCTAT TCCGATTCTAT
 55 GAACGAGTGC TCCGATTCTAT TCCGATTCTAT TCCGATTCTAT TCCGATTCTAT
 AATACCATGT AGATGGGGTG CCGCGGAAGT GATGAGGACG GATGAGGACG
 AGTGGTTAGT CCGCGGAAGT GATGAGGACG GATGAGGACG GATGAGGACG
 TTGGACTGGA CACCCGACTC CATCTACGAG CTGGCGGACG CATCGCGGAC
 60 ACATCGACTC CATCTACGAG CTGGCGGACG CATCGCGGAC CATCGCGGAC
 CATCGCGGAC CATCTACGAG CTGGCGGACG CATCTACGAG CTGGCGGACG
 CCACCGCGCC CGAAGAGAGC TGCCCCAAGG GCGTTGGCCT TTTTCTCCTC
 AGCCGACGCC GCGTTGGCCT TTTTCTCCTC TTTTCTCCTC TTTTCTCCTC
 65 CGAGGTCGAT GCGTTGGCCT TTTTCTCCTC TTTTCTCCTC TTTTCTCCTC
 GGTGCATATA TTTTCTCCTC TTTTCTCCTC TTTTCTCCTC TTTTCTCCTC
 TCCAGTGCGC ACAACTGGAT CTGCTGAAAA TGAAAGCCAC GATGTAGCAA
 GAAGCTGTTG CTGCTGAAAA TGAAAGCCAC GATGTAGCAA GATGTAGCAA
 70 GTGGAGAACA ACACAGAGGA AAGATTCGAT TACAGATGCA GGTGGTGGAC
 TAGGCGGCAT ACACAGAGGA AAGATTCGAT TACAGATGCA GGTGGTGGAC
 CTCAAGCCCC GATGTAGCAA GGTGGTGGAC GGTGGTGGAC GGTGGTGGAC
 TACCAGATAA CAGATGAATA CGACGCATAC TCGAGCGTG CGACGCATAC
 CGACATTTCT CCCGGCAAG CTCCGGTGCT AGAACACATG ACCCACCATG
 TGCCGTCGCG TCAACCTTTT ACACACATG GAAAGATATG GAAGAATATG
 75 GTCACGCAAA TGCGAGAACT AGAATATTGG CCATGGACAG CCATGGACAG
 CCTGCTCCGG AATTTGAAAG CCATGGACAG CCATGGACAG CCATGGACAG
 CTATGTGACC AATTTGAAAG CCATGGACAG CCATGGACAG CCATGGACAG
 CAGGAGGTTT CCATGGACAG CCATGGACAG CCATGGACAG CCATGGACAG
 ACACAGAGGA AAGATTCGAT TACAGATGCA GGTGGTGGAC GGTGGTGGAC
 CATCCACAGG GAATTAGTAA GGAGACTAAA TGCTGTGTTA TGCTGTGTTA
 TCGGCCGAAG ACTTTGACGC CGACAAAAGC CAGGACGACT CTTGGCTCT
 TTGCATCATT TACCTGCTGG ACTTGATCGA GGCAGCCTTT GGCAGCCTTT

5 GGCACGCGCT TCAAGTTCGG AGCTATGATG AAATCGGGCA TGTTCCTGAC TTTGTTTATT AACACTGTTT
 TGAACATCAC CATAGCAAGC AGGGTACTGG AGCAGAGACT CACTGACTCC GCCTGTGCGG CCTTCATCGG
 CGACGACAAC ATCGTTTACG GAGTGATCTC CGACAGAGCT ATGGCGGAGA GGTGCGCGTC GTGGGTCAAC
 ATGGAGGTGA AGATCATTGA CGCTGTCATG GGGCAAAAAC CCCCATATTT TTGTGGGGGA TTCATAGTTT
 10 TTGACAGCGT CACACAGACC GCCTGCCGTG TTTTCAGACCC ACTTAAGCGC CTGTTCGAAGT TGGGTAAGCC
 GCTAACAGCT GAAGACAAGC AGGACGAAGA CAGGCGACGA GCACTGAGTG ACGAGGTTAG CAAGTGGTTC
 CGGACAGGCT TGGGGGGCCGA ACTGGAGGTG GCACTAACAT CTAGGTATGA GGTAGAGGGC TGCAAAAGTA
 TCCTCATAGC CATGGCCACC TTGGCGAGGG ACATTAAGGC GTTTAAGAAA TTGAGAGGAC CTGTTATACA
 CCTCTACGGC GGTCTAGAT TGGTGCGTTA ATACACAGAA TTCTGATTgg atccCGGGTA ATTAATTGAA
 15 TTATGCCCT ACGCAAACGT TTTACGGCCG CCGGTGGCGC CCAGCAGAT TAGGCCTCCC AAACCAAGA AGAAGAAGAC
 CAGGCCACTC GCGTGGCTCC CGTCTGTCCTT TTTACGCTCC GACTTCCAGG CCCAGCAGAT GCAGCAACTC ATCAGCGCCG
 TAAATGCGCT GACAATGAGA CAGAACGCAA GCGAGCCCAA GAAGTCAAC GGAACCAACG AGCAGCAAAA GAAGAAAGAC
 AACCAGGCG ACAAGAAGAA GAAGAAACCC GGAAGAAAGAG AAAGAATGTG CATGAAGATT GAAATGACT
 20 GTATCTTCGT ATGCGGCTAG CCACAGTAAC CCACAGTAAC CTGAGGTCG CGCTATCCTG GTGCTGGTTG TGGTCACTTG
 TGCAGAAAAA CTCGGGTGGT CTGGGGGCCCT TTAGGGTAGG CAATGAATACG GATATAGCAA AACAAGAAAG
 CATTGGGCTC CCGAGATAAG GCAATGGCAT ATAACCATAA CTGTATAACT TGTAACAAAG CGCAACAAGA CCTGCGCAAT
 GTTAGGGTAA GTCCGCCTCA CGGAACTCG GGGCAACTCA TATTGACACA TTTATTTTGC AATTGGTTTT TAATATTTCC
 25 GCTTACATAA GCCTTAATTG ACGAATAATT TTTATTTTGC AAAAAAATAA AAAAAAATAA AAAAAAATAA
 AAAAAAATAA AAAAAAATAA AAAAAAATAA AAAAAAATAA AAAAAAATAA AAAAAAATAA AAAAAAATAA
 CTAGTgatca taatcagcca taccacattt gtagaggttt tacttgcttt taacttgctt attgcagctt ataattggtta
 ccctgaacct gaaacataaa caaatgtcac aaataaagca tttttttcac tttttttcac tttttttcac tttttttcac
 30 caataaagc aatagcatca caaatgtcac aaataaagca tttttttcac tttttttcac tttttttcac tttttttcac
 tccaaactca tcaatgtatc ttatcatgtc ttatcatgtc ttatcatgtc ttatcatgtc ttatcatgtc ttatcatgtc
 AGAGGCGGTT TGGGTATTGG CACTCAAAGG GCCAGCAAAA CGAGCATCAC TTTCCCTTG TTTCCCTTG TTTCCCTTG
 TGCGGCGAGC GGTATCAGT GCGGCTCTCC CCGTAATACG CGGTAATACG CGGTAATACG CGGTAATACG
 35 AAAGAACATG TGAGCAAAAG GCGGCTCTCC CCGTAATACG CGGTAATACG CGGTAATACG CGGTAATACG
 CATAGGCTCC GCGGCTCTCC CCGTAATACG CGGTAATACG CGGTAATACG CGGTAATACG CGGTAATACG
 GACTATAAAG ATACCAGGCG TTTCCCTTG TTTCCCTTG TTTCCCTTG TTTCCCTTG TTTCCCTTG
 40 TACCGGATAC CTGTCCGCTT TCGCTCCAAG CCGTCTTGAGT GAGCGAGGTA AGTATTTGGT AAACAAACCA
 CTCAGTTCCG CGGTAACAT GAGCGAGGTA AGTATTTGGT AAACAAACCA CCGCTGGTAG TCCTTTGATC
 GCGCCTTATC CGGTAACAT GAGCGAGGTA AGTATTTGGT AAACAAACCA CCGCTGGTAG TCCTTTGATC
 45 CACTGGTAAC AGGATTAGCA CTAGAAGGAC TTGATCCGGC AACGAAACT CACGTTAAGG AATGTTTTAA
 TACGGCTACA TTGATCCGGC AACGAAACT CACGTTAAGG AATGTTTTAA AATCAGTGAG CACACCTATG
 TTGGTAGCTC TTGATCCGGC AACGAAACT CACGTTAAGG AATGTTTTAA AATCAGTGAG CACACCTATG
 50 TACGCGCAGA AAAAAAGAT TGTATCCAGT TGCTTACAGG AGGCGAGTGA AGGCGAGTGA AGGCGAGTGA
 AACGAAACT CACGTTAAGG AATGTTTTAA AATCAGTGAG CACACCTATG AATCAGTGAG CACACCTATG
 55 ATTAATAATG AAGTTTTAAA GCACCTATG GGGCTTACCA GCAATAAACC CTATTAATTG TGCTTACAGG
 AATCAGTGAG CACACCTATG AATCAGTGAG CACACCTATG AATCAGTGAG CACACCTATG AATCAGTGAG
 TAGATAACTA CGATACGGGA AGATTTATCA CTATTAATTG TGCTTACAGG AGGCGAGTGA AGGCGAGTGA
 60 CACCGGCTCC AGATTTATCA CTATTAATTG TGCTTACAGG AGGCGAGTGA AGGCGAGTGA AGGCGAGTGA
 TTTATCCGCT TCCATCCAGT TGCTTACAGG AGGCGAGTGA AGGCGAGTGA AGGCGAGTGA AGGCGAGTGA
 65 TTGCGCAACG TGTGTTGCCAT TGTGTTGCCAT TGTGTTGCCAT TGTGTTGCCAT TGTGTTGCCAT TGTGTTGCCAT
 GCTCCGGTTC CCAACGATCA AGGCGAGTGA AGGCGAGTGA AGGCGAGTGA AGGCGAGTGA AGGCGAGTGA
 CGGTCTCCG ATCGTTGTCA ATCCGTAAGA CCGAGTTGCT TCATTGGAAA ACCCACTCGT GCAACCACT
 AATTCTCTTA TATGCGGCGA AAAGTGCTCA GTTTCGATGA AGCAAAAACA GGAAGGCAAA ATGCCGCAAA
 70 GAGAATAGTG TATGCGGCGA AAAGTGCTCA GTTTCGATGA AGCAAAAACA GGAAGGCAAA ATGCCGCAAA
 CAGAACTTTA AAAGTGCTCA GTTTCGATGA AGCAAAAACA GGAAGGCAAA ATGCCGCAAA ATGCCGCAAA
 TTGAGATCCA GTTTCGATGA AGCAAAAACA GGAAGGCAAA ATGCCGCAAA ATGCCGCAAA ATGCCGCAAA
 TTTCTGGGTG AGCAAAAACA GGAAGGCAAA ATGCCGCAAA ATGCCGCAAA ATGCCGCAAA ATGCCGCAAA
 AATACTCATA CTCTTCTTTT TTCAATATTA AAATAAACAA ATAGGGGTTT TAACCTATAA CTGACACATG
 75 ATATTTGAAT GTATTTAGAA AACCATTATT ATCATGACAT GTGAAAACCT CAGGCGCGGT TGAGAGTGCA
 ACGTCTAAGA AACCATTATT ATCATGACAT GTGAAAACCT CAGGCGCGGT TGAGAGTGCA AGGCGCGGTG
 CGCGCGTTTC GGTGATGACG CCGGGAGCAG CAGATTGTAC TACTAGGTTG AGTTCATAGC AACGACCCCT
 TAAGCGGATG CCGGGAGCAG CAGATTGTAC TACTAGGTTG AGTTCATAGC AACGACCCCT TACTAGGTTG
 80 TTAACATATG GGCATCAGAG AGCAGCCAG CCGGGTCAAT CTGACCGCCC GACGTCATG GACGTCATG
 TGCAATAGGA AGCAGCCAG CCGGGTCAAT CTGACCGCCC GACGTCATG GACGTCATG GACGTCATG
 TAATCAATTA CCGGGTCAAT CTGACCGCCC GACGTCATG GACGTCATG GACGTCATG GACGTCATG
 85 GCCCGCCTGG CTGACCGCCC GACGTCATG GACGTCATG GACGTCATG GACGTCATG GACGTCATG
 GCCAATAGGG ACTTTCCATT ATATGCCAAG GACCTTATGG TTTTGGCAGT ACATCAATGG GAGTTTGT
 CAAGTGTATC ATATGCCAAG GACCTTATGG TTTTGGCAGT ACATCAATGG GAGTTTGT ATGGGCGGTA
 CCCAGTACAT GACCTTATGG TTTTGGCAGT ACATCAATGG GAGTTTGT ATGGGCGGTA TTATCGAAAT
 GGTGATGCGG TTTTGGCAGT ACATCAATGG GAGTTTGT ATGGGCGGTA TTATCGAAAT TAATACGACT
 CACCCCATTTG ACGTCAATGG ATTGACGCAA GAACCCACTG CTTAACTGGC TTATCGAAAT TAATACGACT
 ACTCCGCCCC ATTGACGCAA GAACCCACTG CTTAACTGGC TTATCGAAAT TAATACGACT CACTATAGGG
 GCTAACTAGA GAACCCACTG CTTAACTGGC TTATCGAAAT TAATACGACT CACTATAGGG AGACCGGAAG CTTGAATTC

pcDNA3-E7-Hsp70

SEQ ID NO:18

The E7-Hsp70 fusion sequence is shown in bold, caps

1	gacggatcgg	gagtcctccc	gatacccttat	ggctgactct	cagtacaatc	tgctctgatg	ccgcatagtt	aagccagtat	80
81	ctgtccctcg	ctgtgtgtgt	ggaggtcgct	gagtagtgcg	cgagcaaaat	ttaaagctaca	acaaggcaag	gcttgaccga	160
161	caattgcctg	aagaatctgc	ttagggtttag	gcgttttgcg	ctgcttcgcg	atgtacggcg	cagatatagc	cggtgacatt	240
241	gattattgac	tagttatttaa	tagtaatacaa	ttacgggggc	attagttcat	agcccatata	tggagttccg	cgttacataa	320
321	cttacggtaa	atggccccgc	tggtcgaccg	cccaacgacc	ccggcccat	gacgtcaata	atgacgtatg	ttcccatagt	400
401	aacgccaata	gggactttcc	attgacgtca	atgggtggag	tatttacggt	aaactggccc	atgggcagta	catcaagtgt	480
481	atcatatgcc	aagtacgccc	cctattgacg	tcaatgacgg	taaaatggccc	gcttggcatt	atggcccagta	catgacctta	560
561	tgggactttc	ctacttggca	gtacatctac	gtattagtca	tcgtattac	catggtgatg	cggttttggc	agttacatcaa	640
641	tgggcgtgga	tagcggtttg	actcacgggg	atttccaaat	ctccacccca	ttgacgtcaa	tgggagtttg	ttttggcacc	720
721	aaaatcaacg	ggactttcca	aaatgtcgta	acaactccgc	cccatggacg	caaatggggc	gtaggcgtgt	acgggtgggag	800
801	gtctatataa	gcagagctct	ctggctaact	agagaaccca	ctgcttactg	gcttatcgaa	attaatacga	ctcactatag	880
881	ggagacccaa	gctgggctagc	gttttaaacgg	gccctctaga	ctcgagcggc	cgccactgtg	ctggatatct	gcagaattcc	960
961	accacactgg	actagtggtat	ccatgcatgg	AGATACACCT	ACATTGCATG	AAATATATGT	AGATTGGCAA	CCAGAGACAA	1040
1041	CTGATCTCTA	CTGTTATGAG	CAATTAATG	ACAGCTCAGA	GGAGGAGGAT	GAAATAGATG	GTCCAGCTGG	ACAAGCAGAA	1120
1121	CCGGACACAG	CCCATTACAA	TATTGTAACC	TTTTGTTGCA	AGTGTGACTC	TACGCTTCGG	TTGTGCGTAC	AAAGCACACA	1200
1201	CGTAGACATT	CGTACTTTGG	AAGACCTGTT	AATGGGCACA	TAGGAATATG	TGTGCCCAT	CTGTTCTCAA	GGATCCATGG	1280
1281	CTCGTGCGET	CGGGATCGAC	CTCGGGACCA	CCAACTCCGT	CGTCTCGGTT	CTGGAAGGTG	GCGACCCCGT	CGTCGTCGCC	1360
1361	AACTCCGAGG	GCTCCAGGAC	CACCCCGTCA	ATTGTCCGCT	TCGCCCGCAA	CGGTAGAGTG	CTGGTCGGCC	AGCCCGCCAA	1440
1441	GAACCAAGCA	GTGACCAACG	TCGATCGCAC	CGTGCCTCG	GTCAAGCGAC	ACATGAGGCG	CGACTGGTCC	ATAGAGATTG	1520
1521	ACGGCAAGAA	ATACACCCGC	CCGGAGATCA	CGCCCCGCT	TCTGATGAAG	CTGAAGCGCG	ACGCCGAGGC	CTACTCGGT	1600
1601	GAGGACATTA	CCGACGCGGT	TATCACGACG	CCCGCTACT	TCAATGACGC	CCAGCGTCAG	GCCACCAAGG	ACGCCGGCCA	1680
1681	GATCGCGCGC	CTCAACGTGC	TGCGGATCGT	CAACGAGCGG	ACCGCGGCCG	CGCTGGCCTA	CGGCCCTCGA	AAGGGCGAGA	1760
1761	AGGAGCAGCG	AATCCTGGTC	TTGCACTTGG	GTGGTGGCAC	TTTCGACGTT	TCCCTGCTGG	AGATCGGCCA	GGGTGTGGTT	1840
1841	GAGTCCGTG	CCACTTCGGG	TGACAACCAC	TCGATCTGAC	CAGTCCACCT	CCAGCGGGTC	GTCGATTGGC	TGGTGGACAA	1920
1921	GTTCAGGGC	ACCAGCGGCA	AGTTCGAGT	GAGTCCAGCT	CAGTCCAGCT	CCAGCGCTGC	GGAAGCCGCC	GAGAAGGCAA	2000
2001	AGATCGAGCT	GAGTTCGAGT	AGCTGACCCG	AGCTGACCCG	AGCTGACCCG	AGCTGACCCG	CCGACCAAGAA	CCCGTTGTTT	2080
2081	TTAGACGAGC	AGCTGACCCG	GACACCGGCA	TTTCGGTGTG	GGAGATCGAT	CACGTTGTGC	GTCAACCCCG	ATGAGGTTGT	2160
2161	GGTGATCGCT	GACACCGGCA	TTTCGGTGTG	GGAGATCGAT	CACGTTGTGC	GTCAACCCCG	ATGAGGTTGT	CGCGGTGGGA	2240
2241	TGACCGATCT	GGTCAAGGAA	CTCACCGGCG	CCTCAAGGGC	GAGGTGAAAG	ACGTTCTGCT	ACCCGCTGTA	GCCTGGGTAT	2320
2321	GCCGCTCTGC	AGGCCCGGCG	CTCACCGGCG	CCTCAAGGGC	GAGGTGAAAG	ACGTTCTGCT	ACCCGCTGTA	GCCTGGGTAT	2400
2401	CGAGACCAAG	GCGGGGGTGA	TGACCGAGCT	CATCAGCGCG	AACACCCACG	TCCCAACCAA	GCGTGGGAG	ACTTTCACCA	2480
2481	CCGCGGACGA	CAACCAACCG	CTCGGTCAGA	TCCAGGTCTA	TCAGGGGGAG	CGTGAGATCG	CCGCGCACAA	CAAGTTGCTC	2560
2561	GGGTCTCTCG	AGCTGACCCG	CATCCCGCGG	CGCCCGCGGG	GGATTCCGCA	GATCGAGGTC	ACTTTCGACA	TCGACGCCAA	2640
2641	CGGCATTGTG	CAGTGCACCG	CCAAGGACAA	GGGCACCGGC	AAGGAGAACA	CGATCCGAAT	CCAGGAAGGC	TCGGGCCCTGT	2720
2721	CCAAGGAAGA	CATTGACCCG	ATGATCAAGG	AGCCCGAAGC	GCACGCCGAG	GAGGATCGCA	AGCGTCGCGA	GGAGGCCGAT	2800
2801	GTTCGTAATC	AAGCCGAGAC	ATTGGTCTAC	CAGACGGAGA	AGTTCGTCAA	AGAACAGCGT	GAGGCCGAGG	GTGGTTCGAA	2880
2881	GGTACCTGAA	GACACGCTGA	ACAAGTTTGA	TGCCCGGGTG	CGGGAAGCGA	AGGCGGCACCT	TGGCGGATCG	GATATTTCCG	2960
2961	CCATCAAGTC	GGCGATGGAG	AAGCTGGGCC	AGGATCGCA	GGCTCTGGG	CAAGCGATCT	ACGAAGCAGC	TCAGGCTGCG	3040
3041	TCACAGGCCA	CTGGCGCTGC	CCACCCCGGC	TCGGCTGATG	AAAGCTTaaag	tttaaacccgc	tgatagcct	cgactgtgcc	3120

3121	ttctagttgc	cagccatctg	ttgtttgccc	ctcccccggt	ccttccttga	ccctggaagg	tgccactccc	actgtccttt	3200
3201	cctaataaaa	tgaggaaatt	gcatcgcaat	gtctgagtag	gtgtcattct	attctggggg	gtggggtggg	gcaggacagc	3280
3281	aagggggagc	atgggaaga	caatagcagg	catgctgggg	atgcggtggg	ctctatggct	tctgagggcg	aaagaaccag	3360
3361	ctggggctct	agggggatc	ccacgcgcc	ctgtagcggc	gcattaagcg	cggcggtgtg	ggtggttacg	cgcagcgtga	3440
3441	ccgctacact	tgccagcgg	ctagcgccc	ctcctttcgc	ttcttccct	tcctttctcg	ccacgttcgc	cggctttccc	3520
3521	cgtcaagctc	taaatcggg	catccttta	gggttccgag	ttagtctttt	acggacactc	gaccccaaaa	aacttgatta	3600
3601	gggtgatgg	tcacgtagtg	ggccatcgcc	ctgatagacg	gttttttcgc	ctttgacgtt	ggagtccacg	ttctttaata	3680
3681	gtggactctt	gttccaaact	ggaacaacac	tcaaccctat	ctcggctctat	tccttttgatt	tataagggat	tttgggggatt	3760
3761	tcggcctatt	gggttaaaaa	tgagctgatt	taacaaaaat	ttaacgcgaa	ttaattctgt	ggaatgtgtg	tcagtttaggg	3840
3841	tgtggaaagt	ccccaggctc	cccaggcagg	cagaagtatg	caaaagcatg	atctcaatta	gcagcaacc	aggtgtggaa	3920
3921	agtccccagg	ctccccagca	ggcagaagta	tgcaaaagcat	gcacttcaat	tagtcagcaa	ccatagtccc	gcccctaact	4000
4001	ccgcccctcc	cgcccctaac	tcggcccagt	tccgcccatt	ctccgcccc	tggtgacta	attttttta	tttatgcaga	4080
4081	ggccgaggcc	gcctctgcct	ctgagctatt	ccagaagttag	tgaggaggct	tttttggagg	cctaggcttt	tgcaaaaagc	4160
4161	tcccgggagc	ttgtatatcc	attttcggat	ctgatcaaga	gacaggatga	ggatcgtttc	gcatgattga	acaagatgga	4240
4241	ttgcacgcag	gttctccggc	cgcttgggtg	gagaggctat	tcggctatga	ctgggcacaa	cagacaatcg	gctgctctga	4320
4321	tgccgcgctg	ttccggctgt	cagcgacagg	gcccgcggtt	ctttttgtca	agacgcacct	gtccggtgcc	ctgaatgaac	4400
4401	tgccaggacga	ggcagcgccg	ctatcgtggc	tgcccgcgac	ggggttctct	tgcgacgtg	tgctgcactg	tgctcactgaa	4480
4481	gcgggaaagg	actggctgct	attggggcgaa	gtgcccgggc	aggatctcct	gtcatctaac	ctgtcctctg	ccgagaaaagt	4560
4561	atccatcatg	gctgatgcaa	tgccggcggt	gcatacgttt	ggtccggcta	cctgcccact	cgaccaccaa	gcgaaaacatc	4640
4641	gcacgcagcg	agcacgtact	cggtggaag	ccggtcttgt	cgatcaggat	gatctggacg	aagagcatca	ggggctcgcg	4720
4721	ccagccgaac	tggttcgccg	gctcaaggcg	cgcatacgcc	acggcgagga	tctcgtcgtg	acctatggcg	atgcctgctt	4800
4801	gccgaatc	atggtggaaa	atggccgctt	ttctggattc	atcgactgtg	gcggctggg	tgtggcgac	cgctatcagg	4880
4881	acatagcggt	ggctaccgtg	gatatgtcgt	aagagcttgg	cggcgaaatg	gctgaccgct	ttctcgtgct	ttacgggtatc	4960
4961	gcgcgtcccg	atcgcagcg	catcgccctc	tatcgccctc	ttgacgagtt	cgtcttctat	gaaaggttgg	gcttcggaat	5040
5041	accgaccaa	cgacgccc	cctgccaatc	ccagcgccg	gatctcatgc	tggaagtctt	cgccccccc	aacttgttta	5120
5121	cgttttccgg	gacgcggct	gaatgatcct	ccagcgccg	gatctcatgc	aatattcaca	ttttttcact	gcattctagt	5200
5201	ttgcagctta	taatggttac	aaataaagca	atagcatcac	aaatttcaca	gacctctagc	tagagcttgg	cgtaatcatg	5280
5281	tgtggtttgt	ccaaactcat	caatgtatct	tccgctcaca	gttgcgctca	ctgccccgct	tcagtcgagc	aaagtataag	5360
5361	gtcatagctg	tttccctgtg	gaaattgtta	cattaattgc	gttgcgctca	ctgccccgct	tcagtcgagc	aaagtataag	5440
5441	cctgggggtgc	ctaagtgtg	agctaactca	ccagcgccg	gcaaaaagcc	aggaaccgta	aaaaggccgc	gttgctggcg	5520
5521	tgccagctgc	attaatgaat	cgctcggtcg	ccctggaag	ctcctcgtg	ttccgacgag	ggcgaaacc	cgacaggact	5600
5601	tgaactcgctg	cgctcggtcg	cgcccaacgc	ccctggaag	ccctcgtg	ttccgacgag	ggcgaaacc	cgacaggact	5680
5681	caggggataa	cgcaggaaa	agcatgtgag	caaaaagcca	atcgacgctc	agtcagagag	tgccgaaacc	cgacaggact	5760
5761	ttttccata	ggctccgccc	ccctggaag	ccctcgtg	ttccgacgag	ggcgaaacc	cgacaggact	cgacaggact	5840
5841	ataaagatac	cagcgcttcc	ccctggaag	ccctcgtg	ttccgacgag	ggcgaaacc	cgacaggact	cgacaggact	5920
5921	cgcccttctc	cccttccgga	agcgtggcgc	tttctcaatg	ctcagcgtgt	aggtatctca	gttcgggtga	ggtcgttcgc	6000
6001	tccaagctgg	cctgtgtgca	cgcaaccccc	ggtaagcccc	accgtggcg	cttatccggt	aactatcgtc	ttgagtcaca	6080
6081	ccgggtaaga	cacgacttat	cgccactggc	agcagccact	ggtaacagga	ttagcagagc	gaggtatgta	ggcggtgcta	6160
6161	cagagttctt	gaagtgttgg	cctaactacg	gctacactag	aaggacagta	tttggatctc	ggcgtctgct	gaagccagtt	6240
6241	accttcggaa	aaagagttgg	tagctcttga	tcgggcaaac	aaaccacgcg	tggttagcgtt	ggtttttttg	tttgcagaca	6320
6321	gcagattacg	cgcagaaaaa	aaggatctca	agaagatcct	ttgatctttt	ctacgggggtc	tgacgctcag	tggaaacgaaa	6400
6401	actcacgtta	agggattttg	gtcatgagat	tatcaaaaag	gatcttccac	tagatccttt	taaaataaaa	atgaagtttt	6480
6481	aaatcaatct	aaagtatata	tgagtaaaact	tggtctgaca	gttaccacatg	cttaatacagt	gaggcaccta	tctcagcgat	6560

6561	ctgtctattt	cgttcatcca	tagttgcctg	actcccgc	gtgtagataa	ctacgatacg	ggaggggctta	ccatctggcc	6640
6641	ccagtgtgc	aatgataccg	cgagaccac	gctcacggc	tccagattta	tcagcaataa	accagccagc	cggaaggcc	6720
6721	gagcgaag	gtgttcctgc	aactttatcc	gcctccatcc	agtcatttaa	ttgttgccgg	gaagctagag	taagtagttc	6800
6801	gccagttaat	agtttgccga	acgtttgtgc	cattgctaca	ggcatcgtgg	tgtaacgctc	gtcgtttggt	atggcttcat	6880
6881	tcagtcocgg	ttcccaacga	tcaaggcgag	ttacatgac	ccccatgttg	tgcaaaaaag	cggtttagctc	cttcgggtcct	6960
6961	ccagtcgttg	tcagaagtaa	gttgcccgca	gtgttatcac	tcattggttat	ggcagcactg	cataattctc	ttactgtcat	7040
7041	gccatccgta	agatgctttt	ctgtgactgg	tgaagtaaca	accaagtcac	tctgagaata	gtgtatggg	cgaccgagtt	7120
7121	gctcttgccc	ggcgtcaata	cggtacaata	cgcgcacac	tagcagaact	ttaaaaagtc	tcatactgg	aaaacgttct	7200
7201	tcggggcgaa	aactctcaag	gatcttaccg	ctgttgagat	ccagttcgat	gtaaccacac	cgtgcaccca	actgatcttc	7280
7281	agcatctttt	actttcacca	gcgtttctgg	gtgagcaaaa	acaggaaggc	aaaatgccgc	aaaaaaggga	ataaggcgga	7360
7361	cacggaaatg	ttgaatactc	atactcttcc	ttttcaata	ttattgaagc	atttatcagg	gttattgtct	catgagcgga	440
7441	tacataattg	aatgtattta	gaaaaataaa	caaatagggg	ttccgcgcac	atttccccga	aaagtgcac	ctgacgtc	7518

SINrep5-E7-Hsp70

SEQ ID NO:19 The E7-Hsp70 fusion sequence is shown in bold, caps

1	attgacggcg	tagtacacac	tattgaatca	aacagccgac	caattgcact	accatcacaa	tggagaagcc	agtagtaaac	80
81	gtagacgtag	accccagag	tccgtttgtc	gtgcaactgc	aaaaagctt	cccgaattt	gaggtagtag	cacagcaggt	160
161	cactccaaat	gaccatgcta	atgccacagc	cttttgcgat	ctggccagta	aactaatacg	gctggagggtt	cctaccacag	240
241	cgacgatctt	ggacatagcg	agcgcacccg	ctcgtagaaat	gttttcgcag	caccagatct	attgtgctgt	ccccatcgct	320
321	agtcacagaag	acccggacgc	catgatgaaa	tacgccagta	aactggcgga	aaaagcgtgc	aagattacaa	acaaagaactt	400
401	gcatgagaag	attaaggatc	tccggaccgt	acttgatcac	cgggatgctg	aaacaccatc	gctctgcttt	cacaacagctt	480
481	ttacctgcaa	catcgctggc	gaatatccgc	tcatgcagga	cgtgtatctc	aacgctcccg	gaactatcta	tcatacaggt	560
561	atgaaaaggcg	tgcggaccct	gtactggatt	ggcttcgaca	ccaccagtt	catgttctcg	gctatggcag	gttcgtaccc	640
641	tgcgtacaac	acaaattgtcg	ataatgagga	agaaggagtt	gaagcccggt	tcgcggggtt	atttctccgt	aggatcgaca	720
721	gtaggacagg	aaaattgtcg	ataatgagga	agaaggagtt	gaagcccggt	gttccacttg	aattggaaaagc	agtcgtacac	800
801	ctttatccag	aacacagagc	cagcttgtag	aggtctacga	ttccatcggt	tcaccatcag	tcccgggatc	acggagagaa	880
881	ttgcccgtgt	gatacagtg	tgagttgcga	aggtcttctt	gctatgcata	gttactgaca	cagtaaaagg	agaacgggta	960
961	ccgtgggata	cgcggttaca	cacaatagcg	acccatgcga	atcagatgac	tggtataatg	gcaacggata	tataccctga	1040
1041	togttccctg	tgtgcacgta	catccccgcc	accatagcgc	atcagatgac	gtgactaacg	gaggaacacc	aacaccatgc	1120
1121	cgatgcacaa	aaactttctg	ttgggctcaa	ccagcgaatt	gtcattaacg	gtaggactaa	caggaaacac	taacagagaaa	1200
1201	aaaattacct	tctgcagatc	atagcacaaag	gggttcagcaa	atgggctaaag	gagcgaaggg	atgatcttga	taacagagaaa	1280
1281	atgctgggta	ctagagaacg	caagcttacg	tatggctgct	tggtggcggtt	tcgcactaaag	aaagtacatt	cgtttttatcg	1360
1361	cccacctgga	acgcagacct	gcgtaaaaag	cccagcctct	tttagcgctt	ttcccatgct	gtccgtatgg	acgacctctt	1440
1441	tgccccatgtc	gctgagggcag	aaattgaaac	tgccattgca	accaaaagaa	gaggaataaac	tgctgcaggt	ctcggaggaa	1520
1521	ttagtcatgg	aggccaaggc	tgctttttgag	gatgctcagg	aggaagccag	agcggagaag	ctccgagaag	cacttccacc	1600
1601	attagtggga	gacaaaaggca	tcgaggcgagc	cgcagaagtt	gtctgcgaag	tggagggggt	ccaggcgagc	atcggagcag	1680
1681	cattagtgtg	aaccccgccg	ggtaacgttaa	ggataatacc	tcaagcaaat	gaccgtatga	tcggacagta	tatcgttgtc	1760
1761	tcgccaaact	ctgtgctgaa	gaattgccaaa	ctcgaccacg	cgcaccgcct	agcagatcag	gttaagatca	taaacacactc	1840
1841	cggaaagatca	ggaaggtagc	cggtcgaaac	atacgacgct	aaagtactga	tgccagcagg	aggtgccgta	ccatggccag	1920
1921	aattccctagc	actgagtgag	agcggccacgt	tagtgtacaa	cgaagagagag	tttgtgaacc	gcaaacata	ccacattgcc	2000

2001	atgcattggcc	ccgccaaagaa	tacagaagag	gagcagtaca	aggttacaaa	ggcagagcct	gcagaaacag	agtactgtgt	2080
2081	tgactggag	aagaagcgtt	gcgttaagaa	ggaagaagcc	tcaggtctgg	tcctctcggg	agaactgacc	aacctccct	2160
2161	atcatgagct	agctctggag	ggactgaaga	ccgacctgc	ggtcccgtac	aaggtcgaaa	caataggagt	gataggcaca	2240
2241	ccgggtcgg	gcaagtcagc	tattatcaag	tcaactgtca	cgccacgaga	tcttggtacc	agcgaaagaa	aagaaaattg	2320
2321	tcgcgaatt	gagggcgagc	tgctaagact	gaggggtatg	cagattacgt	cgaagacagt	agattcgggt	atgctcaacg	2400
2401	gatgccaaa	agccgtagaa	gtgctgtacg	ttgacgaagc	gttcgcgtgc	cagcgaggag	cactacttgc	cttgattgct	2480
2481	atcgtcaggc	cccgcgaagaa	ggtagtacta	tgccgaagcc	ccatgcaatg	cggtattcttc	aacatgatgc	aactaaaggt	2560
2561	acatttcaat	caccctgaaa	aagacatata	caccaagaca	ttctacaaag	atctctcccg	gcgttgaca	cagccagtta	2640
2641	cagctattgt	atcgacactg	cattacatag	gaaagatgaa	atccacgaag	cgttgcaaga	agaacattga	aatcgatat	2720
2721	acagggggcca	caaagccgaa	gccaggggat	atcatcctga	catgtttccc	cggttggtgt	aagcaattgc	aaatcgacta	2800
2801	tcgccgacat	gaagtaatga	cagccgcggc	ctcacaaagg	ctaaccagaa	aaggagtgt	tgccgtccgg	caaaaagtca	2880
2881	atgaaaaccc	actgtacgcg	atcacatcag	agcatgtgaa	cgtgttgctc	accgcactg	aggacaggt	agtgtgaaa	2960
2961	accttgtagg	gcgacccatg	gattaaagcag	ccacttaaca	tacctaaagg	aaactttcag	gctactatag	aggactggga	3040
3041	agctgaacac	aagggataaa	ttgctgcaat	aaacagcccc	actccccgtg	ccaatccgtt	cagctgaaag	accaacgttt	3120
3121	gctggggcgaa	agcattggaa	ccgatactag	ccacggccgg	tatcgtactt	accggttgcc	agtggagcga	actgttccca	3200
3201	cagtttgccg	atgacaaaac	acattcggcc	atttaacgct	tagacgtaat	ttgcattaa	tttttcggca	tggacttgac	3280
3281	aagcggactg	ttttctaacc	agagcatccc	actaacgtac	catcccgcg	attcagcgag	gccggtagct	cattgggaca	3360
3361	acagccaggg	aacgcgaag	tatgggttacg	atcagggcat	tgccgcgaa	ctctcccgt	gatttcgggt	gttccagcta	3440
3441	gctgggaagg	gcacacaact	tgatttgacg	acggggagaa	ccagcgttaa	ctctgcacag	cataaactgg	tccgggtgaa	3520
3521	ccgcaatctt	cctcacgcct	tagtccccga	gtacaaggag	agcaaacctg	gcccgttcaa	aaattctttg	aaccagtcca	3600
3601	aacaccactc	agtactttgt	gtatcagag	aaaaaattga	agctccccgt	aagagaatcg	aatggatcgc	cccgattggc	3680
3681	atagccgggtg	cagataagaa	ctacaacctg	gctttcgggt	ttccgcgcga	ggcacggtac	gacctgggt	tcatacaact	3760
3761	tggaaactaaa	tacagaaacc	accactttca	gcagtgcgaa	gacctgcgg	cgaccttaaa	aaccttttcg	cgttcggccc	3840
3841	tgaattgcct	taaccacgga	ggcaccctcg	tggtagaagtc	ctatggctac	ctcaagcaat	acagaaatgt	acctgatttt	3920
3921	gctcttgcca	gaaagtgtgt	caggggtgtct	gcagcgagac	cagattgtgt	attgcgtgat	ttcgtccgtg	tatgagggt	4000
4001	ccgacaacta	gacaacagcc	gtacacggca	attcaccccc	caccatctga	attgctgact	gtcaagagga	agcagttgtc	4080
4081	caagagatgg	agttggagcc	gcgccgtcat	accgcaccac	aaggggagaat	attgctgact	gtcaagagga	agcagttgtc	4160
4161	aacgcagcca	atccgctggg	tagaccaggg	gaaggagtct	gccgtgccat	ctataaacgt	tgcccgacca	gtttaccga	4240
4241	ttcagccacg	gagacaggca	ccgcaagaat	gactgtgtgc	ctaggaaaga	aagtgatcca	cgcggtcggc	cctgatttcc	4320
4321	ggaagcaccc	agaagcagaa	gccttgaaat	tgctacaaaa	cgctaccat	gcaagtggcag	acttagtaaa	tgaacataac	4400
4401	atcaagtctg	tcgccattcc	actgctatct	acaggcatct	acgcagccgg	aaaagaccgc	cttgaagtat	cacttaactg	4480
4481	cttgacaacc	gcgctagaca	gaactgacgc	ggacgttaacc	atctattgcc	tggataagaa	gtggaaaggaa	agaatcgacg	4560
4561	cggcactcca	acttaaggag	tctgtaacag	agctgaaggga	tgaagatatg	gagatcgacg	atgagttagt	atggatttcat	4640
4641	ccagacagtt	gcttgaaagg	aagaaaggga	ttcagttacta	caaaaggaaa	attgtatttcg	tacttcgaag	gcaccaaat	4720
4721	ccatcaagca	gcaaaagaca	tggcggagat	aaaggtctctg	ttccctaag	accaggaaag	taatgaacaa	ctgtgtgctt	4800
4801	acatatggg	tgagaccatg	gaagcaatcc	gcgaaaagtg	cccggtcgac	cataaacctg	cgtctagccc	gccccaaacg	4880
4881	ttgccgtgcc	tttgcatgta	tgccatgacg	ccgaaaaggg	ttcacagact	tagaagcaat	aacgtcaaa	aagttacagt	4960
4961	atgctcctcc	accccccttc	ctaagcacaa	aattaagaat	gttcagaagg	ttcagtgac	gaaagtatct	ctgtttaatc	5040
5041	cgcacactcc	cgcattcgtt	ccgcctcgta	agttacataga	agtgccagaa	cagcctacgg	ctcctcctgc	acaggccgag	5120
5121	gagggccccc	agttgttagc	gacaccgtca	ccatctacag	ctgataacac	ctcgtttgat	gtcacagaca	tctcactgga	5200
5201	tatggatgac	agtagcgaag	gctcactttt	ttcgagctttt	agcggtatcg	acaactctat	tactagtatg	gacagttgggt	5280
5281	cgtcaggacc	tagttcacta	gagatagtag	accgaaggca	gggtgggtgtg	gctgacgttc	atgcgctcca	agagcctgcc	5360
5361	cctattccac	cgccaaggct	aaagaagatg	gcccgccctgg	cagcggcgaag	aaaagagccc	actccaccgg	caagcaatag	5440

5441	ctctgagtc	ctccacctct	cttttgggtg	ggtatccatg	tcctctggat	caattttcga	cggagagacg	gcccgcag	5520
5521	cagcgtaca	acccctggca	acagggccca	cggatgtg	tatgtctttc	gcatgtttt	ccgacggaga	gattgatgag	5600
5601	ctgacccga	gagtaactga	gtccgaaccc	gtccctgtttg	gatcatttga	accggcgaa	gtgaactcaa	ttatatcgtc	5680
5681	cagatcagc	gtatcttttc	cactacgcaa	gcagagacgt	agacgcagga	gcagagagac	tgaatactga	ctaaccgggg	5760
5761	taggtgggtg	catatctttc	acggacacag	gcccctggga	cttgcaaaag	aagtccgttc	tcgagaacca	gcttacagaa	5840
5841	ccgaccttgg	agcgcaatgt	cctggaaaga	attcatgccc	cgggtctcga	cagctcga	gaggaacca	tcaaaactcag	5920
5921	gtaccagatg	atgcccccg	aagccaacaa	aagtaggtac	cagctctcga	aagtagaaaa	tcagaaagc	ataaccactg	6000
6001	agcgactact	gtcaggacta	cgactgtata	actctgccac	agatcagcca	gaatgtctata	agataactc	tcgaaacca	6080
6081	ttgtactcca	gtagcgtacc	ggcgaactac	tcggtaccac	agttcgctgt	agctgtctgt	aacaactatc	tgcatgagaa	6160
6161	ctatccgaca	gtagcatctt	atcagattac	tgacgagtag	gatgcttact	tggatatggt	agacgggaca	gtcgccctgc	6240
6241	tggatactgc	aaccttctgc	cccgtaagc	ttagaagtta	ccgaaaaaaa	catgagtata	gagccccgaa	tatccgcagt	6320
6321	gcggttccat	cagcgatgca	gaacacgcta	caaaatgtgc	tcattgccc	actaaaaa	aattgcaacg	tcacgcagat	6400
6401	gcgtgaactg	ccaacactgg	actcagcgac	attcaatgtc	gaatgctttc	gaaaatatgc	atgtaatgac	gagtattggg	6480
6481	aggagtctgc	tcggaagcca	attaggatta	ccactgagtt	tgtcacccga	tatgtagcta	gactgaaagg	ccctaaggcc	6560
6561	gccgactat	ttgcaaaagc	gtataatttg	gtcccattgc	aagaagtgc	tatggataga	ttcgtcatgg	acatgaaaag	6640
6641	agacgtgaaa	gttacaccag	gcacgaacaa	attcaccggg	agaccgaaag	tacaagtgat	acaagccgca	gaacccctgg	6720
6721	cgactgctta	cttatgcggg	ttttgtagca	atcatagcag	taggcttacc	gocgtcttgc	ttccaaact	tcacacgctt	6800
6801	tttgacatgt	cggcgaggga	ttttgtagca	atcatagcag	aacacttcaa	gcaaggcgac	ccggtactgg	agacggatat	6880
6881	cgcatacttc	gacaaaagcc	aagacgacgc	tatggcggtta	accggtctga	tgatcttggg	ggacttgggt	gtggatcaac	6960
6961	caactactga	cttgatcag	tgcccttttg	gagaatatc	atccacccat	ctacctacgg	tactctgttt	taaattcggg	7040
7041	gcgatgatga	aatccggga	gttctctaca	ctttttgtca	acacagtttt	gaatgtcgtt	atcgccagca	gagtactaga	7120
7121	agagcggcct	aaaacgtcca	gatgtgcagc	gttcaattggc	gacgacaaca	tcatacatgg	agtagtatct	gacaaaagaa	7200
7201	tggtgtgagag	gtgcgccacc	tggtctcaaca	tgagaggttaa	gacatcagac	gcagtcacgc	gtgagagacc	accttacttc	7280
7281	tgcggcggat	ttatcttgca	agattcgggt	acttccacag	cgtgcccgt	ggcggaatccc	ctgaaaaggc	tggttaagtt	7360
7361	gggtaaaccc	ctcccagccg	acgacgagca	agacgaagac	agaagacgcg	ctctgctaga	tgaacaaaag	gcgtggttta	7440
7441	gagtaggtat	aacaggcaat	ttagcagtg	cogtgacgac	ccggtatgag	gtagacaata	ttacacctgt	ccctactggca	7520
7521	ttgagaactt	ttgcccagag	caaaagagca	ttccaaagcca	tcagagggga	aataaagcat	ctctacgggtg	gtccctaaata	7600
7601	gtcagcatag	tacatttcat	ctgactaata	ctacaacacc	accacctcta	gactcgagcg	gccgccactg	tgctggatat	7680
7681	ctgcagaatt	ccaccacact	ggactagtgg	atccatgcac	ggagatatca	ctacatttgc	tgatgagac	accttacttc	7760
7761	AACGAGAGAC	AACGATCTC	TACTGTTATG	AGCAATTAA	TGACAGCTCA	GAGGAGGAGG	ATGAAATAGA	TGGTCCAGCT	7840
7841	GGACAAGCAG	AACGGGACAG	AGCCCATTTAC	AATATTGTAA	CCTTTTGTG	CAAGTGTGAC	TCTACGCTTC	GGTTGTGCGT	7920
7921	ACAAAGCACA	CACGTAGACA	TTCTGACTTT	GGAAAGACCTG	TAAATGGGCA	CACATAGGAAT	TGTTGCCCCC	ATCTGTTCTC	8000
8001	AAGGATCCAT	GGCTGTCGG	GTCGGGATCG	ACCACCCCGT	CACCAACTCC	GTCGTCTCGG	TTCTGGAAGG	TGGCGACCCG	8080
8081	GTCGTCTGTCG	CCAACCTCGA	GGGCTCCAGG	ACCACCCCGT	CAATTGTGCG	GTTCCGCCCG	AACGTTGAGG	TGCTGTTGCG	8160
8161	CCAGCCCGCC	AAGAACCCAG	CAGTGACCAA	CGTCGATCGC	ACCGTGCGGT	CGGTCAAGCG	ACACATGGGC	AGCGACTGTT	8240
8241	CCATAGAGAT	TGACGGCAAG	AAATACACCG	CGCCGGAGAT	CAGCGCCCGC	ATTCTGATGA	AGCTGAAGCG	CGACGCCGAG	8320
8321	GCCTACCTCG	GTGAGGACAT	TACCGACGCG	GTTATACAGA	CGCCCGGCTA	CTTCAATGAC	GCCACGCTC	AGGCCACCAA	8400
8401	GGACGCCGCG	CAGATCGCCG	GCCTCAACGT	GCTGCGGATC	GTCAACGACG	CGACCCGCGC	CGCGCTGGCC	TACGGCTCG	8480
8481	ACAAAGGCGA	GAAGGAGCAG	CGAATCCCTGG	TCTTCGACTT	GGGTGGTGGC	ACTTTCGACG	TTTCCCTGCT	GGAGATCGGC	8560
8561	GAGGTGTGTCG	TTGAGGTCCG	TGCCACTTCG	GGTGACAACC	ACCTCGGCGG	CGACGACTGG	GACCAGCGGG	TCGTGATTTG	8640
8641	GCTGGTGGAC	AAGTTCGAAG	GCACCAGCGG	CATCGATCTG	ACCAAGGACA	AGATGGCGAT	GCAGCGGCTG	CGGGAAGCCG	8720
8721	CCGAGAAGGC	AAAGATCGAG	CTGAGTTCGA	GTCACTCCAC	CTCGATCAAC	CTGCCCTACA	TCACCGTCCA	CGCCGACAAG	8800
8801	AACCCGTTGT	TCTTAGACGA	GCAGCTGACC	CGCGCGGAGT	TCCAACGGAT	CACTCAGGAC	CTGCTGGACC	GCACCTCGCA	8880

8881	GCGCTTCCAG	TCGGTGATCG	CTGACACCCG	CATTTCGGTG	TCGAGATCG	ATCACGTTGT	GCTCGTGGGT	GGTTCGACCC	8960
8961	GGATGCCCGC	GGTGACCGAT	CTGGTCAAGG	AACTCACCCG	CGGCAAGGAA	CCCAACAAGG	CGGTCAACCC	CGATGAGGTT	9040
9041	GTCGGGTGGG	GAGCGGTCTT	GCAGGCCGGC	GTCTCTAAGG	GCGAGGTGAA	AGACGTTCTG	CTGCTTGATG	TTACCCCGCT	9120
9121	GAGCCTGGGT	ATCGAGACCA	AGGCGGGGGT	GATGACCAGG	CTCATCGAGC	GCAACACCA	GATCCCCACC	AAGCGGTCCG	9200
9201	AGACTTTTAC	CACGCCCGAC	GACAAACCA	CGTCGGTGCA	GATCCAGGTC	TATCAGGGGG	AGCGTGAGAT	CGCCGCGCAC	9280
9281	AACAAGTTGC	TCGGGTCTTT	CGAGTGACCC	GGCATCCCGC	CGGCGCCGGC	GGGGAATCCG	CAGATCGAGG	TCACTTTTCA	9360
9361	CATCGACGCC	AACGGCATTTG	TGCACGTGAC	CGCCAAGGAC	AAGGCGACCG	GCAAGGAGAA	CACGATCCGA	ATCCAGGAAG	9440
9441	GCTCGGGCCT	GTCCAAGGAA	GACATTGACC	GCATGTGCTT	GGACGCCGAA	GAAGTTCGTC	AGGAGGATCG	CAAGCGTCCG	9520
9521	GAGGAGGCCG	ATGTTCTGTA	TCAAGCCGAG	ACATTTGGTT	ACCAGACGGA	TGGCGGAAGC	GAAGCGGCA	CTTGCGGGAT	9600
9601	GGGTGGTTTC	AAGGTACCTG	AAGACACGCT	GAACAAGTTT	GATCGCGCGG	CAGGCTCTGG	GGCAAGCGAT	CTTGGCGGAT	9680
9681	CGGATATTTC	GGCCATCAAG	TCGGCGATGG	AGAACCTGGG	CCAGGAGTCC	GAGGCTCTGA	GGCAAGCGAT	CTTACGAAGCA	9760
9761	GCTCAGGCTG	CGTCACAGGC	CACCTGGCGCT	GCCACACCCG	GCTCGGCTGA	TGAAAGCTTA	agtttgtgag	catgcaggcc	9840
9841	ttggggccaa	tgatccgacc	agcaaaactc	gatgtacttc	cgaggaaactg	atgtgcataa	tgcatcaggc	tggtacatta	9920
9921	gatcccgct	taccggggc	aatatagcaa	cactaaaac	tcgatgtact	tcggaggaaag	cgcagtgcat	aatgctgcgc	10000
10001	agtgttgcca	cataaaccat	atattaacca	tttatctagc	ggacgccaac	aactcaatgt	atttctgagg	aagcgtggtg	10080
10081	cataatgcc	cgcagcgtct	gcataacttt	tattattctt	tttattaac	aacaaaattt	tgtttttaac	atttcaaaa	10160
10161	aaaaaaaaa	aaaaaaaaa	aaaaaaaaa	aagggaattc	ctcgattaat	taagcgccgc	ctcgagggga	attaattctt	10240
10241	gaagacgaaa	gggccaagtg	gcacttttgc	gggaaatgtg	cgcggaaccc	ctatttgttt	atttttctaa	atacattcaa	10320
10321	atatgtatcc	gctcatgaga	caataaccct	gataaatgct	tcaataatat	tgaaaaagga	agagtatgag	tattcaacat	10400
10401	ttcgtgtctg	cccttattcc	cttttttgcg	gcattttgct	ttcctgtttt	tgctcaccac	gaaacgctgg	tgaaagtaaa	10480
10481	agatgctgaa	gatcagttgg	gtgcagagat	gggtttacatc	gaactggatc	tcaacagcgc	taagatcctt	gagagtcttc	10560
10561	gcccgaaga	acgtttttcca	atgatagaca	cttttaaatg	tctgctaatg	ggcgcggtat	tatcctgtgt	tgacgcccgc	10640
10641	caagagcaac	tcggtcgccg	catacactat	tctcagaagt	acttggttga	gtactaccca	gtccaaacta	agcatcttac	10720
10721	ggatggcatg	acagtaagag	aattatgcag	tgctgacata	accatgagtg	ataacactgc	gccttgatcg	ttgggaaccc	10800
10801	cgatcggagg	accgaaggag	ctaaccgctt	tttgcacaa	catgggggat	catgtaactc	acaacgttgc	gcaaacattt	10880
10881	gagctgaatg	aagccatacc	aaacgacgag	cgtgacacca	cgatgctgtg	agcaatggca	tggaagtcga	ggaccacttc	10960
10961	aactggcgaa	ctacttactc	tagcttcccg	gcaacaatta	atagactgga	tggaagtcga	tggtctgcgc	tatcattgca	11040
11041	tgcgctcggc	ccttcgggt	ggctgggtta	ttgctgataa	atctggagcc	ggtagcgtg	ggtagcgtg	atgaacgaaa	11120
11121	gactgggggc	cagatggtaa	gcccctccgt	atcgtagtta	tctacacgac	gggagatcag	gcaactatgg	atgaacgaaa	11200
11201	tagacagatc	gctgagatag	gtccctccgt	gtgtagatc	tggttaactgt	cagaccaagt	ttactcata	atactttaga	11280
11281	ttgatttaaa	acttcatctt	taatttaaaa	gattaaagcat	tggttaactgt	tttgataatc	tcattgaccaa	aatcccttaa	11360
11361	cgtgagtttt	cgttccactg	agcgtcagac	ccgttagagg	gaagatcctt	atcttcttga	gactcttttt	ttctgcggt	11440
11441	aactctgctg	ttgcaaacaa	aaaaaccacc	gctaccagcg	agatcaaaag	tgccggatca	agagctacca	actctttttc	11520
11521	cgaaggtaac	tggtctcagc	agagcgcaga	taccaaaatac	gtgtcttcta	gttagcgtg	agtaggcca	ccacttcaag	11600
11601	aactctgtag	caccgcctac	atacctcgtc	ctgctaatacc	tggttaactgt	ggctgcgtgc	agtgccgata	agtcgtgtct	11680
11681	taccgggttg	gactcaagac	gatatgtacc	ggataaggcg	cagcggctgcg	gctgaacgcg	gggttctgtg	acacagccca	11760
11761	gcttggagcg	aacgaactac	accgaactga	gatacctaca	gcgtgagcat	tgagaaagcg	ccacgcttcc	cgaaggagga	11840
11841	aaggcgggaca	ggatccgggt	aagcggcagg	gtcggaaacag	gagagcgcac	gagggagcct	ccaggggggaa	acgcctggtta	11920
11921	tcctttatag	cctgtcgggt	ttcgccacct	ctgacttgag	cgtcgatttt	tgtagatcgc	gtcagggggg	cggagcctat	12000
12001	ggaaaaacgc	cagcaacgcg	agctcgtatg	gacatatgtg	cgttagaacg	cggtacaac	taatacataa	ccttatgtat	12080
12081	catacacata	cgatttaggg	gacactatag						12110

pSCA1-E7-Hsp70 SEQ ID NO:20 The E7-Hsp70 fusion sequence is shown in bold, caps

1	atggcggatg	10	tgtagacatac	20	acgacgcgcaa	30	aagattttgt	40	tccagctcct	50	gccacactccg	60	ctacgcgaga	70	gattaaccac	80
81	ccacgatggc		cgccaaagt		catgttgata		ttgaggctga		cagcccatc		atcaagtctt		tgcaagaagc		atttccgtcg	160
161	ttcgaaggtg		agtcattgca		ggtcacacca		aattgacctg		caaatgccag		agcattttcg		cacctggcta		ccaaattgat	240
241	cgagcaggag		actgacaaag		acacactcat		cttggtatc		ggcagtgccg		cttccaggag		aatgatgtct		acgcacaaa	320
321	accactcggt		atgcccctatg		cgacgcgag		aaatcccg		acgtgcgac		aggtctgat		agaaactggc		agcgccctcc	400
401	gggaaggtgc		tggtatagaga		gatcgagga		aaatcccg		acgtgcgac		cggtatccg		agccagagc		ctgaatctcc	480
481	taccttttgc		ctgcatacag		acgtcacgtg		tcgtacggca		gcccgaagt		ccgtatccg		ggacgtgtat		cctgtacatg	560
561	caccaacatc		gctgtaccat		caggtcgatg		aaggtgtcag		aacggcgat		tggtattgggt		ttgacaccac		ccggtttatg	640
641	tttgacgcgc		tagcaggcgc		gtatccaacc		tacgcccaaa		actgggccga		cgagcagggtg		ttacaggcca		ggaacatagg	720
721	actgtgtgca		gcatccttga		ctgagggaag		actcgcaaaa		ctgtccattc		tcgcgaagaa		gcaattgaaa		ccttgcgaca	800
801	cagtcattgt		gtaaacaatc		ctttacctgt		aggtgcgata		ccatcgatc		ctgaggagct		ggcacttacc		ctccgtattc	880
881	tatgtgcccc		ggcctgtacg		gtaaaacggt		aggtgtcat		tcccgtgatg		gcactacgtc		ttctctagt		agaaaatcac	960
961	cagacactgt		caaaggagaa		agagtctcat		gacagaggtc		gacacagctc		attgtaaggt		ttctctagt		tgcaagacca	1040
1041	atactagcga		cgacgctcac		accggaggac		gcacagaagt		tgtagtggtg		attgtaaggt		aggatagttg		aatgactggc	1120
1121	aacacagcga		aacactaaca		cgatgaagaa		ctatctgtct		ccgattgtg		ccgtcgcat		tagcaagtgg		tgaacggaag	1200
1201	acaaggcaga		ccttgatgat		gaaaaacctc		tggtgttccg		agagaggtca		cttactgtg		gctgcttgg		gcgagggaa	1280
1281	acaggggaag		tgcaacacct		cagccactgc		aatccagctc		agacaaatgt		gaaggtgcct		tcagagttaa		actcgttctg	1360
1361	catcccagag		ctatgttcta		caggcctcgc		gacgcgtcgt		cctctgcccc		tgtagtggtg		attgtaaggt		gagccagagc	1440
1441	gaggttaat		acctgtcttc		ccttaccacc		gctgtggaac		acctcgcagc		gcgttgaaag		tcacgcgaca		gctactactg	1520
1521	gtatcacgca		ggtgcagggtg		tcgtggaac		acctcgcagc		tgctcaagag		gcccctgtgc		tgacggcagg		gctcctactac	1600
1601	gaaattacgt		agttctgtcc		ccgcagaccg		tgctcaagag		gctcctactac		gacgcgagct		gagccagctg		gagccagctg	1680
1681	gaaattacgt		agttctgtcc		ccgcagaccg		tgctcaagag		gctcctactac		gacgcgagct		gagccagctg		gagccagctg	1760
1761	gaaattacgt		agttctgtcc		ccgcagaccg		tgctcaagag		gctcctactac		gacgcgagct		gagccagctg		gagccagctg	1840
1841	gaaattacgt		agttctgtcc		ccgcagaccg		tgctcaagag		gctcctactac		gacgcgagct		gagccagctg		gagccagctg	1920
1921	gaaattacgt		agttctgtcc		ccgcagaccg		tgctcaagag		gctcctactac		gacgcgagct		gagccagctg		gagccagctg	2000
2001	gaaattacgt		agttctgtcc		ccgcagaccg		tgctcaagag		gctcctactac		gacgcgagct		gagccagctg		gagccagctg	2080
2081	gaaattacgt		agttctgtcc		ccgcagaccg		tgctcaagag		gctcctactac		gacgcgagct		gagccagctg		gagccagctg	2160
2161	gaaattacgt		agttctgtcc		ccgcagaccg		tgctcaagag		gctcctactac		gacgcgagct		gagccagctg		gagccagctg	2240
2241	gaaattacgt		agttctgtcc		ccgcagaccg		tgctcaagag		gctcctactac		gacgcgagct		gagccagctg		gagccagctg	2320
2321	gaaattacgt		agttctgtcc		ccgcagaccg		tgctcaagag		gctcctactac		gacgcgagct		gagccagctg		gagccagctg	2400
2401	gaaattacgt		agttctgtcc		ccgcagaccg		tgctcaagag		gctcctactac		gacgcgagct		gagccagctg		gagccagctg	2480
2481	gaaattacgt		agttctgtcc		ccgcagaccg		tgctcaagag		gctcctactac		gacgcgagct		gagccagctg		gagccagctg	2560
2561	gaaattacgt		agttctgtcc		ccgcagaccg		tgctcaagag		gctcctactac		gacgcgagct		gagccagctg		gagccagctg	2640
2641	gaaattacgt		agttctgtcc		ccgcagaccg		tgctcaagag		gctcctactac		gacgcgagct		gagccagctg		gagccagctg	2720
2721	gaaattacgt		agttctgtcc		ccgcagaccg		tgctcaagag		gctcctactac		gacgcgagct		gagccagctg		gagccagctg	2800
2801	gaaattacgt		agttctgtcc		ccgcagaccg		tgctcaagag		gctcctactac		gacgcgagct		gagccagctg		gagccagctg	2880
2881	gaaattacgt		agttctgtcc		ccgcagaccg		tgctcaagag		gctcctactac		gacgcgagct		gagccagctg		gagccagctg	2960
2961	gaaattacgt		agttctgtcc		ccgcagaccg		tgctcaagag		gctcctactac		gacgcgagct		gagccagctg		gagccagctg	3040
3041	gaaattacgt		agttctgtcc		ccgcagaccg		tgctcaagag		gctcctactac		gacgcgagct		gagccagctg		gagccagctg	3120
3121	gaaattacgt		agttctgtcc		ccgcagaccg		tgctcaagag		gctcctactac		gacgcgagct		gagccagctg		gagccagctg	3200
3201	gaaattacgt		agttctgtcc		ccgcagaccg		tgctcaagag		gctcctactac		gacgcgagct		gagccagctg		gagccagctg	3280
3281	gaaattacgt		agttctgtcc		ccgcagaccg		tgctcaagag		gctcctactac		gacgcgagct		gagccagctg		gagccagctg	3360

3361	aaggatgtat	ggattcaatg	ccgcaacagc	tgccaggctg	gaagctagac	ataccttctt	gaaggggag	ttcctatcaa	tgccatacgg	3440
3441	gcagcagcc	agttatcgca	gaaagaaaa	tccaaccgct	ttctgtgctg	gacaatgtaa	ttcctatcaa	ttcctatcaa	ccgagggctg	3520
3521	ccgacgccc	tgggtgctga	gtacaagacg	gttaaaaggca	gtaggggtga	gtggctggtc	aataaagtaa	gaggggtacca		3600
3601	cgtctgtgctg	gtgagtgagt	acaacttggc	tttgccttga	cgcaggggtca	cttggtgtgc	accgtgaat	gtcacagggc		3680
3681	ccgatagggtg	ctacgacctg	agtttaggac	ttccggctga	cgcggcgagg	ttcagacttg	gggggtgtaa	cattcacacg		3760
3761	gaattcagaa	tccaccacta	ccagcagtg	gtcgaccacg	ccatgaagct	gcagatgctt	gggggtgtaa	cgtacagact		3840
3841	gctaaaaacc	ggcgcatct	tgatgagagc	ttacggatcc	gcccataaaa	tcagcgaagc	cgttggttcc	tccttaagca		3920
3921	gaaagtcttc	gtctgcaaga	gtgtgcgcc	cggattgtgt	caccagcaat	acagaagtgt	ttctgtgtt	ctccaacttt		4000
4001	gacaacggaa	agagaccctc	tacgtatcac	cagatgaata	ccaagctgag	tgccgtgtat	gcccggagaag	ccatgcacac		4080
4081	ggccgggtgt	gcaccatcct	acagagttaa	gagagcagc	atagccacgt	gcacagaagc	ggctgtggtt	aacgcagcta		4160
4161	acgcccgttg	caatgtaggg	gatggcgtag	gcaggccgct	ggcgaagaaa	tggcgttcag	cttttaagg	agcagaaca		4240
4241	ccagtgggca	aattaaaaac	agtcattgac	ggctcgtacc	ccgtcatcca	cgtgtagc	cctaatttct	ctgccacgac		4320
4321	tgaagcgga	ggggaccgag	aattggccg	tgctaccg	gcagtgccg	ccgaagttaa	ccctcaacca	ctgagcagcg		4400
4401	tagccatccc	gctgctgtcc	acaggagtgt	tcagcgccg	agagatagg	ctgcagcaat	ccctcaacca	tctattcaca		4480
4481	gcaatggagc	ccacggacgc	tgactcaatga	tgactgtgag	gagacaaaag	ttgggagaag	aaatccagg	aagccattga		4560
4561	catgagagc	gctgtggagt	tgactgtgac	tgactgtgag	ctgaccacag	acttgggtgag	agtgacccc	gacagcagcc		4640
4641	tgggtggctg	taagggtctac	gagcttgtag	acgggtcgct	gtactcgtac	tttgaaggtg	cgaaattcaa	ccaggctgct		4720
4721	attgatattg	cagagatact	gacgttgttg	ccgaactg	tccagctgac	ggaacagata	tgccatac	cgtgggga		4800
4801	aacaattgac	aacatcagat	gcaatgttcc	ggtagaagat	tccgacttcc	caacacttcc	gggagcag	ccctgctgt		4880
4881	gcccgtacgc	aatgacagca	gaacggatgc	cccgcttag	gtcacaccaa	gttaaaagca	tgggtgtt	ctcatctttt		4960
4961	ccccctccga	aataccatgt	agatggggtg	cagaaggtaa	agtgcgagaa	ggcttctcctg	ttcgaccgga	cggtaacctt		5040
5041	agtgtgtagt	ccgcggaagt	atgcccgcac	tacgacggac	cactcagatc	ggctgttacc	aggggttgac	ttggactgga		5120
5121	ccaccgactc	gtcttccact	gccagcgata	ccatgtcgt	accagttt	cagtcgtgtg	acatcgatc	gatctacgag		5200
5201	ccaatggctc	ccatagtagt	gacggctgac	gtcacactt	aaccgcgag	catcgcgagc	ctggcggcag	atgtgcacc		5280
5281	tgaaccgcga	gacctgtg	acctcgagaa	cccgactt	cccccgcg	ggaagagagc	tgcatactt	gcctcccgc		5360
5361	cggcgagcgc	accgtgtg	gcccgcagaa	agccgacgc	tgcccccaag	actcggtt	ggaacagct	gcctttgac		5440
5441	ttcggcgact	ttgacgagca	cgaagtcgat	gcgttggcct	ccgggattac	tttcggagac	ttcgacgagc	tcctgcgact		5520
5521	aggccgcgcg	ggtgcatata	ttttctctc	ggacactggc	agcggaat	tacaacaaa	atccgtagg	cagcacaatc		5600
5601	tccagtgcgc	acaactggat	gcggtccagg	aggagaaaat	gtacccgcca	aaattggata	ctgagagggg	gaagctgtt		5680
5681	ctgctgaata	tgcagatgca	cccatcgag	gctaataaga	gtcgatacca	gtctcgcaaa	gtggaagaca	tgaaagccac		5760
5761	ggtggtggac	aggctcacat	cgggggcccag	attgtacacg	ggagcggagc	taggcccgc	accacatac	gcggttcggt		5840
5841	acccccgcgc	cgtgtactcc	ctacacgtga	tgcgaagatt	ctcaagccc	gatgtagcaa	tcgcagcgtg	caacgaatac		5920
5921	ctatccagaa	attacccaac	agtggcgctg	taccagataa	cagatgaata	gcagcatac	ttggacatgg	ttgacgggtc		6000
6001	ggatagttgc	ttggacagag	gcacattctg	cccgcggaag	ctccggtgct	accgaaaca	tcattgctac	caccagcga		6080
6081	ctgtacgcag	tgccgtccc	tcaccccttc	agaacacact	acagaacgtg	ctagcgccg	ccaccaagag	aaactgcaac		6160
6161	gtcacgcaaa	tgcgagaact	accacacatg	gactcggcag	tgttcaacgt	ggagtgtctc	aaagcgtatg	cctgtctcgg		6240
6241	agaaatattg	gaagaatattg	ctaaacaacc	tatccggata	accactgaga	acatcactac	ctatgtgacc	aaattgaaag		6320
6321	gcccgaatgg	tgctgctt	ttcgttaaga	cccaacactt	ggttccgctg	caggaggttc	ccatggacag	attcacggtc		6400
6401	gacatgaac	gagatgtcaa	agtcactcca	ggagcgaacc	gaatagtaa	aagaccctaa	gtccaggtaa	ttcaagcagc		6480
6481	ggagcattg	gcgaccgtt	acctgtctga	ctccacacac	gatcatgcc	ggagactaaa	tgctgtgtta	gccttaacg		6560
6561	tgacacatt	gtttgatattg	tcggccgaag	actttgacgc	gatcatgcc	ccttggtctt	accagcagga	cccggttcta		6640
6641	gagacggaca	ttgcatcatt	cgacaaaagc	caggacgact	ccttggtctt	tacaggttta	atgatctctg	aagattctag		6720
6721	ggtggatcag	tacctgtg	acttgatcga	ggcagccttt	ggggaatat	ccagctgtca	cttacaact	ggcagcgtc		6800
6801	tcaagttcgg	agctatgatg	aaactcgga	tggttctgac	ttgtttttt	aacactgttt	tgaacatcac	catagcaagc		6880
6881	agggacttgg	agcagagact	cactgactcc	gcctgtgctg	cttcatcgg	cgacgacaac	atcgttcacg	gagtgatctc		6960
6961	cgacaagctg	atggcgagga	ggtgcgcgct	gtgggtcaac	atggaggtga	agatcattga	cgtgtgtcatg	ggcgaaaaac		7040

7041 ccccatattt ttgtggggga ttcatagttt ttgacagcgt cacacagacc gctgcccgtg tttcagacc acttaagcgc 7120
 7121 ctgttaagt tgggtaagcc gtaaacagct gaagacaagc aggacgaaga caggcgacga gcactgagt acgaggttag 7200
 7201 caagtgttc cggacaggtc tggggccga actggaggtg gactaacat gctagatga ggtagagggc tgcaaaagta 7280
 7281 tccatagc ctggtccacc ttggcaggg acattaaagc gtttaagaaa gtagagggac ctgtatata cctctacgac 7360
 7361 ggtcttagat ttgtgcgtta atacacagaa ttctgattgg atccatgcac ggagatgac ctacattgca tgaatatatg 7440
 7441 ttagatttgc aaccagagac aaactgatct tactgttatg agcaattaaa tgacagctca gaggagagg atgaatatga 7520
 7521 tgggtcagct ggacaagcag aaccggacag agcccatatc aatattgtaa cctttgtttg caagtgtgac tctacgcttc 7600
 7601 ggttggcgt acaaagcaca cactgagaca ttctgacttt ggaagacctg acctgggac caactatcc gtcgtctcgg ttctggagg 7680
 7681 atctgttctc aagatccat ggctcgtcgg ccaactccga gggctccagg gctcgacatg gctcgcccta cctcagcgtc 7760
 7761 tggcgacctg ccagcccggc aagaaccagg agactccagc agctgacatg cagcgccgca gctcgcccta cctcagcgtc 7840
 7841 tgggtgctgg ccagcccggc aagaaccagg agactccagc agctgacatg cagcgccgca gctcgcccta cctcagcgtc 7920
 7921 agcgaactgg ccataagat gctacactcg gctacactcg gctacactcg gctacactcg gctacactcg gctacactcg 8000
 8001 cgacgcccag gctacactcg gctacactcg gctacactcg gctacactcg gctacactcg gctacactcg gctacactcg 8080
 8081 agccacacaa ggacgcccgg cagatcgccg gtagggacat taccgacgg gctacactcg gctacactcg gctacactcg 8160
 8161 tacggcctcg caaaggcga gaaggagcag gaattcctgg gctacactcg gctacactcg gctacactcg gctacactcg 8240
 8241 ggagatcgcc gaggtgtgg tggaggtcgg tggaggtcgg tggaggtcgg tggaggtcgg tggaggtcgg tggaggtcgg 8320
 8321 tctgtcattg cctgtgtgac cctgtgtgac cctgtgtgac cctgtgtgac cctgtgtgac cctgtgtgac cctgtgtgac 8400
 8401 cgggaagccc cctgtgtgac cctgtgtgac cctgtgtgac cctgtgtgac cctgtgtgac cctgtgtgac cctgtgtgac 8480
 8481 cgccgacaag aaccggtgtg tctgtgtgac cctgtgtgac cctgtgtgac cctgtgtgac cctgtgtgac cctgtgtgac 8560
 8561 gcaactcgaa gccgttccag ggtgacccag ggtgacccag ggtgacccag ggtgacccag ggtgacccag ggtgacccag 8640
 8641 ggttgcaccc ggtgacccag ggtgacccag ggtgacccag ggtgacccag ggtgacccag ggtgacccag ggtgacccag 8720
 8721 ccatgaggtt gacgctgggt gacgctgggt gacgctgggt gacgctgggt gacgctgggt gacgctgggt gacgctgggt 8800
 8801 ttaccccgct gagcctgggt gagcctgggt gagcctgggt gagcctgggt gagcctgggt gagcctgggt gagcctgggt 8880
 8881 aagcgtcgga agactttcac cagcgtcgga agactttcac cagcgtcgga agactttcac cagcgtcgga agactttcac 8960
 8961 cgccgcccac acaatttgc cagcgtcgga agactttcac cagcgtcgga agactttcac cagcgtcgga agactttcac 9040
 9041 tcaatttgc cagcgtcgga agactttcac cagcgtcgga agactttcac cagcgtcgga agactttcac cagcgtcgga 9120
 9121 atccaggaag gctcggccct gctcggccct gctcggccct gctcggccct gctcggccct gctcggccct gctcggccct 9200
 9201 caagcgtcgc gagagggccc agactttcac cagcgtcgga agactttcac cagcgtcgga agactttcac cagcgtcgga 9280
 9281 gtgagggcca ggtgtgttc ggtgtgttc ggtgtgttc ggtgtgttc ggtgtgttc ggtgtgttc ggtgtgttc 9360
 9361 ctggcggtat cggatatttc ggtgtgttc ggtgtgttc ggtgtgttc ggtgtgttc ggtgtgttc ggtgtgttc 9440
 9441 ctacgaagca gctcagctg ttacatccct acgcaaacgt ttacgagcgt cgggtggcgc ccgcccgcgt tggccggtg 9520
 9521 attaatgaa ttacatccct acgcaaacgt ttacgagcgt cgggtggcgc ccgcccgcgt tggccggtg 9600
 9601 caggccactc cgtgtgttc cgtgtgttc cgtgtgttc cgtgtgttc cgtgtgttc cgtgtgttc cgtgtgttc 9680
 9681 gacaatgaga cagaacgcaa ttgtctctgc taggctctcc aaacaaaga agaaagac aacaaacca aagccgaaaa 9760
 9761 cgcagcccaa gaagatcaac ggaacaaacg agcagcaaaa gaagaaagc aagaagac aacaaacca aagccgaaaa 9840
 9841 ggaacaaagc aagaatgtg catgaagatt gaaatgact gtaatttctg taatttctg taatttctg taatttctg 9920
 9921 cagacatgac gggcaccgca ctatcatggg tgcagaaaa ctggtgtgtg ttggtgtgtg ttggtgtgtg ttggtgtgtg 10000
 10001 gtgtgtgtg ttgtcacttg ccaatggcat taaccataa cttgataatg taatttctg taatttctg taatttctg 10080
 10081 aacagaaaaa gttagggtaa gcaatggcat taaccataa cttgataatg taatttctg taatttctg taatttctg 10160
 10161 tggcccgctg gtcgcgctca cggaaactcg gggcaactca ttgtgataa taatttctg taatttctg taatttctg 10240
 10241 gcttaattcg acgaataatt ggaattttat ttattttg aatttttt taatttctg taatttctg taatttctg 10320
 10321 aaaaaaaa aaaaaaaa aaaaaaaa aaaaaaaa aaaaaaaa aaaaaaaa aaaaaaaa aaaaaaaa 10400
 10401 gttagggttt tacttgcttt aaaaaactc ccaactctc ccaactctc ccaactctc ccaactctc ccaactctc 10480
 10481 taactgttt attgcagctt ataattggtt aaaaataagc aatagcatca caaatttcac aaaaataagc aaaaataagc 10560
 10561 tgcattctag ttgtgtgtttg tccaaactca tcaatgtatc ttatcatgct tggatctagt ggcattat gaatcggcca 10640
 10641 acgcgcgggg agagggcgggt tgcgtattgg ggcgtcttcc gcttctctgc tcaatgacgc gctgctgctg gctgttcgac 10720

10721	tgcggcgagc	ggtatcagct	cactcaaaag	cggtaataacg	ggtatccaca	gaatcagggg	ataacgcagg	aaagaacatg	10800
10801	tgcagcaaa	gcccagaaac	ggcaggaag	cgtaaaaaag	ccgcgttgct	ggcgtttttc	cataggctcc	gccccctga	10880
10881	cgagcatcac	aaaaatcgac	gtcgaagtca	gaggtggcga	aaccggacag	gactataaaag	ataccaggcg	tttccccctg	10960
10961	gaagtcctct	atgctgcctc	cctgttccga	ccctggcgct	taccggatac	ctgtccgcct	ttctcccttc	gggaagcgtg	11040
11041	gcgctttctc	aatgctcgcg	ctgtagggtat	ctcagttcgg	tgtagttcgt	tcgctccaag	tcgggtcgtg	tgacacgaac	11120
11121	ccccgttcag	cccgcaccgt	gcgcctttat	cggttaactat	cgctcttgag	ccaacccggg	aagacacgac	ttatcgccac	11200
11201	tggcagcagc	cactggtaac	aggtattagca	gagcgaagta	tgtaggcggt	gttagagcgt	tcttgaagtg	gtggcctaac	11280
11281	tacgggtaca	ctagaaggac	agttatttgg	atctgcgctc	tgtgaagcc	agttaccttc	ggaaaaagag	ttggtagctc	11360
11361	ttgatccggc	aaacaaacca	ccgctggtag	cggtgggttt	tttggttgca	agcagcagat	tacgcgcaga	aaaaaaggat	11440
11441	ctcaagaaga	tcttttgatc	tttctactcg	ggcattctga	cgctcagttg	aacgaaaaat	cacgttaagg	gattttggtc	11520
11521	atgagattat	caaaaaggat	cttcacctag	atccttttaa	attaaaaaat	aagttttaaa	tcaatctaaa	gtatatatga	11600
11601	gtaaaattgg	cttgacagtt	accaatgctt	aatcagtgag	gcacctatct	cagcgatctg	tctatttcgt	tcattccatg	11680
11681	ttgcctgact	ccccgtcgtg	tagataacta	cgatacggga	gggcttacca	tctggcccca	gtgctgcaat	gataccgcga	11760
11761	gaccacgct	caccggctcc	agatttatca	gcaataaacc	agccagccgg	aagggccgag	cgcaagtg	gtcctgcaac	11840
11841	tttatccgcc	tccatccagt	ctattaattg	ttgcccggaa	gctagagtaa	gtagttcgc	agttaatagt	ttgcgcaacg	11920
11921	ttgttgccat	tgctacagcc	atcgtgggtg	cacgctcgtc	gittggtag	gcttcattca	gtcccggttc	ccaacgatca	12000
12001	agcgagttta	catgatcccc	catgttgtgc	aaaaagcgg	ttagctcctt	cggtcctcgg	atcgttgtca	gaagtaagtt	12080
12081	ggccgcagtg	ttatcactca	tggttatggc	agcaatcgat	aattctctta	ctgtcatgcc	atccgtaaga	gtcctttctg	12160
12161	tgacttggta	gtactcaacc	aagtcattct	gagaatagtg	tatgcggcga	ccgagtttgc	cttgcccggc	gtcaaatcgg	12240
12241	gataataccg	cgccacatag	cagaaactta	accactcgt	gcacccaact	gatcttcagc	atcttttact	ttcaccagcg	12320
12321	cttacccgctg	ttgagatcca	gttcgatgta	atgccgcaaa	aaaggggaata	agggcgacac	ggaaatgttg	aatactcata	12400
12401	ttcttgcgtg	agcaaaaaa	ttgaagcatt	tatcagggtt	attgtctcat	gagcggtac	atatttgaat	gtatttagaa	12480
12481	ctcttccitt	ttcaatatta	gttcgatgta	tcccgaaaa	gtgccacctg	acgtctaaga	aaccattatt	atcatgacat	12560
12561	aaataaaca	aatagggttc	cgcgcacatt	tcccgaaat	cgccggtttc	ggtagtgacg	gtgaaaacct	ctgacacatg	12640
12641	taacctataa	aatagggttc	atcacgaggg	taagcggatg	ccgggagcag	acaagcccg	caaggcgcgt	cagcgggtgt	12720
12721	cagctccccg	agacgggtcac	agcttctgtc	ggcatcagag	cagattgtac	tgagagtgca	ccatatcgac	gctctccctt	12800
12801	tggcgggtgt	cggggctggc	ttaactatgc	tactagggtg	agggcgttga	gcaccggccg	cgcaaggaat	ggtagcgtcg	12880
12881	atgcgactcc	tgcattaggga	agcagccccg	ccatatatgg	agttccgcgt	tacataactt	acggtaaatg	gcccgcctgg	12960
12961	taatcaatta	cgggggtcatt	agttcatagc	ccatatatgg	agttccgcgt	ccatagtaac	gccaataggg	actttccatt	13040
13041	ctgaccgccc	aacgaccccc	gcccattgac	gtcaataatg	acgtatgttc	caagtgtatc	atatgccaag	tacgccccct	13120
13121	gacgtcaatg	ggtggagtat	ttacggtaaa	ctgcccactt	ggcagtaacat	gaccttattg	gactttccta	cttgccagta	13200
13201	attgacgtca	atgacggtaa	atggcccgc	tggtcattatg	ccagttatg	acatcaatgg	gcgtggatag	cgttttgact	13280
13281	catctacgta	ttagtcatcg	tattaccat	ggtgatgcgg	ttttggcagt	tggcaccaaa	atcaacggga	ctttccaaaa	13360
13361	cacggggatt	tccaagtctc	caccccattg	acgtcaatgg	gagtttgttt	gtggcagggc	tataataagca	gagctctctg	13440
13441	tgctcgtaaca	actccgcccc	attgacgcaa	atgggcggta	ggcgtgtacg	gtgggaggtc	gtgggaggtc	gtgggaggtc	13520
13521	gctaactaga	gaaccactg	cttaactggc	ttatcgaaat	taatacgact	cactataagg	agaccgggaag	cttgaattc	13599
	10	20	30	40	50	60	70	80	

pcDNA3-FL-E7 SEQ ID NO:21 **FL-E7** fusion sequence is shown in bold, caps

1	10	20	30	40	50	60	70	80	
81	gacgagtcgg	gagatctccc	gatccccctat	ggtcgactct	cagtacaatc	tgctctgatg	cgcacatagt	aagccagtat	80
161	ctgctccctg	cttgtgtgtt	ggaggtcgct	gagtagtgcg	cgagcaaat	ttaagctaca	acaaggcaag	gcttgaccga	160
241	caattgcagt	aagaatctgc	ttagggttag	gcgttttgcg	ctgcttcgcg	atgtacgggc	cagatatacg	gcttgacatt	240
321	gattattgac	tagttattaa	tagtaataca	ttacgggggtc	attagttcat	agcccatata	tggagttcgc	cggtacataa	320
401	cttacgggtaa	atggcccgcc	tggtcgaccg	cccaagacc	cccgcccat	gacgtcaata	atgacgtatg	ttcccatagt	400
481	aacgcaata	gggactttcc	attgacgtca	atgggtggac	tatttacggg	aaactgccc	cttggcagta	catcaagtgt	480
561	atcatatgcc	aagtacgccc	cttattgacg	tcaatgacgg	taaatggccc	gocctggcatt	atgcccagta	catgacctta	560
641	tgggactttc	ctactttgga	gtacatctac	gtttccaagt	ctccaccca	ttgacgtcaa	cggttttggc	agtacatcaa	640
721	tggggtgga	tagcgttttg	actcacgggg	atttccaaagt	ctccaccca	ttgacgtcaa	tgggagtttg	ttttggcacc	720
801	aaaatacaacg	ggactttcca	aaatgtcgta	acaactccgc	cccattgacg	caaatgggcg	gtaggcgtgt	acggtgggag	800
881	gtctatataa	gcagagctct	ctggctaact	agagaaccca	ctgcttactg	gcttatcgaa	attaatacga	ctcactatag	880
961	ggagacccaa	gctggctagc	gtttaaacgg	gccctctaga	ATGACAGTGC	TGGCGCCAGC	CTGGAGCCCA	AATTCCTCCC	960
1041	TGTTGCTGCT	GTTGCTGCTG	CTGAGTCTCT	GCTGCGGGG	GACACCTGAC	TGTTACTTCA	GCCACAGTCC	CATCTCCTCC	1040
1121	AACTTCAAAG	TGAAGTTTAG	AGAGTTGACT	GACCACCTGC	TAAAGATTA	CCAGTCACT	GTGGCGGTCA	ATCTTCAGGA	1120
1201	CGAGAAGCAC	TGCAAGGCTT	TGTGGAGCCT	CTTCCTAGCC	CAGCGCTGGA	TAGAGCAACT	GAAGACTGTG	GCAGGGTCTA	1200
1281	AGATGCAAAAC	GCTTCTGGAG	GACGTCAACA	CCGAGATACA	TTTTGTCAAC	TCATGTACCT	TCCAGCCCCCT	ACCAGAAATGT	1280
1361	CTGCGATTTC	TCCAGACCAA	CATCTCCAC	CTCTGAAGG	ACACCTGCAC	ACAGTGTCTT	GCTGTGAAGC	CCTGTATCGG	1360
1441	GAAGGCTTGC	CAGAAATTCT	CTCGGTGCCT	GAGGTGTCAG	TGCCAGCCGG	ACTCTCCAC	CTGCTGCCCC	CCAAGGAGTC	1440
1521	CCATAGCCCT	AGAAGCCACG	GAGTCTCCAG	AGCTCGGCC	CAGGCGAGGA	TCCATGCATG	GACATACACC	TACATTGCAT	1520
1601	GAATATATGT	TAGATTGCA	ACCAGAGACA	ACTGATCTCT	ACTGTTATGA	GCAATTAAAT	GACAGCTCAG	AGGAGGAGGA	1600
1681	TGAATAATAGT	GGTCCAGCTG	GACAAGCAGA	ACCGACAGA	GCCCATTACA	ATATTGTAAC	CTTTTGTGTC	AAGTGTGACT	1680
1761	CTACGCTTCG	GTTGTGCGTA	CAAAGCACAC	ACGTAGACAT	TCGTACTTTG	GAAGACCTGT	TAATGGGCAC	ACTAGGAATT	1760
1841	GTGTGCCCCA	TCTGTCTCA	GGATAAGCTT	aagtttaaac	cgtgatcag	cctcgactgt	gocctctagt	tgcccagccat	1840
1921	ctgttgtttg	ccccctcccc	gtgccttccct	tgacctctgga	aggtgcccact	cccactgtcc	tttcctaata	aaatgaggaa	1920
2001	attgcatcgc	attgtctgag	taggtgtcat	tctattcttg	gggttggggt	ggggcaggac	agcaaagggg	aggattggga	2000
2081	agacaatagc	aggcatgctg	gggatgcggt	gggctctatg	gcttctgag	cggaagaac	cagctggggc	tctagggggg	2080
2161	atccccacgc	gcoctgtagc	ggcgcattaa	gcgcggcggg	tgtggtgggt	acgcgcagcg	tgaccgctac	acttgcccagc	2160
2241	gcccctagcg	cggtcccttt	cgctttcttc	ccttcccttc	tcgcccagtt	cgccggcctt	ccccgtcaag	ctctaaatcg	2240
2321	gggcatccct	ttagggttcc	gatttagtgc	tttacggcac	ctcgacccca	aaaaacttga	ttagggtgat	ggttcacgta	2320
2401	gtggggccatc	gcocctgatag	acgggttttc	gccctttgac	gttgagagtc	acgttcttta	atagtggaat	cttgttccaa	2400
2481	actgggaacaa	cactcaaccc	tatctcggtc	tattcttttg	atttataagg	gattttgggg	atttcggcct	attggttaaa	2480
2561	aaatgagctg	atttaacaaa	aatttaacgc	gaattaatc	tgttggaatgt	gtgtcagta	gggtgtggaa	agtcocagg	2560
2641	ctccccagga	aggcagaagt	atgcaaaaga	tgcattctcaa	ttagtacgca	accagggtgtg	gaaagtcccc	aggtccccca	2640
2721	gcaggcagaa	gtatgcaaa	catgcatctc	aattagtcag	caaccatagt	cccgccctta	actccgcca	tcccggccct	2720
2801	aactccgccc	agttccgccc	attctccgcc	ccatggctga	gcttttttgg	tttattatgc	agaggccgag	gcgcctctgt	2800
2881	cctctgagct	attccagaag	tagttaggag	gcttttttgg	cttcctagtc	ttttgcaaaa	agctccggg	agcttgtata	2880
2961	tccattttcg	gatctgatca	agagacagga	ttaggatcgt	ttcgcatagc	tgaacaagat	ggattgcacg	cagggttctcc	2960
3041	ggccgcttgg	gtggagaggc	tattcggtca	tgactgggca	caacagacaa	tcggctgctc	tgatgcgcgc	gtgttccggc	3040
3121	tgtcagcgca	ggggcgcccc	gttctttttg	tcaagaccca	cctgtccggt	gcccctgaatg	aactgcagga	cgaggcagcg	3120

3121	cggtatcgt	ggctggccac	gacggggcgtt	ccttgccgag	ctgtgctcga	cgttgtcact	gaagcgggaa	gggactggct	3200
3201	gctattgggc	gaagtgcgg	ggcaggatct	cctgtcatct	caccttgctc	ctgcgcagaa	agatccatc	atggctgatg	3280
3281	caatgcggcg	gctgcatacg	cttgatccgg	ctacctgcc	attcgaccac	caagcgaac	atcgcatcga	gcgagcacgt	3360
3361	actcggatgg	aagccggtct	tgtgatcag	gatgatctgg	acgaagagca	tcagggggctc	gcgcagccg	aactgttcgc	3440
3441	caggctcaag	gcgcgcatgc	ccgacggcga	ggatctcgtc	gtgacctatg	gcgatgcctg	cttgccgaat	atcatggtgg	3520
3521	aaaatggcgg	cttttctgga	ttcatcgact	gtggccggct	gggtgtggcg	gacgcgtatc	aggacatagc	gttggtctacc	3600
3601	cgtgatattg	ctgaagagct	tggcggcgaa	tgggtgacc	gcttctcgt	gctttacggt	atgcgcgtc	ccgattcgca	3680
3681	gcgcategoc	ttctatcgcc	ttcttgacga	gttcttctga	gcgggactct	ggggttcgaa	atgcgcgacc	aagcgacgcc	3760
3761	caacctgcca	tcacgagatt	togatccac	cgccgccttc	tatgaagggt	tgggcttcgg	aatcgtttct	cgggacgccg	3840
3841	gctggatgat	cctccagcgc	ggggatctca	tgtcggaagt	cttcgcccac	cccaacttgt	ttattgcagc	ttataatggt	3920
3921	tacaaataaa	gcaatagcat	cacaaatttc	acaaataaag	catttttttc	actgcattct	agttgtgggt	tgtccaaact	4000
4001	catcaatgta	tcttatcatg	tctgtatacc	gtcgacctct	agctagagct	tggcgtaaat	atggttcatag	ctgtttcctg	4080
4081	tgtgaattg	ttatccgctc	acaattccac	acaacatacg	agccggaagc	ataaagtgt	aagcctgggg	tgcctaatga	4160
4161	gtgagctaac	tcacattaat	tgcgttgccg	tcactgcgc	ctttccagtc	gggaaacctg	tcgtgccagc	tgcattaatg	4240
4241	aatcgccaa	gcgcgggga	gagggcgttt	gcgtattggg	cgctcttcgc	cttctctgct	cactgactcg	ctgcgctcgg	4320
4321	tcgttcggct	gcgcgagcg	gtatcagctc	actcaaaagg	ggtaatacgg	ttatccacag	aatcagggga	taacgcagga	4400
4401	aagaacatgt	gagcaaaaag	ccagcaaaa	cccaagacc	gtaaaaaggc	cgcttgctg	gcgtttttcc	ataggctcgc	4480
4481	ccccctgac	gagcatcaca	aaaatcgacg	ctcaagtcag	aggtggcgaa	accgacaggg	actataaaga	taccaggcgt	4560
4561	ttccctctgg	aagctccctc	gtgcgctctc	ctgttccgac	cctgcccgtt	accggatacc	tgtccgcctt	tctccctcgc	4640
4641	ggaagcgtgg	cgctttctca	atgctcacgc	tgtaggtatc	tcagttcggg	gtaggtcgtt	gcgtccaagc	tgggctgtgt	4720
4721	gcacgaacc	ccggttcagc	ccgaccgctg	cgcttatcc	ggtaactatc	gtcttgagtc	caaccocggt	agacacgact	4800
4801	tatcgccact	ggcagcagcc	actggtaaca	ggattagcag	agcgaggtat	gtaggcgggtg	ctacagagtt	cttgaaagtgg	4880
4881	tgggctaact	acggctacac	tagaaggaca	gtatttggt	tctgcgtct	gctgaagcca	gttaccttcg	gaaaaagagt	4960
4961	tggtagctct	tgatccggca	aacaaaccac	cgctggtagc	gggtggtttt	ttggttgcaa	gcagcagatt	acgcgcagaa	5040
5041	aaaaaggatc	tcaagaagat	ccttggatct	tttctacggg	gtctgacgct	cagtggaagt	aaaactcacg	taaagggtat	5120
5121	ttggtcatga	gattatcaaa	aaggatcttc	acctagatcc	ttttaaat	aaaatgaagt	tttaaatcaa	tctaaagtat	5200
5201	atatgagtaa	acttggtctg	acagttacca	atgcttaatc	agtgaggcac	ctatctcagc	gatctgtcta	tttcggtcat	5280
5281	ccatagttgc	ctgactcccc	gtcgtgtaga	taactacgat	acgggagggc	ttaccatctg	gccccagtgc	tgcactgata	5360
5361	ccgcgagacc	cacgctcaac	ggctccagat	ttatcagcaa	taaaccagcc	agcggaaagg	gccgagcgca	gaagtgggtcc	5440
5441	tgcaacttta	tccgcctcca	tccagttctat	taattgttgc	cgggaagcta	gagtaagtag	ttcgccagtt	aatagtttgc	5520
5521	gcaacgttgt	tgccattgct	acaggcatcg	tgggtgtcacg	ctcgtcgttt	ggtaggctt	cattcagctc	cggttcccaa	5600
5601	cgatcaaggc	gagttacatg	atcccccatg	ttgtgcaaaa	aagcgggttag	ctccttcggt	cctccgatcg	ttgtcagaag	5680
5681	taagttggcc	gcagtggtat	cactcatggt	tatggcagca	ctgcataatt	ctcttactgt	catgccatcc	gtaagatgct	5760
5761	tttctgtgac	tgggtgagtac	tcaaccaagt	cattctgaga	atagtgtatg	cggcgaccga	gttgcctctg	ccggcgctca	5840
5841	atacgggata	ataccgcgcc	acatagcaga	actttaaaag	tgtctcatcat	tggaaaaact	tcttcggggc	gaaaaactctc	5920
5921	aaggatctta	ccgctgttga	gatccagttc	gatgtaacc	actcgtgcac	ccaactgatc	ttcagcatct	tttactttca	6000
6001	ccagcgtttc	tgggtgagca	aaaacaggaa	ggcaaaaatg	cgcaaaaatg	ggaataaggg	cgacacggaa	atggttgaata	6080
6081	ctcatactct	tcccttttca	atattattga	agcatttatc	aggggtattg	tctcatgagc	ggatacatat	ttgaatgtat	6160
6161	ttagaaaaat	aaacaaatag	gggttcgcgc	cacatttccc	cgaaaagtgc	cacctgacgt	c	6221	

There is no page 40

GENERAL RECOMBINANT DNA METHODS

Basic texts disclosing general methods of molecular biology, all of which are incorporated by reference, include: Sambrook, J *et al.*, *Molecular Cloning: A Laboratory Manual*, 2nd Edition, Cold Spring Harbor Press, Cold Spring Harbor, NY, 1989; Ausubel, FM *et al.* *Current Protocols in Molecular Biology*, Vol. 2, Wiley-Interscience, New York, (current edition); Kriegler, *Gene Transfer and Expression: A Laboratory Manual* (1990); Glover, DM, ed, *DNA Cloning: A Practical Approach*, vol. I & II, IRL Press, 1985; Albers, B. *et al.*, *Molecular Biology of the Cell*, 2nd Ed., Garland Publishing, Inc., New York, NY (1989); Watson, JD *et al.*, *Recombinant DNA*, 2nd Ed., Scientific American Books, New York, 1992; and Old, RW *et al.*, *Principles of Gene Manipulation: An Introduction to Genetic Engineering*, 2nd Ed., University of California Press, Berkeley, CA (1981).

Unless otherwise indicated, a particular nucleic acid sequence is intended to encompass conservative substitution variants thereof (*e.g.*, degenerate codon substitutions) and a complementary sequence. The term “nucleic acid” is synonymous with “polynucleotide” and is intended to include a gene, a cDNA molecule, an mRNA molecule, as well as a fragment of any of these such as an oligonucleotide, and further, equivalents thereof (explained more fully below). Sizes of nucleic acids are stated either as kilobases (kb) or base pairs (bp). These are estimates derived from agarose or polyacrylamide gel electrophoresis (PAGE), from nucleic acid sequences which are determined by the user or published. Protein size is stated as molecular mass in kilodaltons (kDa) or as length (number of amino acid residues). Protein size is estimated from PAGE, from sequencing, from presumptive amino acid sequences based on the coding nucleic acid sequence or from published amino acid sequences.

Specifically, cDNA molecules encoding the amino acid sequence corresponding to the fusion polypeptide of the present invention or fragments or derivatives thereof can be synthesized by the polymerase chain reaction (PCR) (see, for example, U.S. 4,683,202) using primers derived the sequence of the protein disclosed herein. These cDNA sequences can then be assembled into a eukaryotic or prokaryotic expression vector and the resulting vector can be used to direct the synthesis of the fusion polypeptide or its fragment or derivative by appropriate host cells, for example COS or CHO cells.

This invention includes isolated nucleic acids having a nucleotide sequence encoding the novel fusion polypeptides that comprise a MHC-I-PP or a DC-PP and an antigen, fragments thereof or equivalents thereof. The term nucleic acid as used herein is intended to include such fragments or equivalents. The nucleic acid sequences of this invention can be DNA or RNA.

A cDNA nucleotide sequence the fusion polypeptide can be obtained by isolating total mRNA from an appropriate cell line. Double stranded cDNA is prepared from total mRNA. cDNA can be inserted into a suitable plasmid, bacteriophage or viral vector using any one of a number of known techniques.

In reference to a nucleotide sequence, the term “equivalent” is intended to include sequences encoding structurally homologous and/or a functionally equivalent proteins. For example, a natural polymorphism Hsp70 or FL nucleotide sequence (especially at the third base of a codon) may be manifest as “silent” mutations which do not change the amino acid sequence. Furthermore, there may be one or more naturally occurring isoforms or related, immunologically cross-reactive family members of these proteins. Such isoforms or family members are defined as proteins that share function amino acid sequence similarity to, for example, Hsp70 or FL.

Fragment of Nucleic Acid

A fragment of the nucleic acid sequence is defined as a nucleotide sequence having fewer nucleotides than the nucleotide sequence encoding the full length MHC-I-PP or DC-PP protein, antigenic polypeptide or the fusion thereof.. This invention includes such nucleic acid fragments that encode polypeptides which retain (1) the ability of the fusion polypeptide to induce increases in frequency or reactivity of T cells, preferably CD8+ T cells, that are specific for the antigen part of the fusion polypeptide.

For example, a nucleic acid fragment as intended herein encodes a FL polypeptide that retains the ability to improve the immunogenicity of an antigen when administered as a fusion polypeptide with an antigenic polypeptide or peptide.

Generally, the nucleic acid sequence encoding a fragment of a FL or Hsp70 polypeptide comprises of nucleotides from the sequence encoding the mature protein (or an active fragment thereof such as the FL ECD or the C-terminal domain of Hsp70).

Nucleic acid sequences of this invention may also include linker sequences, natural or modified restriction endonuclease sites and other sequences that are useful for manipulations related to cloning, expression or purification of encoded protein or fragments. These and other modifications of nucleic acid sequences are described herein or are well-known in the art.

The techniques for assembling and expressing DNA coding sequences for MHC-I-PP or DC-pp types of proteins, and DNA coding sequences for antigenic polypeptides, include synthesis of oligonucleotides, PCR, transforming cells, constructing vectors, expression systems, and the like; these are well-established in the art such that those of ordinary skill are familiar with standard resource materials, specific conditions and procedures.

EXPRESSION VECTORS AND HOST CELLS

This invention includes an expression vector comprising a nucleic acid sequence encoding a MHC-I-PP/antigen fusion polypeptide or a DC-PP/antigen fusion polypeptide operably linked to at least one regulatory sequence.

The term “expression vector” or “expression cassette” as used herein refers to a nucleotide sequence which is capable of affecting expression of a protein coding sequence in a host compatible with such sequences. Expression cassettes include at least a promoter operably linked with the polypeptide coding sequence; and, optionally, with other sequences, *e.g.*, transcription termination signals. Additional factors necessary or helpful in effecting expression may also be included, *e.g.*, enhancers.

“Operably linked” means that the coding sequence is linked to a regulatory sequence in a manner that allows expression of the coding sequence. Known regulatory sequences are selected to direct expression of the desired protein in an appropriate host cell. Accordingly, the term “regulatory sequence” includes promoters, enhancers and other expression control elements. Such regulatory sequences are described in, for example, Goeddel, *Gene Expression Technology. Methods in Enzymology*, vol. 185, Academic Press, San Diego, Calif. (1990)).

Thus, expression cassettes include plasmids, recombinant viruses, any form of a recombinant “naked DNA” vector, and the like. A “vector” comprises a nucleic acid which can infect, transfect, transiently or permanently transduce a cell. It will be recognized that a vector can be a naked nucleic acid, or a nucleic acid complexed with protein or lipid. The vector optionally comprises viral or bacterial nucleic acids and/or proteins, and/or membranes (*e.g.*, a cell membrane, a viral lipid envelope, *etc.*). Vectors include, but are not limited to replicons (*e.g.*, RNA replicons (see Example 1, below), bacteriophages) to which fragments of DNA may be attached and become replicated. Vectors thus include, but are not limited to RNA, autonomous self-replicating circular or linear DNA or RNA, *e.g.*, plasmids, viruses, and the like (U.S. Patent No. 5,217,879), and includes both the expression and nonexpression plasmids. Where a recombinant microorganism or cell culture is described as hosting an “expression vector” this includes both extrachromosomal circular and linear DNA and DNA that has been incorporated into the host chromosome(s). Where a vector is being maintained by a host cell, the vector may either be stably replicated by the cells during mitosis as an autonomous structure, or is incorporated within the host’s genome.

Those skilled in the art appreciate that the particular design of an expression vector of this invention depends on considerations such as the host cell to be transfected and/or the type of protein to be expressed.

The present expression vectors comprise the full range of nucleic acid molecules encoding the various embodiments of the fusion polypeptide and its functional derivatives (defined herein) including polypeptide fragments, variants, *etc.*

Such expression vectors are used to transfect host cells (*in vitro*, *ex vivo* or *in vivo*) for expression of the DNA and production of the encoded proteins which include fusion proteins or peptides. It will be understood that a genetically modified cell expressing the fusion polypeptide may transiently express the exogenous DNA for a time sufficient for the cell to be useful for its stated purpose.

The present invention provides methods for producing the fusion polypeptides, fragments and derivatives. For example, a host cell transfected with a nucleic acid vector that encodes the fusion polypeptide is cultured under appropriate conditions to allow expression of the polypeptide.

Host cells may also be transfected with one or more expression vectors that singly or in combination comprise DNA encoding at least a portion of the fusion polypeptide and DNA encoding at least a portion of a second protein, so that the host cells produce yet further fusion polypeptides that include both the portions.

A culture typically includes host cells, appropriate growth media and other byproducts. Suitable culture media are well known in the art. The fusion polypeptide can be isolated from medium or cell lysates using conventional techniques for purifying proteins and peptides, including ammonium sulfate precipitation, fractionation column chromatography (e.g. ion exchange, gel filtration, affinity chromatography, etc.) and/or electrophoresis (see generally, "Enzyme Purification and Related Techniques", *Methods in Enzymology*, 22:233-577 (1971)). Once purified, partially or to homogeneity, the recombinant polypeptides of the invention can be utilized in pharmaceutical compositions as described in more detail herein.

The term "isolated" as used herein, when referring to a molecule or composition, such as a MHC-I-PP or nucleic acid coding therefor, means that the molecule or composition is separated from at least one other compound (protein, other nucleic acid, *etc.*) or from other contaminants with which it is natively associated or becomes associated during processing. An isolated composition can also be substantially pure. An isolated composition can be in a homogeneous state and can be dry or in aqueous solution. Purity and homogeneity can be determined, for example, using analytical chemical techniques such as polyacrylamide gel electrophoresis (PAGE) or high performance liquid chromatography (HPLC). Even where a protein has been isolated so as to appear as a homogenous or dominant band in a gel pattern, there are trace contaminants which co-purify with it.

Prokaryotic or eukaryotic host cells transformed or transfected to express the fusion polypeptide or a homologue or functional derivative thereof are within the scope of the invention. For example, the fusion polypeptide may be expressed in bacterial cells such as *E. coli*, insect cells (baculovirus), yeast, or mammalian cells such as Chinese hamster ovary cells (CHO) or human cells. Other suitable host cells may be found in Goeddel, (1990) *supra* or are otherwise known to those skilled in the art.

Expression in eukaryotic cells leads to partial or complete glycosylation and/or formation of relevant inter- or intra-chain disulfide bonds of the recombinant protein.

Although preferred vectors are described in the Examples, other examples of expression vectors are provided here. Examples of vectors for expression in yeast *S. cerevisiae* include pYepSec1 (Baldari
5 *et al.*, (1987) *EMBO J.* 6:229-234), pMFa (Kurjan *et al.* (1982) *Cell* 30:933-943), pJRY88 (Schultz *et al.*, (1987) *Gene* 54:113-123), and pYES2 (Invitrogen Corporation, San Diego, Calif.). Baculovirus vectors available for expression of proteins in cultured insect cells (SF 9 cells) include the pAc series (Smith *et al.*, (1983) *Mol. Cell Biol.* 3:2156-2165,) and the pVL series (Lucklow, V. A., and Summers, M. D., (1989) *Virology* 170:31-39). Generally, COS cells (Gluzman, Y., (1981) *Cell* 23:175-182) are
10 used in conjunction with such vectors as pCDM 8 (Aruffo A. and Seed, B., *supra*, for transient amplification/expression in mammalian cells, while CHO (*dhfr*-negative CHO) cells are used with vectors such as pMT2PC (Kaufman *et al.* (1987), *EMBO J.* 6:187-195) for stable amplification/expression in mammalian cells. The NS0 myeloma cell line (a glutamine synthetase expression system.) is available from Celltech Ltd.

15 Often, in fusion expression vectors, a proteolytic cleavage site is introduced at the junction of the reporter group and the target protein to enable separation of the target protein from the reporter group subsequent to purification of the fusion protein. Proteolytic enzymes for such cleavage and their recognition sequences include Factor Xa, thrombin and enterokinase.

Typical fusion expression vectors include pGEX (Amrad Corp., Melbourne, Australia), pMAL
20 (New England Biolabs, Beverly, Mass.) and pRIT5 (Pharmacia, Piscataway, NJ) which fuse glutathione S-transferase, maltose E binding protein, or protein A, respectively, to the target recombinant protein.

Inducible non-fusion expression vectors include pTrc (Amann *et al.*, (1988) *Gene* 69:301-315) and pET 11d (Studier *et al.*, *Gene Expression Technology: Methods in Enzymology* 185, Academic Press, San Diego, Calif. (1990) 60-89). While target gene expression relies on host RNA polymerase
25 transcription from the hybrid trp-lac fusion promoter in pTrc, expression of target genes inserted into pET 11d relies on transcription from the T7 gn10-lacO fusion promoter mediated by coexpressed viral RNA polymerase (T7gn1). This viral polymerase is supplied by host strains BL21(DE3) or HMS174(DE3) from a resident λ prophage harboring a T7gn1 under the transcriptional control of the lacUV 5 promoter.

30 One embodiment of this invention is a transfected cell which expresses novel fusion polypeptide.

Vector Construction

Construction of suitable vectors containing the desired coding and control sequences employs standard ligation and restriction techniques which are well understood in the art. Isolated plasmids, DNA sequences, or synthesized oligonucleotides are cleaved, tailored, and re-ligated in the form desired.

5 The DNA sequences which form the vectors are available from a number of sources. Backbone vectors and control systems are generally found on available "host" vectors which are used for the bulk of the sequences in construction. For the pertinent coding sequence, initial construction may be, and usually is, a matter of retrieving the appropriate sequences from cDNA or genomic DNA libraries. However, once the sequence is disclosed it is possible to synthesize the entire gene sequence *in vitro* starting from the individual nucleotide derivatives. The entire gene sequence for genes of sizeable length, e.g., 500-1000 bp may be prepared by synthesizing individual overlapping complementary oligonucleotides and filling in single stranded nonoverlapping portions using DNA polymerase in the presence of the deoxyribonucleotide triphosphates. This approach has been used successfully in the construction of several genes of known sequence. See, for example, Edge, M. D., *Nature* (1981) 292:756; Nambair, K. P., *et al.*, *Science* (1984) 223:1299; and Jay, E., *J Biol Chem* (1984) 259:6311.

15 Synthetic oligonucleotides are prepared by either the phosphotriester method as described by references cited above or the phosphoramidite method as described by Beaucage, S. L., and Caruthers, M. H., *Tet Lett* (1981) 22:1859; and Matteucci, M. D., and Caruthers, M. H., *J Am Chem Soc* (1981) 103:3185 and can be prepared using commercially available automated oligonucleotide synthesizers. Kinase treatment of single strands prior to annealing or for labeling is achieved using an excess, e.g., about 10 units of polynucleotide kinase to 1 nmole substrate in the presence of 50 mM Tris, pH 7.6, 10 mM MgCl₂, 5 mM dithiothreitol, 1-2 mM ATP, 1.7 pmoles γ -³²P-ATP (2.9 mCi/mmmole), 0.1 mM spermidine, 0.1 mM EDTA.

20 Once the components of the desired vectors are thus available, they can be excised and ligated using standard restriction and ligation procedures. Site-specific DNA cleavage is performed by treating with the suitable restriction enzyme (or enzymes) under conditions which are generally understood in the art, and the particulars of which are specified by the manufacturer of these commercially available restriction enzymes. See, e.g., New England Biolabs, Product Catalog. In general, about 1 mg of plasmid or DNA sequence is cleaved by one unit of enzyme in about 20 ml of buffer solution; in the examples
30 herein, typically, an excess of restriction enzyme is used to insure complete digestion of the DNA substrate. Incubation times of about one hour to two hours at about 37°C. are workable, although variations can be tolerated. After each incubation, protein is removed by extraction with phenol/chloroform, and may be followed by ether extraction, and the nucleic acid recovered from aqueous fractions by precipitation with ethanol. If desired, size separation of the cleaved fragments may

be performed by polyacrylamide gel or agarose gel electrophoresis using standard techniques. A general description of size separations is found in *Methods in Enzymology* (1980) 65:499-560.

Restriction cleaved fragments may be blunt ended by treating with the large fragment of *E. coli* DNA polymerase I (Klenow) in the presence of the four deoxynucleotide triphosphates (dNTPs) using incubation times of about 15 to 25 min at 20° to 25° C. in 50 mM Tris pH 7.6, 50 mM NaCl, 6 mM MgCl₂, 6 mM DTT and 0.1-1.0 mM dNTPs. The Klenow fragment fills in at 5' single-stranded overhangs but chews back protruding 3' single strands, even though the four dNTPs are present. If desired, selective repair can be performed by supplying only one of the, or selected, dNTPs within the limitations dictated by the nature of the overhang. After treatment with Klenow, the mixture is extracted with phenol/chloroform and ethanol precipitated. Treatment under appropriate conditions with S1 nuclease or BAL-31 results in hydrolysis of any single-stranded portion.

Ligations are typically performed in 15-50 ml volumes under the following standard conditions and temperatures: for example, 20 mM Tris-HCl pH 7.5, 10mM MgCl₂, 10 mM DTT, 33 µg/ml BSA, 10-50mM NaCl, and either 40 µM ATP, 0.01-0.02 (Weiss) units T4 DNA ligase at 0° C. (for "sticky end" ligation) or 1mM ATP, 0.3-0.6 (Weiss) units T4 DNA ligase at 14° C. (for "blunt end" ligation). Intermolecular "sticky end" ligations are usually performed at 33-100 µg/ml total DNA concentrations (5-100 nM total end concentration). Intermolecular blunt end ligations are performed at 1 mM total ends concentration.

In vector construction employing "vector fragments", the fragment is commonly treated with bacterial alkaline phosphatase (BAP) or calf intestinal alkaline phosphatase (CIAP) in order to remove the 5' phosphate and prevent self-ligation. Digestions are conducted at pH 8 in approximately 10 mM Tris-HCl, 1 mM EDTA using BAP or CIAP at about 1 unit/mg vector at 60° for about one hour. The preparation is extracted with phenol/chloroform and ethanol precipitated. Alternatively, re-ligation can be prevented in vectors which have been double digested by additional restriction enzyme and separation of the unwanted fragments.

Any of a number of methods are used to introduce mutations into the coding sequence to generate the variants of the invention. These mutations include simple deletions or insertions, systematic deletions, insertions or substitutions of clusters of bases or substitutions of single bases.

For example, modifications of the MHC-I-PP or DC-pp or the antigenic polypeptide DNA sequence are created by site-directed mutagenesis, a well-known technique for which protocols and reagents are commercially available (Zoller, MJ *et al.*, *Nucleic Acids Res* (1982) 10:6487-6500 and Adelman, JP *et al.*, *DNA* (1983) 2:183-193)). Correct ligations for plasmid construction are confirmed, for example, by first transforming *E. coli* strain MC1061 (Casadaban, M., *et al.*, *J Mol Biol* (1980) 138:179-207) or other suitable host with the ligation mixture. Using conventional methods,

transformants are selected based on the presence of the ampicillin-, tetracycline- or other antibiotic resistance gene (or other selectable marker) depending on the mode of plasmid construction. Plasmids are then prepared from the transformants with optional chloramphenicol amplification optionally following chloramphenicol amplification ((Clewell, DB *et al.*, *Proc Natl Acad Sci USA* (1969) 62:1159; Clewell, D. B., *J Bacteriol* (1972) 110:667). Several mini DNA preps are commonly used. See, e.g., Holmes, DS, *et al.*, *Anal Biochem* (1981) 114:193-197; Birnboim, HC *et al.*, *Nucleic Acids Res* (1979) 7:1513-1523. The isolated DNA is analyzed by restriction and/or sequenced by the dideoxy nucleotide method of Sanger (*Proc Natl Acad Sci USA* (1977) 74:5463) as further described by Messing, *et al.*, *Nucleic Acids Res* (1981) 9:309, or by the method of Maxam *et al.* *Methods in Enzymology* (1980) 65:499.

Vector DNA can be introduced into mammalian cells via conventional techniques such as calcium phosphate or calcium chloride co-precipitation, DEAE-dextran-mediated transfection, lipofection, or electroporation. Suitable methods for transforming host cells can be found in Sambrook *et al. supra* and other standard texts.

Often, in fusion expression vectors, a proteolytic cleavage site is introduced at the junction of the reporter group and the target protein to enable separation of the target protein from the reporter group subsequent to purification of the fusion protein. Proteolytic enzymes for such cleavage and their recognition sequences include Factor Xa, thrombin and enterokinase.

Known fusion expression vectors include pGEX (Amrad Corp., Melbourne, Australia), pMAL (New England Biolabs, Beverly, Mass.) and pRIT5 (Pharmacia, Piscataway, NJ) which fuse glutathione S-transferase, maltose E binding protein, or protein A, respectively, to the target recombinant protein.

Inducible non-fusion expression vectors include pTrc (Amann *et al.*, (1988) *Gene* 69:301-315) and pET 11d (Studier *et al.*, *Gene Expression Technology: Methods in Enzymology* 185, Academic Press, San Diego, Calif. (1990) 60-89). While target gene expression relies on host RNA polymerase transcription from the hybrid trp-lac fusion promoter in pTrc, expression of target genes inserted into pET 11d relies on transcription from the T7 gn10-lacO fusion promoter mediated by coexpressed viral RNA polymerase (T7gn1). This viral polymerase is supplied by host strains BL21(DE3) or HMS174(DE3) from a resident λ prophage harboring a T7gn1 under the transcriptional control of the lacUV 5 promoter.

Promoters and Enhancers

A promoter region of a DNA or RNA molecule binds RNA polymerase and promotes the transcription of an "operably linked" nucleic acid sequence. As used herein, a "promoter sequence" is the nucleotide sequence of the promoter which is found on that strand of the DNA or RNA which is

transcribed by the RNA polymerase. Two sequences of a nucleic acid molecule, such as a promoter and a coding sequence, are “operably linked” when they are linked to each other in a manner which permits both sequences to be transcribed onto the same RNA transcript or permits an RNA transcript begun in one sequence to be extended into the second sequence. Thus, two sequences, such as a promoter
5 sequence and a coding sequence of DNA or RNA are operably linked if transcription commencing in the promoter sequence will produce an RNA transcript of the operably linked coding sequence. In order to be “operably linked” it is not necessary that two sequences be immediately adjacent to one another in the linear sequence.

The preferred promoter sequences of the present invention must be operable in mammalian cells and may be either eukaryotic or viral promoters. Although preferred promoters are described in the
10 Examples, other useful promoters and regulatory elements are discussed below. Suitable promoters may be inducible, repressible or constitutive. A “constitutive” promoter is one which is active under most conditions encountered in the cell’s environmental and throughout development. An “inducible” promoter is one which is under environmental or developmental regulation. A “tissue specific” promoter
15 is active in certain tissue types of an organism. An example of a constitutive promoter is the viral promoter MSV-LTR, which is efficient and active in a variety of cell types, and, in contrast to most other promoters, has the same enhancing activity in arrested and growing cells. Other preferred viral promoters include that present in the CMV-LTR (from cytomegalovirus) (Bashart, M. *et al.*, *Cell* 41:521 (1985)) or in the RSV-LTR (from Rous sarcoma virus) (Gorman, C.M., *Proc. Natl. Acad. Sci. USA*
20 79:6777 (1982). Also useful are the promoter of the mouse metallothionein I gene (Hamer, D., *et al.*, *J. Mol. Appl. Gen.* 1:273-288 (1982)); the TK promoter of Herpes virus (McKnight, S., *Cell* 31:355-365 (1982)); the SV40 early promoter (Benoist, C., *et al.*, *Nature* 290:304-310 (1981)); and the yeast *gal4* gene promoter (Johnston, S.A., *et al.*, *Proc. Natl. Acad. Sci. (USA)* 79:6971-6975 (1982); Silver, P.A., *et al.*, *Proc. Natl. Acad. Sci. (USA)* 81:5951-5955 (1984)). Other illustrative descriptions of transcriptional
25 factor association with promoter regions and the separate activation and DNA binding of transcription factors include: Keegan *et al.*, *Nature* (1986) 231:699; Fields *et al.*, *Nature* (1989) 340:245; Jones, *Cell* (1990) 61:9; Lewin, *Cell* (1990) 61:1161; Ptashne *et al.*, *Nature* (1990) 346:329; Adams *et al.*, *Cell* (1993) 72:306. The relevant disclosure of all of these above-listed references is hereby incorporated by reference.

The promoter region may further include an octamer region which may also function as a tissue
30 specific enhancer, by interacting with certain proteins found in the specific tissue. The enhancer domain of the DNA construct of the present invention is one which is specific for the target cells to be transfected, or is highly activated by cellular factors of such target cells. Examples of vectors (plasmid or retrovirus) are disclosed in (Roy-Burman *et al.*, U.S. Patent No. 5,112,767). For a general discussion

of enhancers and their actions in transcription, see, Lewin, B.M., *Genes IV*, Oxford University Press, Oxford, (1990), pp. 552-576. Particularly useful are retroviral enhancers (*e.g.*, viral LTR). The enhancer is preferably placed upstream from the promoter with which it interacts to stimulate gene expression. For use with retroviral vectors, the endogenous viral LTR may be rendered enhancer-less and substituted with other desired enhancer sequences which confer tissue specificity or other desirable properties such as transcriptional efficiency.

The nucleic acid sequences of the invention can also be chemically synthesized using standard techniques. Various methods of chemically synthesizing polydeoxynucleotides are known, including solid-phase synthesis which, like peptide synthesis, has been fully automated with commercially available DNA synthesizers (See, *e.g.*, Itakura *et al.* U.S. Pat. No. 4,598,049; Caruthers *et al.* U.S. Pat. No. 4,458,066; and Itakura U.S. Pat. Nos. 4,401,796 and 4,373,071, incorporated by reference herein).

PROTEINS AND POLYPEPTIDES

The terms “polypeptide,” “protein,” and “peptide” when referring to compositions of the invention are meant to include variants, analogues, and mimetics with structures and/or activity that substantially correspond to the polypeptide or peptide from which the variant, *etc.* was derived.

The present invention includes an “isolated” fusion polypeptide comprising a MHC-I-PP and/or a DC-PP linked to an antigenic polypeptide. A preferred MHC-I-PP is Hsp70. A preferred DC-PP is FL or the ECD or FL. Preferred fusion polypeptides are Hsp70/E7 (SEQ ID NO:8) and FL-E7 (SEQ ID NO:12). While the present disclosure exemplifies the full length Hsp70 and the ECD of FL, it is to be understood that homologues of Hsp70 from other bacteria or from eukaryotic origin, homologues of FL or its ECD, and mutants thereof that possess the characteristics disclosed herein are intended within the scope of this invention.

The term “chimeric” or “fusion” polypeptide or protein refers to a composition comprising at least one polypeptide or peptide sequence or domain that is chemically bound in a linear fashion with a second polypeptide or peptide domain. One embodiment of this invention is an isolated or recombinant nucleic acid molecule encoding a fusion protein comprising at least two domains, wherein the first domain comprises an MHC-I-PP or a DC-PP *e.g.*, Hsp60 or FL, and the second domain comprising an antigenic epitope, *e.g.*, an MHC class I-binding peptide epitope. Additional domains can comprise a polypeptide, peptide, polysaccharide, or the like. The “fusion” can be an association generated by a peptide bond, a chemical linking, a charge interaction (*e.g.*, electrostatic attractions, such as salt bridges, H-bonding, *etc.*) or the like. If the polypeptides are recombinant, the “fusion protein” can be translated from a common mRNA. Alternatively, the compositions of the domains can be linked by any chemical or electrostatic means. The chimeric molecules of the invention (*e.g.*, MHC-I-PP- fusion proteins) can also include additional sequences, *e.g.*, linkers, epitope tags, enzyme cleavage recognition sequences,

signal sequences, secretion signals, and the like. Alternatively, a peptide can be linked to a carrier simply to facilitate manipulation or identification/ location of the peptide.

Also included is a “functional derivative” of Hsp70 or FL, which refers to an amino acid substitution variant, a “fragment,” or a “chemical derivative” of the protein, which terms are defined below. A functional derivative retains measurable (a) Hsp70-like or (b) FL-like activity, preferably that of promoting immunogenicity of one or more antigenic epitopes fused thereto by either (a) promoting presentation by class I pathways or (b) promoting maturation or activation of APCs, which permits the “functional derivative’s” utility in accordance with the present invention. “Functional derivatives” encompass “variants” and “fragments” regardless of whether the terms are used in the conjunctive or the alternative herein.

A functional homologue must possess the above biochemical and biological activity. In view of this functional characterization, use of homologous Hsp70 or FL proteins including proteins not yet discovered, fall within the scope of the invention if these proteins have sequence similarity and the recited biochemical and biological activity.

To determine the percent identity of two amino acid sequences or of two nucleic acid sequences, the sequences are aligned for optimal comparison purposes (*e.g.*, gaps can be introduced in one or both of a first and a second amino acid or nucleic acid sequence for optimal alignment and non-homologous sequences can be disregarded for comparison purposes). In a preferred method of alignment, Cys residues are aligned.

In a preferred embodiment, the length of a sequence being compared is at least 30%, preferably at least 40%, more preferably at least 50%, even more preferably at least 60%, and even more preferably at least 70%, 80%, or 90% of the length of the reference sequence (*e.g.*, Hsp70, SEQ ID NO:4 and FL-ECD, SEQ ID NO:10). The amino acid residues (or nucleotides) at corresponding amino acid (or nucleotide) positions are then compared. When a position in the first sequence is occupied by the same amino acid residue (or nucleotide) as the corresponding position in the second sequence, then the molecules are identical at that position (as used herein amino acid or nucleic acid “identity” is equivalent to amino acid or nucleic acid “homology”). The percent identity between the two sequences is a function of the number of identical positions shared by the sequences, taking into account the number of gaps, and the length of each gap, which need to be introduced for optimal alignment of the two sequences.

The comparison of sequences and determination of percent identity between two sequences can be accomplished using a mathematical algorithm. In a preferred embodiment, the percent identity between two amino acid sequences is determined using the Needleman and Wunsch (*J. Mol. Biol.*

48:444-453 (1970) algorithm which has been incorporated into the GAP program in the GCG software package (available at <http://www.gcg.com>), using either a Blossom 62 matrix or a PAM250 matrix, and a gap weight of 16, 14, 12, 10, 8, 6, or 4 and a length weight of 1, 2, 3, 4, 5, or 6. In yet another preferred embodiment, the percent identity between two nucleotide sequences is determined using the GAP
5 program in the GCG software package (available at <http://www.gcg.com>), using a NWSgapdna.CMP matrix and a gap weight of 40, 50, 60, 70, or 80 and a length weight of 1, 2, 3, 4, 5, or 6. In another embodiment, the percent identity between two amino acid or nucleotide sequences is determined using the algorithm of E. Meyers and W. Miller (CABIOS, 4:11-17 (1989)) which has been incorporated into the ALIGN program (version 2.0), using a PAM120 weight residue table, a gap length penalty of 12 and
10 a gap penalty of 4.

The nucleic acid and protein sequences of the present invention can further be used as a "query sequence" to perform a search against public databases, for example, to identify other family members or related sequences. Such searches can be performed using the NBLAST and XBLAST programs (version 2.0) of Altschul *et al.* (1990) *J. Mol. Biol.* 215:403-10. BLAST nucleotide searches can be performed
15 with the NBLAST program, score = 100, wordlength = 12 to obtain nucleotide sequences homologous to Hsp70 or FL nucleic acid molecules. BLAST protein searches can be performed with the XBLAST program, score = 50, wordlength = 3 to obtain amino acid sequences homologous to HVP22 protein molecules. To obtain gapped alignments for comparison purposes, Gapped BLAST can be utilized as described in Altschul *et al.* (1997) *Nucleic Acids Res.* 25:3389-3402. When utilizing BLAST and
20 Gapped BLAST programs, the default parameters of the respective programs (*e.g.*, XBLAST and NBLAST) can be used. See <http://www.ncbi.nlm.nih.gov>.

Thus, a homologue of Hsp70 or of FL described above is characterized as having (a) functional activity of native Hsp70 or FL and (b) sequence similarity to a native Hsp70 protein (such as SEQ ID NO:4) or native FL (SEQ ID NO:10) when determined as above, of at least about 20% (at the amino acid
25 level), preferably at least about 40%, more preferably at least about 70%, even more preferably at least about 90%.

It is within the skill in the art to obtain and express such a protein using DNA probes based on the disclosed sequences of Hsp70 or FL. Then, the fusion protein's biochemical and biological activity can be tested readily using art-recognized methods such as those described herein, for example, a T cell
30 proliferation, cytokine secretion or a cytolytic assay, or an *in vivo* assay of tumor protection or tumor therapy. A biological assay of the stimulation of antigen-specific T cell reactivity will indicate whether the homologue has the requisite activity to qualify as a "functional" homologue.

A "variant" refers to a molecule substantially identical to either the full protein or to a fragment thereof in which one or more amino acid residues have been replaced (substitution variant) or which has

one or several residues deleted (deletion variant) or added (addition variant). A "fragment" of Hsp70 or FL refers to any subset of the molecule, that is, a shorter polypeptide of the full-length protein.

A number of processes can be used to generate fragments, mutants and variants of the isolated DNA sequence. Small subregions or fragments of the nucleic acid encoding the spreading protein, for example 1-30 bases in length, can be prepared by standard, chemical synthesis. Antisense oligonucleotides and primers for use in the generation of larger synthetic fragment.

A preferred group of variants are those in which at least one amino acid residue and preferably, only one, has been substituted by different residue. For a detailed description of protein chemistry and structure, see Schulz, GE *et al.*, *Principles of Protein Structure*, Springer-Verlag, New York, 1978, and Creighton, T.E., *Proteins: Structure and Molecular Properties*, W.H. Freeman & Co., San Francisco, 1983, which are hereby incorporated by reference. The types of substitutions that may be made in the protein molecule may be based on analysis of the frequencies of amino acid changes between a homologous protein of different species, such as those presented in Table 1-2 of Schulz *et al.* (*supra*) and Figure 3-9 of Creighton (*supra*). Based on such an analysis, conservative substitutions are defined herein as exchanges within one of the following five groups:

1	Small aliphatic, nonpolar or slightly polar residues	Ala, Ser, Thr (Pro, Gly);
2	Polar, negatively charged residues and their amides	Asp, Asn, Glu, Gln;
3	Polar, positively charged residues	His, Arg, Lys;
4	Large aliphatic, nonpolar residues	Met, Leu, Ile, Val (Cys)
5	Large aromatic residues	Phe, Tyr, Trp.

The three amino acid residues in parentheses above have special roles in protein architecture. Gly is the only residue lacking a side chain and thus imparts flexibility to the chain. Pro, because of its unusual geometry, tightly constrains the chain. Cys can participate in disulfide bond formation, which is important in protein folding.

More substantial changes in biochemical, functional (or immunological) properties are made by selecting substitutions that are less conservative, such as between, rather than within, the above five groups. Such changes will differ more significantly in their effect on maintaining (a) the structure of the peptide backbone in the area of the substitution, for example, as a sheet or helical conformation, (b) the charge or hydrophobicity of the molecule at the target site, or (c) the bulk of the side chain. Examples of such substitutions are (i) substitution of Gly and/or Pro by another amino acid or deletion or insertion of Gly or Pro; (ii) substitution of a hydrophilic residue, *e.g.*, Ser or Thr, for (or by) a hydrophobic residue, *e.g.*, Leu, Ile, Phe, Val or Ala; (iii) substitution of a Cys residue for (or by) any other residue; (iv) substitution of a residue having an electropositive side chain, *e.g.*, Lys, Arg or His, for (or by) a

residue having an electronegative charge, *e.g.*, Glu or Asp; or (v) substitution of a residue having a bulky side chain, *e.g.*, Phe, for (or by) a residue not having such a side chain, *e.g.*, Gly.

Most acceptable deletions, insertions and substitutions according to the present invention are those that do not produce radical changes in the characteristics of the wild-type or native protein in terms of its intercellular spreading activity and its ability to stimulate antigen specific T cell reactivity to an antigenic epitope or epitopes that are fused to the spreading protein. However, when it is difficult to predict the exact effect of the substitution, deletion or insertion in advance of doing so, one skilled in the art will appreciate that the effect can be evaluated by routine screening assays such as those described here, without requiring undue experimentation.

Whereas shorter chain variants can be made by chemical synthesis, for the present invention, the preferred longer chain variants are typically made by site-specific mutagenesis of the nucleic acid encoding the polypeptide, expression of the variant nucleic acid in cell culture, and, optionally, purification of the polypeptide from the cell culture, for example, by immunoaffinity chromatography using specific antibody immobilized to a column (to absorb the variant by binding to at least one epitope).

The term “chemically linked” refers to any chemical bonding of two moieties, *e.g.*, as in one embodiment of the invention, where an **MHC-I-PP** or DC-PP is chemically linked to an antigenic peptide. Such chemical linking includes the peptide bonds of a recombinantly or *in vivo* generated fusion protein.

Chemical Derivatives

“Chemical derivatives” of the polypeptide or fusion polypeptide of the invention contain additional chemical moieties not normally a part of the protein. Covalent modifications of the polypeptide are included within the scope of this invention. Such derivatized moieties may improve the solubility, absorption, biological half life, and the like. Moieties capable of mediating such effects are disclosed, for example, in *Remington's Pharmaceutical Sciences*, 16th ed., Mack Publishing Co., Easton, PA (1980).

Such modifications may be introduced into the molecule by reacting targeted amino acid residues of the polypeptide with an organic derivatizing agent that is capable of reacting with selected side chains or terminal residues. Another modification is cyclization of the protein. Examples of chemical derivatives of the polypeptide follow.

Lysiny and amino terminal residues are derivatized with succinic or other carboxylic acid anhydrides. Derivatization with a cyclic carboxylic anhydride has the effect of reversing the charge of the lysiny residues. Other suitable reagents for derivatizing amino-containing residues include imidoesters such as methyl picolinimidate; pyridoxal phosphate; pyridoxal; chloroborohydride;

trinitrobenzenesulfonic acid; O-methylisourea; 2,4 pentanedione; and transaminase-catalyzed reaction with glyoxylate.

Carboxyl side groups, aspartyl or glutamyl, may be selectively modified by reaction with carbodiimides (R-N=C=N-R') such as 1-cyclohexyl-3-(2-morpholinyl-(4-ethyl) carbodiimide or 1-ethyl-3-(4-azonia-4,4-dimethylpentyl) carbodiimide. Furthermore, aspartyl and glutamyl residues can be converted to asparaginy and glutaminy residues by reaction with ammonia.

Other modifications include hydroxylation of proline and lysine, phosphorylation of hydroxyl groups of seryl or threonyl residues, methylation of the amino group of lysine (Creighton, *supra*, pp. 79-86), acetylation of the N-terminal amine, and amidation of the C-terminal carboxyl groups.

Also included are peptides wherein one or more D-amino acids are substituted for one or more L-amino acids.

Multimeric Peptides

The present invention also includes longer polypeptides in which a basic peptidic sequence obtained from the sequence of either the MHC-I-PP or the DC-PP, or the antigenic polypeptide or peptide unit, is repeated from about two to about 100 times, with or without intervening spacers or linkers. It is understood that such multimers may be built from any of the peptide variants defined herein. Moreover, a peptide multimer may comprise different combinations of peptide monomers and the disclosed substitution variants thereof. Such oligomeric or multimeric peptides can be made by chemical synthesis or by recombinant DNA techniques as discussed herein. When produced chemically, the oligomers preferably have from 2-12 repeats of the basic peptide sequence. When produced recombinantly, the multimers may have as many repeats as the expression system permits, for example from two to about 100 repeats.

In tandem multimers, preferably dimers and trimers, of the fusion polypeptide, the chains bonded by interchain disulfide bonds or other "artificial" covalent bonds between the chains such that the chains are "side-by-side" rather than "end to end."

THERAPEUTIC COMPOSITIONS AND THEIR ADMINISTRATION

A vaccine composition comprising the nucleic acid encoding the fusion polypeptide, or a cell expressing this nucleic acid is administered to a mammalian subject, preferably a human. The vaccine composition is administered in a pharmaceutically acceptable carrier in a biologically effective or a therapeutically effective amount. The composition may be given alone or in combination with another protein or peptide such as an immunostimulatory molecule. Treatment may include administration of an adjuvant, used in its broadest sense to include any nonspecific immune stimulating compound such as

an interferon. Adjuvants contemplated herein include resorcinols, non-ionic surfactants such as polyoxyethylene oleyl ether and n-hexadecyl polyethylene ether.

A therapeutically effective amount is a dosage that, when given for an effective period of time, achieves the desired immunological or clinical effect.

5 A therapeutically active amount of a nucleic acid encoding the fusion polypeptide may vary according to factors such as the disease state, age, sex, and weight of the individual, and the ability of the peptide to elicit a desired response in the individual. Dosage regimes may be adjusted to provide the optimum therapeutic response. For example, several divided doses may be administered daily or the dose may be proportionally reduced as indicated by the exigencies of the therapeutic situation. A
10 therapeutically effective amounts of the protein, in cell associated form may be stated in terms of the protein or cell equivalents.

Thus an effective amount is between about 1 nanogram and about 10 milligram per kilogram of body weight of the recipient, more preferably between about 0.1 μg and 1 $\mu\text{g}/\text{kg}$. Dosage forms suitable for internal administration preferably contain (for the latter dose range) from about 0.01 μg to 100 μg of
15 active ingredient per unit. The active ingredient may vary from 0.5 to 95% by weight based on the total weight of the composition. Alternatively, an effective dose of cells expressing the nucleic acid is between about 10^4 and 10^8 cells. Those skilled in the art of immunotherapy will be able to adjust these doses without undue experimentation.

The active compound may be administered in a convenient manner, *e.g.*, injection by a
20 convenient and effective route. Preferred routes include subcutaneous, intradermal, intravenous and intramuscular routes. Other possible routes include oral administration, intrathecal, inhalation, transdermal application, or rectal administration. For the treatment of existing tumors which have not been completely resected or which have recurred, direct intratumoral injection is also intended.

Depending on the route of administration, the active compound may be coated in a material to
25 protect the compound from the action of enzymes, acids and other natural conditions which may inactivate the compound. Thus it may be necessary to coat the composition with, or co-administer the composition with, a material to prevent its inactivation. For example, an enzyme inhibitors of nucleases or proteases (*e.g.*, pancreatic trypsin inhibitor, diisopropylfluorophosphate and trasylol).or in an appropriate carrier such as liposomes (including water-in-oil-in-water emulsions as well as conventional
30 liposomes (Strejan *et al.*, (1984) *J. Neuroimmunol* 7:27).

As used herein "pharmaceutically acceptable carrier" includes any and all solvents, dispersion media, coatings, antibacterial and antifungal agents, isotonic and absorption delaying agents, and the like. The use of such media and agents for pharmaceutically active substances is well known in the art. Except insofar as any conventional media or agent is incompatible with the active compound, use thereof

in the therapeutic compositions is contemplated. Supplementary active compounds can also be incorporated into the compositions.

Preferred pharmaceutically acceptable diluents include saline and aqueous buffer solutions. Pharmaceutical compositions suitable for injection include sterile aqueous solutions (where water
5 soluble) or dispersions and sterile powders for the extemporaneous preparation of sterile injectable solutions or dispersion. Isotonic agents, for example, sugars, polyalcohols such as mannitol, sorbitol, sodium chloride may be included in the pharmaceutical composition. In all cases, the composition should be sterile and should be fluid. It should be stable under the conditions of manufacture and storage and must include preservatives that prevent contamination with microorganisms such as bacteria
10 and fungi. Dispersions can also be prepared in glycerol, liquid polyethylene glycols, and mixtures thereof and in oils. Under ordinary conditions of storage and use, these preparations may contain a preservative to prevent the growth of microorganisms.

The carrier can be a solvent or dispersion medium containing, for example, water, ethanol, polyol (for example, glycerol, propylene glycol, and liquid polyethylene glycol, and the like), and
15 suitable mixtures thereof. The proper fluidity can be maintained, for example, by the use of a coating such as lecithin, by the maintenance of the required particle size in the case of dispersion and by the use of surfactants.

Prevention of the action of microorganisms can be achieved by various antibacterial and antifungal agents, for example, parabens, chlorobutanol, phenol, ascorbic acid, thimerosal, and the like.

20 Compositions are preferably formulated in dosage unit form for ease of administration and uniformity of dosage. Dosage unit form refers to physically discrete units suited as unitary dosages for a mammalian subject; each unit contains a predetermined quantity of active material (*e.g.*, the nucleic acid vaccine) calculated to produce the desired therapeutic effect, in association with the required pharmaceutical carrier. The specification for the dosage unit forms of the invention are dictated by and
25 directly dependent on (a) the unique characteristics of the active material and the particular therapeutic effect to be achieved, and (b) the limitations inherent in the art of compounding such an active compound for the treatment of, and sensitivity of, individual subjects

For lung instillation, aerosolized solutions are used. In a sprayable aerosol preparations, the active protein may be in combination with a solid or liquid inert carrier material. This may also be
30 packaged in a squeeze bottle or in admixture with a pressurized volatile, normally gaseous propellant. The aerosol preparations can contain solvents, buffers, surfactants, and antioxidants in addition to the protein of the invention.

Other pharmaceutically acceptable carriers for the nucleic acid vaccine compositions according to the present invention are liposomes, pharmaceutical compositions in which the active protein is

contained either dispersed or variously present in corpuscles consisting of aqueous concentric layers adherent to lipidic layers. The active protein is preferably present in the aqueous layer and in the lipidic layer, inside or outside, or, in any event, in the non-homogeneous system generally known as a liposomic suspension. The hydrophobic layer, or lipidic layer, generally, but not exclusively, comprises phospholipids such as lecithin and sphingomyelin, steroids such as cholesterol, more or less ionic surface active substances such as dicetylphosphate, stearylamine or phosphatidic acid, and/or other materials of a hydrophobic nature. Those skilled in the art will appreciate other suitable embodiments of the present liposomal formulations.

ANTIGENS ASSOCIATED WITH PATHOGENS

A major use for the present invention is the use of the present nucleic acid compositions in therapeutic vaccine for cancer and for major chronic viral infections that cause morbidity and mortality worldwide. Such vaccines are designed to eliminate infected cells - this requires T cell responses as antibodies are often ineffective. The vaccines of the present invention are designed to meet these needs.

Preferred antigens are epitopes of pathogenic microorganisms against which the host is defended by effector T cells responses, including cytotoxic T lymphocyte (CTL) and delayed type hypersensitivity. These typically include viruses, intracellular parasites such as malaria, and bacteria that grow intracellularly such as Mycobacteria and Listeria species. Thus, the types of antigens included in the vaccine compositions of this invention are any of those associated with such pathogens (in addition, of course, to tumor-specific antigens). It is noteworthy that some viral antigens are also tumor antigens in the case where the virus is a causative factor in cancer.

In fact, the two most common cancers worldwide, hepatoma and cervical cancer, are associated with viral infection. Hepatitis B virus (HBV) (Beasley, R.P. *et al.*, *Lancet* **2**, 1129-1133 (1981) has been implicated as etiologic agent of hepatomas. 80-90% of cervical cancers express the E6 and E7 antigens (exemplified herein) from one of four "high risk" human papillomavirus types: HPV-16, HPV-18, HPV-31 and HPV-45 (Gissmann, L. *et al.*, *Ciba Found Symp.* **120**, 190-207 (1986); Beaudenon, S., *et al.* *Nature* **321**, 246-249 (1986). The HPV E6 and E7 antigens are the most promising targets for virus associated cancers in immunocompetent individuals because of their ubiquitous expression in cervical cancer. In addition to their importance as targets for therapeutic cancer vaccines, virus associated tumor antigens are also ideal candidates for prophylactic vaccines. Indeed, introduction of prophylactic HBV vaccines in Asia have decreased the incidence of hepatoma (Chang, M.H., *et al.* *New Engl. J. Med.* **336**, 1855-1859 (1997), representing a great impact on cancer prevention.

Among the most important viruses in chronic human viral infections are HPV, HBV, hepatitis C Virus (HCV), human immunodeficiency virus (HIV-1 and HIV-2), herpesviruses such as Epstein Barr

Virus (EBV), cytomegalovirus (CMV) and HSV-1 and HSV-2 and influenza virus. Useful antigens include HBV surface antigen or HBV core antigen; ppUL83 or pp89 of CMV; antigens of gp120, gp41 or p24 proteins of HIV-1; ICP27, gD2, gB of HSV; or influenza nucleoprotein (Anthony, LS *et al.*, *Vaccine* 1999; 17:373-83). Other antigens associated with pathogens that can be utilized as described herein are antigens of various parasites, includes malaria, preferably malaria peptide (NANP)40.

In addition to its applicability to human cancer and infectious diseases,, the present invention is also intended for use in treating animal diseases in the veterinary medicine context. Thus, the approaches described herein may be readily applied by one skilled in the art to treatment of veterinary herpesvirus infections including equine herpesviruses, bovine viruses such as bovine viral diarrhea virus (for example, the E2 antigen), bovine herpesviruses, Marek's disease virus in chickens and other fowl; animal retroviral and lentiviral diseases (*e.g.*, feline leukemia, feline immunodeficiency, simian immunodeficiency viruses, *etc.*); pseudorabies and rabies; and the like.

As for tumor antigens, any tumor-associated or tumor-specific antigen that can be recognized by T cells, preferably by CTL, can be used. In addition to the HPV-E7 antigen exemplified herein is mutant p53 or HER2/neu or a peptide thereof. Any of a number of melanoma-associated antigens may be used, such as MAGE-1, MAGE-3, MART-1/Melan-A, tyrosinase, gp75, gp100, BAGE, GAGE-1, GAGE-2, GnT-V, and p15 (see, US 6,187,306).

The following references set forth principles and current information in the field of basic, medical and veterinary virology and are incorporated by reference: *Fields Virology*, Fields, BN *et al.*, eds., Lippincott Williams & Wilkins, NY, 1996; *Principles of Virology: Molecular Biology, Pathogenesis, and Control*, Flint, S.J. *et al.*, eds., Amer Society for Microbiology, Washington, 1999; *Principles and Practice of Clinical Virology*, 4th Edition, Zuckerman. A.J. *et al.*, eds, John Wiley & Sons, NY, 1999; *The Hepatitis C Viruses*, by Hagedorn, CH *et al.*, eds., Springer Verlag, 1999; *Hepatitis B Virus: Molecular Mechanisms in Disease and Novel Strategies for Therapy*, Koshy, R. *et al.*, eds., World Scientific Pub Co, 1998; *Veterinary Virology*, Murphy, F.A. *et al.*, eds., Academic Press, NY, 1999; *Avian Viruses: Function and Control*, Ritchie, B.W., Iowa State University Press, Ames, 2000; *Virus Taxonomy: Classification and Nomenclature of Viruses: Seventh Report of the International Committee on Taxonomy of Viruses*, by M. H. V. Van Regenmortel, MHV *et al.*, eds., Academic Press; NY, 2000.

DELIVERY OF VACCINE NUCLEIC ACID TO CELLS AND ANIMALS

Examples I-III describe certain preferred approaches to delivery of the vaccines of the present invention: naked DNA, self-replicating RNA and virally-based suicide DNA. A broader description of other approaches including viral and nonviral vectors and delivery mechanisms follow.

DNA delivery involves introduction of a “foreign” DNA into a cell *ex vivo* and ultimately, into a live animal or directly into the animal. Several general strategies for gene delivery (= delivery of nucleic acid vectors) for purposes that include “gene therapy” have been studied and reviewed extensively (Yang, N-S., *Crit. Rev. Biotechnol.* 12:335-356 (1992); Anderson, W.F., *Science* 256:808-813 (1992);
5 Miller, A.S., *Nature* 357:455-460 (1992); Crystal, R.G., *Amer. J. Med.* 92(suppl 6A):44S-52S (1992); Zwiebel, J.A. *et al.*, *Ann. N.Y. Acad. Sci.* 618:394-404 (1991); McLachlin, J.R. *et al.*, *Prog. Nucl. Acid Res. Molec. Biol.* 38:91-135 (1990); Kohn, D.B. *et al.*, *Cancer Invest.* 7:179-192 (1989), which references are herein incorporated by reference in their entirety).

One approach comprises nucleic acid transfer into primary cells in culture followed by
10 autologous transplantation of the *ex vivo* transformed cells into the host, either systemically or into a particular organ or tissue.

The term “systemic administration” refers to administration of a composition or agent such as a molecular vaccine as described herein, in a manner that results in the introduction of the composition into the subject’s circulatory system or otherwise permits its spread throughout the body. “Regional”
15 administration refers to administration into a specific, and somewhat more limited, anatomical space, such as intraperitoneal, intrathecal, subdural, or to a specific organ. The term “local administration” refers to administration of a composition or drug into a limited, or circumscribed, anatomic space, such as intratumoral injection into a tumor mass, subcutaneous injections, intramuscular injections. One of skill in the art would understand that local administration or regional administration may also result in
20 entry of a composition into the circulatory system.

For accomplishing the objectives of the present invention, nucleic acid therapy would be accomplished by direct transfer of a the functionally active DNA into mammalian somatic tissue or organ *in vivo*. DNA transfer can be achieved using a number of approaches described below. These systems can be tested for successful expression *in vitro* by use of a selectable marker (*e.g.*, G418
25 resistance) to select transfected clones expressing the DNA, followed by detection of the presence of the antigen-containing expression product (after treatment with the inducer in the case of an inducible system) using an antibody to the product in an appropriate immunoassay. Efficiency of the procedure, including DNA uptake, plasmid integration and stability of integrated plasmids, can be improved by linearizing the plasmid DNA using known methods, and co-transfection using high molecular weight
30 mammalian DNA as a “carrier”.

Examples of successful “gene transfer” reported in the art include: (a) direct injection of plasmid DNA into mouse muscle tissues, which led to expression of marker genes for an indefinite period of time (Wolff, J.A. *et al.*, *Science* 247:1465 (1990); Acsadi, G. *et al.*, *The New Biologist* 3:71 (1991)); (b) retroviral vectors are effective for *in vivo* and *in situ* infection of blood vessel tissues; (c) portal vein

injection and direct injection of retrovirus preparations into liver effected gene transfer and expression *in vivo* (Horzaglou, M. *et al.*, *J. Biol. Chem.* 265:17285 (1990); Koleko, M. *et al.*, *Human Gene Therapy* 2:27 (1991); Ferry, N. *et al.*, *Proc. Natl. Acad. Sci. USA* 88:8387 (1991)); (d) intratracheal infusion of recombinant adenovirus into lung tissues was effective for *in vivo* transfer and prolonged expression of foreign genes in lung respiratory epithelium (Rosenfeld, M.A. *et al.*, *Science* 252:431 (1991)); (e) Herpes simplex virus vectors achieved *in vivo* gene transfer into brain tissue (Ahmad, F. *et al.*, eds, *Miami Short Reports - Advances in Gene Technology: The Molecular Biology of Human Genetic Disease*, Vol 1, Boehringer Mannheim1 Biochemicals, USA, 1991).

Retroviral-mediated human therapy utilizes amphotrophic, replication-deficient retrovirus systems (Temin, H.M., *Human Gene Therapy* 1:111 (1990); Temin *et al.*, U.S. Patent 4,980,289; Temin *et al.*, U.S. Patent 4,650,764; Temin *et al.*, U.S. Patent No. 5,124,263; Wills, J.W. U.S. Patent 5,175,099; Miller, A.D., U.S. Patent No. 4,861,719). Such vectors have been used to introduce functional DNA into human cells or tissues, for example, the adenosine deaminase gene into lymphocytes, the NPT-II gene and the gene for tumor necrosis factor into tumor infiltrating lymphocytes. Retrovirus-mediated gene delivery generally requires target cell proliferation for gene transfer (Miller, D.G. *et al.*, *Mol. Cell. Biol.* 10:4239 (1990). This condition is met by certain of the preferred target cells into which the present DNA molecules are to be introduced, *i.e.*, actively growing tumor cells. Gene therapy of cystic fibrosis using transfection by plasmids using any of a number of methods and by retroviral vectors has been described by Collins *et al.*, U.S. Patent 5,240,846.

The DNA molecules encoding the fusion polypeptides of the present invention may be packaged into retrovirus vectors using packaging cell lines that produce replication-defective retroviruses, as is well-known in the art (see, for example, Cone, R.D. *et al.*, *Proc. Natl. Acad. Sci. USA* 81:6349-6353 (1984); Mann, R.F. *et al.*, *Cell* 33:153-159 (1983); Miller, A.D. *et al.*, *Molec. Cell. Biol.* 5:431-437 (1985);, Sorge, J., *et al.*, *Molec. Cell. Biol.* 4:1730-1737 (1984); Hock, R.A. *et al.*, *Nature* 320:257 (1986); Miller, A.D. *et al.*, *Molec. Cell. Biol.* 6:2895-2902 (1986). Newer packaging cell lines which are efficient and safe for gene transfer have also been described (Bank *et al.*, U.S. 5,278,056.

This approach can be utilized in a site specific manner to deliver the retroviral vector to the tissue or organ of choice. Thus, for example, a catheter delivery system can be used (Nabel, EG *et al.*, *Science* 244:1342 (1989)). Such methods, using either a retroviral vector or a liposome vector, are particularly useful to deliver the nucleic acid to be expressed to a blood vessel wall, or into the blood circulation of a tumor.

Other virus vectors may also be used, including recombinant adenoviruses (Horowitz, M.S., In: *Virology*, Fields, BN *et al.*, eds, Raven Press, New York, 1990, p. 1679; Berkner, K.L., *Biotechniques* 6:616 9191988), Strauss, S.E., In: *The Adenoviruses*, Ginsberg, HS, ed., Plenum Press, New York, 1984,

chapter 11), herpes simplex virus (HSV) for neuron-specific delivery and persistence. Advantages of adenovirus vectors for human gene delivery include the fact that recombination is rare, no human malignancies are known to be associated with such viruses, the adenovirus genome is double stranded DNA which can be manipulated to accept foreign genes of up to 7.5 kb in size, and live adenovirus is a safe human vaccine organisms. Adeno-associated virus is also useful for human therapy (Samulski, R.J. *et al.*, *EMBO J.* 10:3941 (1991) according to the present invention.

Another vector which can express the DNA molecule of the present invention, and is useful in the present therapeutic setting, particularly in humans, is vaccinia virus, which can be rendered non-replicating (U.S. Patents 5,225,336; 5,204,243; 5,155,020; 4,769,330; Sutter, G *et al.*, *Proc. Natl. Acad. Sci. USA* (1992) 89:10847-10851; Fuerst, T.R. *et al.*, *Proc. Natl. Acad. Sci. USA* (1989) 86:2549-2553; Falkner F.G. *et al.*; *Nucl. Acids Res* (1987) 15:7192; Chakrabarti, S *et al.*, *Molec. Cell. Biol.* (1985) 5:3403-3409). Descriptions of recombinant vaccinia viruses and other viruses containing heterologous DNA and their uses in immunization and DNA therapy are reviewed in: Moss, B., *Curr. Opin. Genet. Dev.* (1993) 3:86-90; Moss, B. *Biotechnology* (1992) 20:345-362; Moss, B., *Curr Top Microbiol Immunol* (1992) 158:25-38; Moss, B., *Science* (1991) 252:1662-1667; Piccini, A *et al.*, *Adv. Virus Res.* (1988) 34:43-64; Moss, B. *et al.*, *Gene Amplif Anal* (1983) 3:201-213.

In addition to naked DNA or RNA, or viral vectors, engineered bacteria may be used as vectors. A number of bacterial strains including *Salmonella*, BCG and *Listeria monocytogenes*(LM) (Hoiseth & Stocker, *Nature* 291, 238-239 (1981); Poirier, TP *et al.* *J. Exp. Med.* 168, 25-32 (1988); (Sadoff, J.C., *et al.*, *Science* 240, 336-338 (1988); Stover, C.K., *et al.*, *Nature* 351, 456-460 (1991); Aldovini, A. *et al.*, *Nature* 351, 479-482 (1991); Schafer, R., *et al.*, *J. Immunol.* 149, 53-59 (1992); Ikonomidis, G. *et al.*, *J. Exp. Med.* 180, 2209-2218 (1994)). These organisms display two promising characteristics for use as vaccine vectors: (1) enteric routes of infection, providing the possibility of oral vaccine delivery; and (2) infection of monocytes/macrophages thereby targeting antigens to professional APCs.

In addition to virus-mediated gene transfer *in vivo*, physical means well-known in the art can be used for direct transfer of DNA, including administration of plasmid DNA (Wolff *et al.*, 1990, *supra*) and particle-bombardment mediated gene transfer (Yang, N.-S., *et al.*, *Proc. Natl. Acad. Sci. USA* 87:9568 (1990); Williams, R.S. *et al.*, *Proc. Natl. Acad. Sci. USA* 88:2726 (1991); Zelenin, A.V. *et al.*, *FEBS Lett.* 280:94 (1991); Zelenin, A.V. *et al.*, *FEBS Lett.* 244:65 (1989); Johnston, S.A. *et al.*, *In Vitro Cell. Dev. Biol.* 27:11 (1991)). Furthermore, electroporation, a well-known means to transfer genes into cell *in vitro*, can be used to transfer DNA molecules according to the present invention to tissues *in vivo* (Titomirov, A.V. *et al.*, *Biochim. Biophys. Acta* 1088:131 ((1991)).

"Carrier mediated gene transfer" has also been described (Wu, C.H. *et al.*, *J. Biol. Chem.* 264:16985 (1989); Wu, G.Y. *et al.*, *J. Biol. Chem.* 263:14621 (1988); Soriano, P. *et al.*, *Proc. Natl.*

Acad. Sci. USA 80:7128 (1983); Wang, C-Y. *et al.*, *Proc. Natl. Acad. Sci. USA* 84:7851 (1982); Wilson, J.M. *et al.*, *J. Biol. Chem.* 267:963 (1992)). Preferred carriers are targeted liposomes (Nicolau, C. *et al.*, *Proc. Natl. Acad. Sci. USA* 80:1068 (1983); Soriano *et al.*, *supra*) such as immunoliposomes, which can incorporate acylated mAbs into the lipid bilayer (Wang *et al.*, *supra*). Polycations such as

5 asialoglycoprotein/polylysine (Wu *et al.*, 1989, *supra*) may be used, where the conjugate includes a molecule which recognizes the target tissue (e.g., asialoorosomucoid for liver) and a DNA binding compound to bind to the DNA to be transfected. Polylysine is an example of a DNA binding molecule which binds DNA without damaging it. This conjugate is then complexed with plasmid DNA according to the present invention for transfer.

10 Plasmid DNA used for transfection or microinjection may be prepared using methods well-known in the art, for example using the Quiagen procedure (Quiagen), followed by DNA purification using known methods, such as the methods exemplified herein.

Having now generally described the invention, the same will be more readily understood through reference to the following examples which are provided by way of illustration, and are not intended to be

15 limiting of the present invention, unless specified.

EXAMPLE I

Enhancement of Sindbis Virus Self-Replicating RNA Vaccine Potency by Linkage of Nucleic Acid encoding *Mycobacterium Tuberculosis* Heat Shock Protein 70 and an Antigen

20 The present study investigated whether DNA linking full-length E7 to Hsp70 can enhance the potency of self-replicating Sindbis RNA vaccines. We showed that a Sindbis RNA vaccine linking E7 with Hsp70 significantly increased expansion and activation of E7-specific CD8⁺ T cells and NK cells, bypassing the requirement for CD4⁺ T cell-mediated help and resulting in potent anti-tumor immunity against E7-expressing tumors. Mechanistic studies confirmed that the Sindbis E7/Hsp70 RNA vaccine

25 induced apoptotic death of host cells and promoted processing of this apoptotic material by dendritic cells (DCs), leading to significantly increased expansion and activation of E7-specific CD8⁺ T cells. This enhanced CD8 response resulted in a state of potent anti-tumor immunity against an E7-expressing tumor cell line.

MATERIALS AND METHODS

Plasmid DNA Constructs and Preparation

30

The generation of pcDNA3-Hsp70, pcDNA3-E7, and pcDNA3-E7/Hsp70 has been described previously (Chen *et al.*, *supra*). The Sindbis virus RNA replicon vector, SINrep5 has also been described previously (Bredenbeek, PJ *et al.*, 1993. *J Virol* 67:6439-46). For the generation of SINrep5-

Hsp70, SINrep5-E7, and SINrep5-E7/Hsp70, DNA fragments encoding *Mtb* Hsp70, HPV-16 E7, and chimeric E7/Hsp70 were isolated by cutting pcDNA3-Hsp70, pcDNA3-E7, and pcDNA3-E7/Hsp70 respectively with Xba I and Pme I restriction enzymes, followed by gel recovery from the digested products. These isolated DNA fragments were further cloned into the corresponding Xba I and Pme I sites of the SINrep5 vector to generate SINrep5-Hsp70, SINrep5-E7, and SINrep5-E7/Hsp70 constructs. SINrep5-E7/GFP constructs were generated to evaluate the effect of linkage of E7 to an irrelevant protein. For the generation of SINrep5-E7/GFP, we first constructed pcDNA3-GFP. For the generation of pcDNA3-GFP, a DNA fragment encoding the green fluorescent protein (GFP) was first amplified with PCR using pEGFPN1 DNA (Clontech, Palo Alto, CA) and a set of primers: 5'-atcggatccatggtgagcaagggcgaggag-3' (SEQ ID NO:24) and 5'-gggaagctttactgtacagctcgccatg-3' (SEQ ID NO:25). The amplified product was digested with BamHI/ HindIII and further cloned into the BamHI/HindIII cloning sites of pcDNA3 vector. For the generation of pcDNA3-E7/GFP, a DNA fragment encoding HPV-16 E7 first amplified with PCR using pcDNA3-E7 as a template and a set of primers: 5'-ggggaattcatgcatggagatacaccta-3' (SEQ ID NO:26) and 5'-ggtggatccttgagaacagatgg-3' (SEQ ID NO:27). The amplified product was then digested with EcoRI/BamHI and further cloned into the EcoRI/BamHI cloning sites of pcDNA3-GFP. E7/GFP was cut with XbaI/PmeI from pcDNA3-E7/GFP and cloned into XbaI/PmeI sites of SIN5rep. The accuracy of these constructs was confirmed by DNA sequencing.

In Vitro RNA Preparation

The generation of RNA transcripts from SINrep5-Hsp70, SINrep5-E7, SINrep5-E7/GFP, SINrep5-E7/Hsp70 and SINrep5 was performed using the protocol described by Mandl *et al.*, (Mandl, CW *et al.*, 1998. *Nat Med* 4:1438-40). SpeI was used to linearize DNA templates for the synthesis of RNA replicons from SINrep5-Hsp70, SINrep5-E7, SINrep5-E7/Hsp70 and SINrep5. RNA vaccines were transcribed *in vitro* and capped using SP6 RNA polymerase and capping analog from the *in vitro* transcription kit (Life Technologies, Rockville, MD) according to vendor's manual. After synthesis, DNA was removed by digestion with DNase I. Synthesized RNA was quantified and analyzed using denaturing formaldehyde agarose gels (Mandl *et al.*, *supra*). The purified RNA was divided into aliquots to be used for vaccination in animals and for transfection of a baby hamster kidney (BHK21) cell line. The protein expression of the transcripts was assessed by transfection of the RNA into BHK21 cells using electroporation.

Cell Lines

BHK21 cells were obtained from the ATCC (Rockville, MD) and grown in Glasgow MEM supplemented with 5% FBS, 10% tryptose phosphate broth, 2 mM glutamine, and antibiotics. Cells

were kept at 37°C in a humidified 5% CO₂ atmosphere and were passaged every 2 days. The production and maintenance of TC-1 cells has been described previously (Lin, KY *et al.*, 1996. *Cancer Research* 56:21-26). On the day of tumor challenge, TC-1 cells were harvested by trypsinization, washed twice with 1X Hanks buffered salt solution (HBSS), and finally resuspended in 1X HBSS to the designated concentration for injection.

ELISA to detect E7 Protein Expression of SINrep5 RNA vaccines

The expression of E7 protein from SINrep5-E7 and SINrep5-E7/Hsp70 RNA was determined by indirect ELISA. The quantity of E7 protein was determined using cell lysates from SINrep5-E7 or -E7/Hsp70 transfected BHK21 cells. Briefly, 10⁷ BHK21 cells were transfected with the 4µg SINrep5, SINrep5-E7, SINrep5-Hsp70, SINrep5-E7/GFP or SINrep5-E7/Hsp70 RNA transcripts respectively via electroporation as described by Liljestrom *et al.*, (Liljestrom, PS *et al.*, 1991. *J Virol* 65:4107-13). We used SINrep5-vector containing the β-gal gene and determined the transfection efficiency. The transfected cells were fixed and stained for lacZ expression using X-Gal (Sanes, JR *et al.*, 1986. *Embo J* 5:3133-42). In general, the transfection efficiency in our electroporation was consistent and measured around 30%. The transfected BHK21 cells were collected 16-20 hrs after electroporation. A 96-microwell plate was coated BHK 21 cell lysates that were transfected with various SINrep5 RNA constructs in a final volume of 100 µl, and were incubated at 4°C overnight. The bacteria-derived HPV-16 E7 proteins were used as a positive control. The wells were then blocked with PBS containing 20% fetal bovine serum. Diluted anti-E7 Ab (Zymed, San Francisco, CA) were added to the ELISA wells, and incubated on 37°C for 2 hr. After washing with PBS containing 0.05% Tween-20, the plate was incubated with 1/2000 dilution of a peroxidase-conjugated rabbit anti-mouse IgG antibody (Zymed, San Francisco, CA) at room temperature for one hour. The plate was washed, developed with 1-Step™ Turbo TMB-ELISA (Pierce, Rockford, IL), and stopped with 1M H₂SO₄. The ELISA plate was read with a standard ELISA reader at 450 nm. The quantity of E7 protein of the cell lysates was then calculated and determined by comparing with the standardized E7 protein.

Mice

6 to 8-week-old female C57BL/6 mice from the National Cancer Institute (Frederick, MD) were purchased and kept in the oncology animal facility of the Johns Hopkins Hospital (Baltimore, MD). All animal procedures were performed according to approved protocols and in accordance with recommendations for the proper use and care of laboratory animals.

RNA Vaccination

All SINrep5 RNA vaccines were generated using *in vitro* transcription as described earlier. RNA concentration was determined by optical density measured at 260 nm. The integrity and quantity of RNA transcripts were further checked using denaturing gel electrophoresis. Mice were vaccinated intramuscularly with 10 µg of SINrep5-Hsp70, SINrep5-E7, SINrep5-E7 mixed with SINrep5-Hsp70, SINrep5-E7/GFP or SINrep5 RNA vaccines in the right hind leg while SINrep5-E7/Hsp70 was administered in 0.1, 1, and 10 µg quantities.

ELISA to Measure Anti-E7 Antibodies

Anti-HPV 16 E7 antibodies in the sera were determined by direct ELISA as described previously (Wu, TC 1995. *Proc Natl Acad Sci USA* 92:11671-5). A 96-microwell plate was coated with 100 µl bacteria-derived HPV-16 E7 proteins (5 µg/ml) and incubated at 4°C overnight. The wells were then blocked with PBS containing 20% fetal bovine serum. Sera obtained from mice on day 14 post-immunization were serially diluted in PBS, added to the ELISA wells, and incubated at 37°C for 2 hr. After washing with PBS containing 0.05% Tween-20, the plate was incubated with 1/2000 dilution of a peroxidase-conjugated rabbit anti-mouse IgG antibody (Zymed, San Francisco, CA) at room temperature for one hour. The plate was washed, developed with 1-Step™ Turbo TMB-ELISA (Pierce, Rockford, IL), and stopped with 1M H₂SO₄. The ELISA plate was read with a standard ELISA reader at 450 nm.

ELISA to Measure IFN γ

Splenocytes were harvested 2 weeks after vaccination and cultured with the E7 peptide (aa 49-57) containing the MHC class I epitope (RAHYNIVTF, SEQ ID NO:22) (Feltkamp, MC *et al.*, 1993. *Eur J Immunol* 23:2242-9) or the E7 peptide (aa 30-67) containing the class II epitope (DSSEEEDEIDGPAGQAE PDRAHYNIVTFCKCDSTLRL; SEQ ID NO:23) (Tindle, RW 1991. *Proc Natl Acad Sci USA* 88:5887-91) in a total volume of 2 ml of RPMI 1640, supplemented with 10% (vol/vol) fetal bovine serum, 50 units/ml penicillin and streptomycin, 2 mM L-glutamine, 1 mM sodium pyruvate, 2 mM nonessential amino acids in a 24-well tissue culture plate for 6 days. The supernatants were harvested and assayed for the presence of IFN γ using a commercial ELISA kit (Endogen, Woburn, MA) according to the manufacturer's protocol.

Cytotoxic T Lymphocyte (CTL) Assays

Cytolysis was determined by quantitative measurements of lactate dehydrogenase (LDH) released from cells using CytoTox96®, a non-radioactive cytotoxicity assay kit (Promega, Madison, WI) according to the manufacturer's protocol. Briefly, splenocytes were harvested and pooled 2 weeks after RNA vaccination. Five mice were used for each vaccinated group. Splenocytes were cultured with the E7 peptide (aa 49-57) in a total volume of 2 ml of RPMI 1640, supplemented with 10% (vol/vol) fetal

bovine serum, 50 units/ml penicillin/ streptomycin, 2mM L-glutamine, 1mM sodium pyruvate, 2mM nonessential amino acids in a 24-well tissue culture plate for 6 days as effector cells. TC-1 tumor cells were used as target cells. The TC-1 cells mixed with splenocytes at various effector/target (E/T) ratios. After 5 hr incubation at 37°C, 50µl of the culture medium was collected to assess the amount of LDH present. The percentage of lysis was calculated from the formula: $100 \times \{(A-B)/(C-D)\}$, where A is the reading of experimental-effector signal value, B is the effector spontaneous background signal value, C is maximum signal value from target cells, D is the target spontaneous background signal value.

In Vivo Tumor Protection Experiments

For the tumor protection study, mice (5 per group) were immunized i.m. with 10µg/mouse of SINrep5-Hsp70, SINrep5-E7, SINrep5-E7 mixed with SINrep5-Hsp70, SINrep5-E7/GFP or SINrep5 RNA, or 0.1µg/mouse, 1 µg/mouse, or 10µg/mouse of SINrep5-E7/Hsp70 RNA. 14 days after immunization, mice were injected intravenously with 1×10^4 cells/mouse TC-1 tumor cells in the tail vein. Three weeks after tumor challenge, mice were euthanized. The number of tumor nodules on the lung surface in each mouse was evaluated and counted by experimenters in a blinded fashion.

In Vivo Depletion of Cells using Monoclonal Antibodies

The procedure was described previously (Lin, KY *et al.*, 1996. *Cancer Research* 56:21-26; Wu, TC *et al.*, 1995. *J. Exp. Med* 182:1415-1421). In brief, each mouse was vaccinated with 1 µg self-replicating SINrep5-E7/Hsp70 RNA i.m. and challenged with 10^4 TC-1 tumor cells i.v. (via tail vein). Depletions were initiated one week prior to tumor challenge. IgG_{2a} antibody (PharMingen, San Diego, CA) was used as a non-specific isotype control. MAb GK1.5 (Dialynas, DP 1983. *J. Immunol.* 131:2445) was used for depletion of CD4+ cells; mAb 2.43 (Sarmiento, MA *et al.*, 1980. *J. Immunol.* 125:2665) was used for depletion of CD8+ cells; and mAb PK136 (Koo, GC *et al.*, *J Immunol.* 137:3742) was used for depletion of NK1.1+ cells. Flow cytometry analysis revealed that >95% of the cells of the appropriate lymphocyte subset were depleted while numbers of cells of other subsets were unchanged. Depletion treatment was discontinued on day 21 after tumor challenge.

Cell Surface Marker Staining and Flow Cytometric Analysis

Splenocytes from naïve or vaccinated mice were immediately stained for cell surface markers according to Ji, H *et al.*, 1999, *Human Gene Therapy* 10:2727-2740. Cells were then washed once in FACScan® buffer and stained with PE-conjugated monoclonal rat anti-mouse NK1.1 antibody or FITC-conjugated monoclonal rat anti-mouse CD3 antibody (PharMingen, San Diego, CA). NK cells are NK1.1+ and CD3-negative. Flow cytometry was used to determine the percent of splenocytes that were NK cells.

In Vitro Analysis of Cell Death

10⁷ BHK21 cells were transfected with 4 µg SINrep5, SINrep5-E7, SINrep5-Hsp70 or SINrep5-E7/Hsp70 RNA transcripts as mentioned above. The transfection efficiency was around 20-30%.

Native BHK21 cells or BHK21 cells that were electroporated without SINrep5 RNA served as controls.

5 BHK21 cells were collected and assessed every 24 hr, until hour 72. The percentages of apoptotic and necrotic BHK21 cells were determined using annexin V apoptosis detection kits (PharMingen, San Diego, CA) according to the manufacturer's protocol, followed by flow cytometry.

Generation and Culture of Dendritic Cells (DCs) from Bone Marrow

10 DCs were generated by culturing bone marrow cells in the presence of GM-CSF as described previously (Lu, Z *et al.*, 2000. *J Exp Med* 191:541-550). Briefly, bone marrow was collected from the tibias of mice. Erythrocytes were lysed, and the remaining cells were passed through a nylon mesh to remove small pieces of bone and debris. The cells were collected and 10⁶ cells/ml were placed in 24-well plates in RPMI 1640, supplemented with 5% FCS, 2mM β-mercaptoethanol, 1% nonessential amino acids, 100 U/ml penicillin and 100 µg/ml streptomycin (Life Technologies, Rockville, MD), and 100
15 U/ml GM-CSF (PharMingen, San Diego, CA). Two-thirds of the medium was replaced every 2 days, and nonadherent cells were harvested on day 7. The collected cells were characterized by flow cytometry for DC markers as previously described (Wang, TL *et al.*, 2000. *J Exp Med* 191:541-550).

CTL Assay Using DCs Pulsed with BHK21 Cells that had been Transfected with Various RNA Transcripts as Target Cells

20 CTL assays using DCs pulsed with BHK21 cells that had been transfected with various RNA transcripts as target cells were performed using a protocol similar to that described by Albert, ML *et al.*, 1998. *Nature* 392:86-9 and Albert ML, *et al.*, 1998. *J Exp Med* 188:1359-68). Briefly, 10⁷ BHK21 cells were transfected with 4µg of various self-replicating SINrep5 RNA constructs via electroporation. The cells were collected 16-20 hr later. The levels of E7 protein expression in BHK21 cells transfected
25 with SINrep5-E7, or SINrep5-E7/Hsp70 RNA transcripts were similar, as determined by ELISA. 3 x 10⁵ transfected BHK21 cells were co-incubated with 10⁵ of bone marrow-derived DCs at 37°C for 48 hr. These "prepared" DCs were then used as target cells (T) and the H-2D^b-restricted E7-specific CD8+ T cells were used as effector cells (E) (Wang, *et al.*, *supra*). CTL assays were performed with 10⁴ target cells and effector cells numbers yielding E/T ratios of 1, 3 and 9, incubated in a final volume of 200 µl.
30 After 5 hrs at 37°C, 50 µl of culture supernatant were collected to assess the amount of LDH as described above. Negative controls included DCs co-incubated with untransfected BHK21 cells, transfected BHK21 cells incubated alone, untreated DCs incubated alone, and cells of the CD8+ T cell line incubated alone.

RESULTS

Construction and Characterization of Self-replicating RNA Constructs

Generation of plasmid DNA constructs and subsequent preparation of self-replicating SINrep5 RNA constructs was performed as described above. The SINrep5 vector includes DNA encoding Sindbis virus RNA replicase and the SP6 promoter ((Bredenbeek, PJ *et al.*, 1993. *J Virol* 67:6439-46)). A schematic diagram of SINrep5, SINrep5-Hsp70, SINrep5-E7, SINrep5-E7/GFP and SINrep5-E7/Hsp70 RNA transcripts using SP6 RNA polymerase is shown in Figure 1. An ELISA to test expression of E7 protein by BHK21 cells transfected with the various self-replicating RNA constructs showed that SINrep5-E7 and SINrep5-E7/Hsp70 expressed similar amounts of E7 protein.

Vaccination with Self-replicating SINrep5-E7/Hsp70 RNA Enhances an E7-Specific Cytotoxic T cell Response

CD8⁺ T lymphocytes are important effectors of anti-tumor immunity. Generation of E7-specific CD8⁺ CTLs following vaccination was assessed. Figure 2 shows results of a study wherein splenocytes from the various self-replicating SINrep5 RNA vaccines were cultured with the E7 peptide (aa 49-57) for 6 days and were examined as effector cells against TC-1 tumor cell targets. The self-replicating SINrep5-E7/Hsp70 generated a significantly greater lytic activity in the lymphocyte population compared to cells from mice vaccinated with the other SINrep5 RNA vaccines ($p < 0.001$, one-way ANOVA). The capacity of SINrep5-E7/Hsp70 RNA to generate lytic activity was approximately 7 times that induced by self-replicating SINrep5-E7 RNA ($57.2 \pm 6.8\%$ versus $8.0 \pm 1.3\%$, E/T ratio 45, $p < 0.001$).

The concentration of IFN- γ in the supernatant of stimulated splenocytes was assessed by ELISA. Splenocytes from mice given the various self-replicating RNA vaccines were cultured *in vitro* with E7 peptide (aa 49-57) (MHC class I epitope (Feltkamp *et al.*, *supra*)) for 6 days. As a negative control, the stimulatory peptide was omitted. Culture supernatants were collected for measurement of IFN- γ concentration. As shown in Figure 3, peptide-stimulated splenocytes from mice vaccinated with self-replicating E7/Hsp70 RNA secreted the highest concentration of IFN- γ compared to cells from mice given the other RNA vaccines ($p < 0.001$, one-way ANOVA). Fusion of Hsp70 to E7 significantly enhanced IFN- γ -secreting E7-specific CD8⁺ T cell activity.

Vaccination with Self-replicating SINrep5-E7/Hsp70 RNA Did Not Enhance IFN- γ -secreting E7-Specific CD4⁺ T cells or Anti-E7 Antibodies

ELISA was used to assess the E7-specific CD4⁺ T cell responses generated by the vaccines by measuring concentration of IFN- γ in the supernatant of cultured splenocytes. Splenocytes were cultured in the presence of with E7 peptide (aa 30-67) (that includes an MHC class II epitope (Tindle *et al.*, *supra*) for 6 days. The peptide was omitted in the negative control. As shown in Figure 4, there was no

significant increase in the concentration of IFN- γ from splenocytes obtained from mice vaccinated with self-replicating E7/Hsp70 RNA compared to the other RNA vaccines. Therefore, a vaccine in which Hsp70 is fused to E7 does appear to enhance IFN- γ -secreting E7-specific CD4⁺ T cell activity.

The quantity of anti- E7 antibodies in the sera of the vaccinated mice was determined by direct ELISA two weeks after vaccination. Sera of mice vaccinated with SINrep5-E7/Hsp70 did have higher titers of E7-specific antibodies compared to mice vaccinated with other RNA vaccine constructs.

Vaccination with Self-Replicating SINrep5-E7/Hsp70 RNA Protects Mice Better Against the Growth of E7-Expressing TC-1 Tumors

An *in vivo* tumor protection experiment was performed using different doses of SINrep5-E7/Hsp70 RNA administered intramuscularly in the right hind leg. Each mouse was vaccinated with 10 μ g of one of the following constructs: self-replicating SINrep5, SINrep5-E7, SINrep5-Hsp70, SINrep5-E7 mixed with SINrep5-Hsp70, SINrep5-E7/GFP or SINrep5-E7/Hsp70 RNA. Self-replicating E7/Hsp70 RNA was also tested at doses of 0.1 and 1 μ g/mouse. E7-expressing TC-1 cells were injected i.v. 14 days later. Such tumor challenge simulates hematogenous spread of the tumor cells, allowing evaluation of vaccine effects on metastasis to the lungs via the bloodstream. Pulmonary nodules were assessed 21 days after tumor challenge. Figure 5 shows a lower mean number of pulmonary nodules in mice vaccinated with the self-replicating E7/Hsp70 RNA vaccines (0.1 μ g, 1 μ g, and 10 μ g) compared to mice given the other RNA vaccines ($p < 0.001$, one-way ANOVA). Representative photographs of the lung tumors (unmagnified) are shown in Figure 6. The results demonstrated that self-replicating RNA SINrep5-E7/Hsp70 vaccines protected mice from i.v. tumor challenge even at the lower dose of 0.1 μ g whereas vaccination with 10 μ g of SINrep5 without insert, SINrep5-E7, SINrep5-Hsp70, SINrep5-E7 mixed with SINrep5-Hsp70, or SINrep5-E7/GFP RNA showed no or little protection, developing numerous tumor nodules. These results also showed that linkage of RNA encoding E7 to RNA encoding an irrelevant protein such as GFP did not result in an antitumor effect, but rather that and that antitumor protection offered by Hsp70 required physical linkage of E7 to Hsp70 at the nucleic acid level.

CD8⁺ T Cells and NK cells Are Important for the Anti-tumor Effect

To determine the types of lymphocytes required for protection against E7-expressing tumors, *in vivo* antibody depletion experiments were done in which depletion was initiated one week before tumor challenge and terminated on day 21 after tumor challenge. As shown in Figure 7, the mean number of pulmonary nodules from mice depleted of CD8⁺ T cells or of NK1.1⁺ cells was significantly higher than that observed in mice treated with control IgG2a isotype antibody (which was similar to no antibody depletion). Furthermore, depletion of NK1.1⁺ cells resulted in a higher number of tumor lung nodules than depletion of CD8⁺ cells. In contrast, the mean number of pulmonary nodules from mice depleted of

CD4⁺ T cells resembled the isotype controls, indicating that CD4⁺ T cells were not critical for this effect. Therefore, it was concluded that (Hariharan *et al.*, *supra*). CD8⁺ T cells are essential for generation of antigen-specific anti-tumor immunity by SINrep5-E7/Hsp70 RNA vaccine and (Berglund, PM *et al.*, 1997. *AIDS Res Hum Retroviruses* 13:1487-95) NK cells play an important role as well.

To investigate whether NK cells were significantly expanded in mice vaccinated with various RNA constructs, flow cytometry analysis was performed, evaluating CD3(-), NK1.1+ cells. The proportion of NK cells was markedly increased in mice vaccinated with each of the all constructs (E7/Hsp70, E7, Hsp70, and control plasmid) relative to naïve mice, indicating that the expansion of NK cells is not a response limited to the E7/Hsp70 vaccine (Figure 8).

Self-Replicating RNA Vaccines Induce Apoptosis

Self-replicating RNA vaccines have been shown to induce apoptotic changes following uptake by cells (Ying *et al.*, *supra*). We evaluated apoptosis in BHK21 cells transfected with various RNA vaccines. Percentages of apoptotic BHK21 cells were normalized for transfection efficiency. As shown in Figure 9, apoptosis was induced in all groups of BHK21 cells transfected with various of the RNA vaccines compared to two negative control groups (untransfected or electroporated without RNA). There were no significant difference between the different RNA vaccines. there A steady decline in apoptosis occurred from 24 hr to 72 hr after transfection (with SIN5-E7/Hsp70: 70.3±3.6% at 24 hr, 49.3±4.2% at 48 hr, 18.0±3.1% at 72 hr, p<0.001, one-way ANOVA). These results confirm that cells transfected with each of these self-replicating RNA vaccines undergo apoptotic changes.

Enhanced Presentation of E7 through the MHC Class I Pathway in Dendritic Cells Pulsed With Cells Transfected with SINrep5-E7/Hsp70 RNA

Enhanced E7-specific CD8⁺ T cell responses *in vivo* may occur as a result of presentation of E7 via the MHC class I pathway resulting from uptake of apoptotic cellular material expressing various E7 constructs by host APCs. An experiment was performed to characterize the MHC class I presentation of E7 in DCs "pulsed" with BHK21 cells that had been transfected with various self-replicating RNA constructs. As noted above, the transfection efficiency and E7 expression is similar in BHK21 cells transfected with the various E7-containing self-replicating RNAs. Here, transfected BHK21 cells were incubated with bone marrow-derived DCs to allow antigen uptake and processing by the DCs. These DCs were then used as target cells for killing by E7-specific CD8⁺ CTL. As shown in Figure 10, DCs incubated with BHK21 cells that had been transfected with SINrep5-E7/Hsp70 RNA were lysed to a higher degree than DCs incubated with BHK21 cells transfected with SINrep5-E7 RNA (p<0.001). These results suggested that the presence of Hsp70 in a fusion protein with E7 (that was "fed" to DCs in

the form of RNA-transfected BHK21 cells) resulted in more effective presentation of E7 by DCs (via the MHC class I pathway) to CD8⁺ T cells as compared to E7 protein alone.

DISCUSSION

A vaccine designated SINrep5-E7/Hsp70, administered *in vivo* significantly enhanced E7-specific CD8⁺ T cell responses compared to the SINrep5-E7 RNA vaccine lacking Hsp70. It is unlikely that this effect results from occurs improved direct MHC class I presentation of E7 to CTLs by the cells that actually express E7/Hsp70 -- a process known as "direct priming". Intramuscular delivery of RNA replicons is believed to deliver RNA into muscle cells, which are not "professional" APCs because they do not express co-stimulatory molecules that are important for efficient activation of T cells. Even if the various SINrep5 constructs are delivered to other cell types after i.m. administration, the self-replicating RNA eventually causes apoptosis of the cells it transfects (Frolov *et al.*, *supra*). The cell initially transfected are therefore unlikely to be efficient direct presenters of antigen.

Rather, the enhanced CD8⁺ T cell responses are likely a result of a process whereby apoptotic cells (and subcellular material) are endocytosed and processed by professional APCs via MHC class I pathways for more effective presentation to CD8⁺ T cells (Albert *et al.*, *supra*) Alternatively, apoptotic cells may release the chimeric E7/Hsp70 protein which is then taken up and processed by APCs via the MHC class I-restricted pathway (Srivastava, PK *et al.*, *Immunogenetics* 39:93-8; Arnold, D *et al.*, 1995. *J Exp Med* 182:885-9; Suto, R *et al.*, 1995. *Science* 269:1585-8). Hsp70 complexes are known to enter professional APCs by binding specifically to the cell surface followed by receptor-mediated endocytosis (Arnold-Schild, D *et al.*, 1999. *J Immunol* 162:3757-60). In recent investigations of receptors for heat shock proteins, CD91 was identified as the receptor for gp96, one member of the HSP family on APCs (Binder, RJ *et al.*, 2000 *Nat. Immunol* 2:151-155).

Another important factor for enhanced activation of antigen specific CD8⁺ T cells by chimeric Hsp70/E7 may be the biology of professional APCs, primarily DCs. Cho *et al.*, recently reported that a mycobacterial HSP fused to an antigen stimulates DCs to upregulate expression of MHC class I, class II and co-stimulatory molecules (Cho, BK 2000. *Immunity* 12:263-272). Thus, induction of DC "maturation" by Hsp70 linked to antigen may augment T cell activity, explaining the results described herein with the chimeric E7/Hsp70 RNA vaccine.

The present study demonstrated that depletion of NK cells reduced the antitumor effect induced by the E7/Hsp70 RNA replicon-based vaccine (Figure 7), indicating that these cells are a necessary component. However, NK cell activity alone cannot account for the observed antitumor effect because other of the RNA replicon-based compositions produced a similar change in NK cell number (Figure 8). The *in vivo* antibody depletion study suggested that CD8⁺ CTLs were important for this antitumor effect

(Figure 7). Thus, it was concluded that both NK cells and CD8⁺ T cells are important contributors to the antitumor effect of the E7/Hsp70 RNA vaccine. Interactions among these two cell populations might also be of interest in understanding the outcome of such vaccination.

A comparison of the study described above with previous studies of the present inventors and their colleagues reveals that different forms of nucleic acid vaccines may activate different subsets of effector cells in the vaccinated host and act via different mechanisms.

Even though NK cells appeared to play a role in the anti-tumor effects induced by E7/Hsp70 RNA replicons, NKs were not essential when the vaccine was a naked E7/Hsp70 DNA vaccine. Thus,, depletion of NK1.1⁺ cells in mice vaccinated with naked E7/Hsp70 DNA did not decrease the anti-tumor immunity (Chen *et al.*, *supra*).

In contrast, CD8⁺ T cells were important for antitumor effects induced by both E7/Hsp70 DNA and E7/Hsp70 RNA replicon-based vaccines.

The apoptotic changes promoted by the self-replicating RNA vaccine raise potential safety concerns. With RNA replicon-based vaccines, increased apoptotic changes and inflammatory responses are localized to the injection sites. However, microscopic examination of the vital organs of E7/Hsp70-vaccinated mice did not show any significant histopathological changes. Potential risks attend the presence of HPV-16 E7 protein in host cells as E7 happens to be a viral oncoprotein that disrupts cell cycle regulation by binding to tumor suppressor pRB protein in nuclei (Lukas, JH *et al.*, 1994. *J Cell Biol* 125:625-38), leading to potential incidence and accumulation of genetic aberrations and eventual malignant transformation. Use of a self-replicating RNA vector eases the concern about oncogenicity of E7 protein since the transfected cells eventually undergo apoptosis.

In summary, the results reveal that fusion of DNA encoding *Mtb* Hsp70 to DNA encoding HPV-16 E7 in an RNA replicon results in a vaccine composition that induces a marked antigen (E7)-specific CD8⁺ T cell-mediated immune response that produces a state of anti-tumor immunity against tumors expressing the antigen. Fusion of Hsp70 DNA to DNA encoding an antigen further enhances the potency of the RNA replicon-based vaccine. These findings are applicable to other tumors and types of cancer where tumor-specific antigens can be identified. Further, these findings are directly applicable to vaccines against organisms responsible for infectious diseases such as viruses, protozoa, fungi and bacteria.

EXAMPLE II

Enhancement of Suicidal DNA Vaccine Potency by Linking *Mycobacterium Tuberculosis* Heat Shock Protein 70 to an Antigen

Recently, RNA replicon vaccines have emerged as an important strategy to alleviate the concerns for potential chromosomal integration and cell transformation noted above (Ying *et al.*, *supra*). These vaccines are self-replicating, self-limiting and may be administered either as RNA or as DNA which is then transcribed into RNA replicons in transfected cells *in vitro* or *in vivo*. DNA-based self-replicating RNA replicons, also known as "suicidal DNA," eventually cause lysis of transfected cells (Berglund *et al.*, *supra*; Leitner *et al.*, *supra*).

The present vaccine was developed using the Semliki Forest virus suicidal DNA vector, pSCA1 (DiCiommo, DP *et al.*, *J Biol Chem* 1998; 273:18060-6). Such vectors alleviate some concern about naked DNA because they eventually cause apoptosis of transfected cells. This feature is particularly desirable for vaccines that encode potentially oncogenic proteins, such as the HPV E7 protein (Wu, TC. *Curr Opin Immunol* 1994; 6:746-754). Because suicidal DNA vectors eventually kill transfected cells, any expression of DNA from these vectors is necessarily transient, conceivably compromising their potency. Therefore, the present inventors have conceived of strategies to enhance the potency of suicidal DNA vaccines.

Disclosed herein are the findings of an investigation of the impact of linking full-length Hsp70 to E7 on the potency of suicidal DNA vaccines. The suicidal DNA vaccine, pSCA1-E7/Hsp70, significantly increased expansion and activation of E7-specific CD8⁺ T cells compared to a vaccine comprising only pSCA1-E7. This enhanced response resulted in potent anti-tumor immunity against E7-expressing tumor cells.

Materials and Methods

Plasmid DNA Constructs and Preparation

pSCA1 vector (DiCiommo *et al.*, *supra*) was a gift from Dr. Bremner at the University of Toronto. This vector contains the human CMV immediate early gene (HCMV IE) promoter upstream of the Semliki Forrest virus (SFV) replicon. The subgenomic promoter is located after the SFV replicon, upstream of a multiple cloning sites for insertion of DNA of interest. For the generation of pSCA1-E7, E7 was cut from pcDNA3-E7 by BamHI/PmeI (Chen *et al.*, *supra*) and cloned into BamHI/SmaI sites of pSCA1. To construct pSCA1-Hsp70, Hsp70 was cut from pcDNA3-Hsp70⁸ by BamHI/PmeI and cloned into BamHI/SmaI sites of pSCA1. For the generation of pSCA1-E7/Hsp70, E7/Hsp70 DNA was cut from pcDNA3-E7/Hsp70 (Chen *et al.*, *supra*) by BamHI/PmeI and cloned into BamHI/SmaI sites of

pSCA1. The accuracy of these constructs was confirmed by DNA sequencing. Plasmid DNA vectors encoding pSCA1-E7, pSCA1-Hsp70, pSCA1-E7/Hsp70 or pSCA1 with no insert were transfected into subcloning efficient DH5TM cells (Life Technologies, USA). The DNA was then amplified and purified (Chen *et al.*, *supra*). The integrity of plasmid DNA and the absence of *Escherichia coli* DNA or RNA were checked in each preparation using 1% agarose gel electrophoresis. DNA concentration was determined by optical density measured at 260 nm. The presence of the inserted E7 fragment was confirmed by restriction enzyme digestion and gel electrophoresis.

Cell Lines

The production and maintenance of TC-1 cells has been described previously (Lin *et al.*, *supra*). In brief, HPV-16 E6, E7 and *ras* oncogene DNA were used to transform primary lung epithelial cells from C57BL/6 mice. The cells were grown in RPMI 1640, supplemented with 10% (vol/vol) fetal bovine serum, antibiotics, L-glutamine, sodium pyruvate, nonessential amino acids at 37°C with 5% CO₂. On the day of tumor challenge, TC-1 cells were harvested by trypsinization, washed twice with 1X Hanks buffered salt solution (HBSS) and finally resuspended in 1X HBSS to the designated concentration for injection.

Mice

6- to 8-week old female C57BL/6 mice from the National Cancer Institute (Frederick, MD) were purchased and kept in the oncology animal facility of the Johns Hopkins Hospital (Baltimore, MD). All animal procedures were performed according to approved protocols and in accordance with recommendations for the proper use and care of laboratory animals.

DNA Vaccination

Gene gun particle-mediated DNA vaccination was performed using a helium-driven gene gun (Bio-rad, Hercules, CA) according to the protocol provided by the manufacturer. Briefly, DNA coated gold particles were prepared by combining 25 mg of 1.6 µm gold microcarriers (Bio-rad, Hercules, CA) and 100 µl of 0.05 M spermidine (Sigma, St, Louis, MO). Plasmid DNA (50 µg) and 1.0 M CaCl₂ (100 µl) were added sequentially to the microcarriers while mixing by vortex. This mixture was allowed to precipitate at room temperature for 10 minutes. The microcarrier/DNA suspension was then centrifuged (10,000 rpm. for 5 sec) and washed 3 times in fresh absolute ethanol before resuspending in 3 ml of polyvinylpyrrolidone (0.1 mg/ml) (Bio-rad, Hercules, CA) in absolute ethanol. The solution was then loaded into tubing and allowed to settle for 4 min. The ethanol was gently removed and the microcarrier/DNA suspension was evenly attached to the inside surface of the tubing by rotating the tube. The tube was then dried using 0.4 liters per minute of flowing nitrogen gas. The dried tubing coated with microcarrier/DNA was then cut to 0.5-inch cartridges and stored in a capped dry bottle at

4°C. As a result, each cartridge contained 1 µg of plasmid DNA and 0.5 mg of gold. The DNA coated gold particles (1 µg DNA/bullet) were delivered to the shaved abdominal region of the mice using a helium-driven gene gun (Bio-rad, Hercules, CA) with a discharge pressure of 400 p.s.i.

Intracytoplasmic Cytokine Staining and Flow Cytometry Analysis

5 Splenocytes from groups naïve or vaccinated mice (5 mice per group) were collected and pooled one week after the last vaccination and incubated either with the E7 peptide (aa 49-57, RAHYNIVTF; SEQ ID NO:22) containing MHC class I epitope or the E7 peptide (aa 30-67) containing MHC class II peptide. The E7 peptide was added at a concentration of 2 µg/ml for 20 hours. To detect E7-specific CD8⁺ T cell precursors and E7-specific CD4⁺ T helper cell responses, CD8⁺ CTL epitopes aa 49-57 or aa 10 30-67 of E7 were used, respectively. Golgistop (Pharmingen, San Diego, CA) was added 6 hr before harvesting the cells from the culture. Cells were then washed once in FACScan buffer and stained with phycoerythrin (PE)-conjugated monoclonal rat anti-mouse CD8 or CD4 antibody (Pharmingen, San Diego, CA). Cells were subjected to intracellular cytokine staining using the Cytofix/Cytoperm kit according to the manufacturer's instructions (Pharmingen). FITC-conjugated anti-IFN-γ or anti-IL-4 15 antibodies and the immunoglobulin isotype control antibody (rat IgG1) were all purchased from Pharmingen. Analysis was done on a Becton-Dickinson FACScan flow cytometer with CELLQuest software (Becton Dickinson Immunocytometry System, Mountain View, CA).

ELISA for anti-E7 antibody (see EXAMPLE I)

In vivo Tumor Protection

20 These studies were generally as Example I, except that different DNA preparations and deliver was used: gene gun with 2 µg of pSCA1-E7, pSCA1-Hsp70, pSCA1-E7/Hsp70 DNA, or pSCA1 without insert. One week later, mice were boosted with the same regimen and, on day 14, were challenged subcutaneously with 10⁴ TC-1 tumor cells in the right leg.

Tumor Therapy

25 The tumor cells and DNA vaccines were prepared as above. Each mouse (5 per group) was challenged i.v. with 10⁴ TC-1 tumor cells on day 0. Three days later, mice were given 2 µg of a vaccine preparation (pSCA1-E7, pSCA1-Hsp70, pSCA1-E7/Hsp70 DNA, or pSCA1 without insert) via gene gun. One week later, animals were boosted using the same regimen and were sacrificed on day 21. The number of tumor nodules on the surface of the lung of each mouse were evaluated and counted by 30 experimenters blinded to sample identity. Statistical significance was tested using one-way ANOVA.

In vivo Antibody Depletion Experiments

The procedure was done as in Example I. Here, vaccination was with 2 µg DNA via gene gun, boosted one week later, and challenged with 5×10^4 TC-1 tumor cells. Depletion treatment was terminated 40 days after tumor challenge.

5 RESULTS

Construction and Characterization of the pSCA1-E7/Hsp70 Suicidal DNA Vaccine

The generation of plasmid DNA constructs and subsequent preparation of DNA-based self-replicating pSCA1 vaccines was performed as described above. The pSCA1 vector includes the HCMV IE promoter and a replicon from the SFV (DiCiommo *et al, supra*). A schematic diagram depicting DNA-based self-replicating pSCA1-E7, pSCA1-Hsp70, and pSCA1-E7/Hsp70 constructs is shown in **Figure 11**.

An ELISA to test expression of E7 protein by BHK21 cells transfected with the various DNA-based self-replicating E7-containing pSCA1 DNA constructs showed that similar amounts of E7 protein were expressed by each of these constructs.

Vaccination with the pSCA1-E7/Hsp70 Suicidal DNA Vaccine Enhances E7-Specific CD8⁺ T Cell-mediated Immune Responses

CD8⁺ T lymphocytes are important effectors of anti-tumor immunity. As a measure of the E7-specific CD8⁺ T cell response generated by the DNA-based self-replicating pSCA1-E7/Hsp70 vaccine, intracellular cytokine staining was evaluated in splenocytes from mice vaccinated intradermally via gene gun. As shown in **Figure 12A**, vaccination of mice with pSCA1-E7/Hsp70 suicidal DNA vaccine generated the highest number of E7-specific IFN-γ⁺ CD8⁺ T cell precursors (40 per 3×10^5 splenocytes) compared to vaccination with pSCA1-E7 DNA (12 per 3×10^5 splenocytes) ($p < 0.01$). pSCA1-E7/Hsp70 DNA immunization led to a nearly 4-fold increase in the number of E7-specific CD8⁺ T cell precursors. The mean number of IFN-γ-producing E7-specific CD8⁺ T cells was determined in the presence (solid columns) and absence (open columns) of E7 peptide aa 49-57 (SEQ ID NO:22) and shown in **Figure 12B**. These results indicated that linkage of Hsp70 to E7 significantly enhanced the frequency of E7-specific CD8⁺ T cell precursors in vaccinated mice.

Vaccination with pSCA1-E7/Hsp70 Suicidal DNA Vaccine Did not Induce Significant E7-Specific T Cell Responses Mediated by CD4⁺ Cells

To examine if various pSCA1 suicidal DNA vaccines stimulated E7-specific CD4⁺ T cell precursors to produce cytokines, double staining flow cytometry for surface CD4 and intracellular IFN-γ or IL-4 was performed to enumerate CD4⁺ cytokine secreting cells in splenocytes from vaccinated mice. Figures 13A and 13B show no significant difference in the number of E7-specific IFN-γ-secreting (or IL-

4-secreting) CD4⁺ cells among the various groups. Thus, linkage of Hsp70 to E7 in a suicidal DNA vaccine did not lead to stimulation of E7-specific CD4⁺ T cell precursors *in vivo*.

Vaccination with pSCA1-E7/Hsp70 Suicidal DNA Vaccine Did Not Induce Anti-E7 Antibodies

The quantity of anti- E7 antibodies in the sera of the vaccinated mice was determined by direct ELISA two weeks after vaccination. Sera of the mice vaccinated with pSCA1-E7/Hsp70 did not have higher titers of E7-specific antibodies compared to those mice vaccinated with pSCA1-E7 vaccine (Figure 14).

Vaccination with pSCA1-E7/Hsp70 Suicidal DNA Vaccine Protects Mice Better Against the Growth of E7-Expressing TC-1 Tumors

An *in vivo* tumor protection experiments was performed using DNA-based self-replicating pSCA1-E7/Hsp70 DNA vaccine and an E7-expressing tumor, TC-1, in C57BL/6 mice. As shown in **Figure 15**, 80% of mice receiving this vaccine remained tumor-free 70 days after TC-1 challenge. In contrast, all mice receiving wild-type pSCA1-E7 and pSCA1-Hsp70 constructs as well as all naïve mice developed tumors within 2 weeks. Therefore, the DNA-based self-replicating pSCA1-E7/Hsp70 vaccine significantly enhanced anti-tumor immunity.

Treatment with pSCA1-E7/Hsp70 Suicidal DNA Vaccine Eradicates Established Tumors in the Lungs

To determine the therapeutic potential of the DNA-based self-replicating pSCA1-E7/Hsp70 vaccine for lung metastases, each mouse was challenged with 10⁴ TC-1 tumor cells i.v. Results are shown in Figure B/6A as the number of pulmonary metastatic tumor nodules \pm SEM. Mice treated with the pSCA1-E7/Hsp70 suicidal DNA vaccine had the lowest number of pulmonary nodules (1.8 ± 0.5) compared to mice vaccinated with wild-type pSCA1-E7 (47.7 ± 4.6), pSCA1-Hsp70 (58.3 ± 1.8), pSCA1 alone (69.0 ± 4.9) or naïve mice (129.5 ± 4.0) (ANOVA, $p < 0.001$). Representative photographs of the lung tumors (unmagnified) are shown in Figure 16. These results indicate that the linkage of Hsp70 to E7 in a suicidal DNA vaccine significantly enhanced the antitumor therapeutic effect.

CD8⁺ T Cells are Essential for Antitumor Effects

To determine the types of lymphocytes required for protection against E7-expressing tumors, *in vivo* antibody depletion experiments were done. As shown in Figure 17, tumors grew out within 10 days of inoculation in all naïve mice and all mice depleted of CD8⁺ T cells. In contrast, 80% of the non-depleted mice remained tumor free 40 days after tumor challenge. Tumor grew within two weeks in 40% of mice depleted of CD4⁺ or of NK1.1⁺ cells. These results suggest that CD8⁺ T cells are essential for E7-specific anti-tumor immunity induced by the pSCA1-E7/Hsp70 suicidal DNA vaccine. CD4⁺ and NK1.1⁺ contributed to a lesser degree to the total antitumor effect.

DISCUSSION

The inventors have demonstrated that linkage of Hsp70 to E7 significantly enhanced the potency of an E7-expressing DNA that was administered as a suicidal DNA vaccine based on a replicating RNA replicon. Such suicidal DNA incorporating Hsp70 fused to E7 generated potent E7-specific CD8⁺ T cell-mediated immunity. Furthermore, this chimeric pSCA1-E7/Hsp70 suicidal DNA vaccine successfully prevented lethal pulmonary metastases in an experimental metastasis model.

Stimulation of CD8⁺ T cell activity is important in antitumor immune responses. (For review, see Chen, CH *et al.*, *J Biomed Sci* 1998, 5:231-52; Pardoll, DM *Nat Med* 1998, 4:525-31). Such immunity was augmented by administration of a pSCA1-E7/Hsp70 suicidal DNA vaccine and was manifest as protection against tumor growth and as therapy of a pre-existing tumor. The importance of this T cell subset is emphasized by the fact that depletion of CD8⁺ CTLs abolished this effect. Activated CTL function as effector cells that kill tumor cells directly or through the release of cytokines that interfere with tumor cell growth or survival. Therefore, the enhanced antigen-specific antitumor CD8⁺ T cell activity is critical to the potency of the pSCA1-E7/Hsp70 vaccine.

One mechanism by which E7-specific CD8⁺ T cell responses are stimulated *in vivo* is the direct MHC class I-restricted presentation of E7 to CD8⁺ T cells by APCs that express E7/Hsp70. This is known as "direct priming". However, because the suicidal DNA vaccine eventually results in the apoptosis of the very cells it transfects, direct priming by directly transfected APCs is unlikely to be effective.

Rather the enhanced CD8⁺ T cell responses observed in pSCA1-E7/Hsp70-vaccinated mice is likely a result of "cross priming," (Huang, AY *et al.*, *Science* 1994; 264:961-5) whereby cells expressing the E7/Hsp70 vaccine DNA release the antigen as an exogenous protein that is subsequently taken up and processed by other APCs via the MHC class I-restricted pathway. Cross-priming is the most likely mechanism for the enhanced CD8⁺ T cell activity because the suicidal DNA composition lyses transfected cells (Frolov *et al.*, *Proc Natl Acad Sci U S A* 1996; 93:11371-7) leading to release of antigen which becomes available to other APCs. Previous studies reported that Hsp70 linked to malaria peptide (NANP)₄₀ (Barrios, C *et al.*, *Clin Exp Immunol* 1994; 98:229-33), HIV-1 p24 (Suzue *et al.*, *supra*), ovalbumin (Suzue, K *et al.*, *Proc Natl Acad Sci U S A* 1997; 94:13146-51), or influenza nucleoprotein (Anthony *et al.*, *supra*) and administered as an exogenous protein enhanced MHC class I presentation of the linked antigens. Hsp70 fusion proteins are likely taken up by professional APCs which are known to be important in presenting exogenous Hsp70-associated antigens through the MHC class I pathway (Mitchell, DA *et al.*, *Eur J Immunol* 1998; 28:1923-33; Suto, R *et al.*, *Science* 1995; 269:1585-8). It was suggested that Hsp70 complexes can enter professional APCs via receptor-mediated endocytosis (Arnold-Schild *et al.*, *supra*). *Mtb* HSP protein fused to antigen stimulated DCs *in vitro* and *in vivo* to

upregulate the level of MHC class I, MHC class II and co-stimulatory molecules (Cho *et al.*, *supra*). According to the present invention, the lytic effect of the pSCA1 vector, the enhancement of MHC class I processing, the maturation of DCs, all of which are induced by Hsp70 fused to the antigen, all contribute to augmentation of CD8⁺ T cell activity resulting from the pSCA1-E7/Hsp70 vaccine via a cross-priming pathway.

Although use of suicidal DNA vectors and their induction of apoptosis alleviate some concerns about DNA vaccine integration into the host genome, the potency of such vaccines may be limited because of that same apoptotic outcome. The present inventors and their colleagues previously demonstrated that linkage of *Mtb* Hsp70 to E7 antigen enhanced the potency of a conventional naked DNA vaccine. (Chen *et al.*, 2000, *supra*). Here, the inventors have successfully extended the chimeric Hsp70 strategy to a suicidal DNA vector. However, the DNA-based RNA replicon vector approach appeared to be less efficacious than the conventional DNA vector approach in generating E7-specific CD8⁺ T cells. For example, approximately 130 E7-specific CD8⁺ T cells were generated per 10⁶ splenocytes when vaccinating mice with the pSCA1-E7/Hsp70 composition. Meanwhile, the previous study of chimeric E7/Hsp70 in a conventional mammalian expression plasmid (pcDNA3) generated about 430 E7-specific CD8⁺ T cells per 10⁶ splenocytes in vaccinated mice. (Chen *et al.*, *supra*). Although DNA-based replicons may be expected produce more E7 than do conventional DNA plasmids because of their self-replicating nature, Leitner *et al.* (*supra*) showed that replicon-based DNA plasmids did not produce more antigen. Furthermore, the apoptotic outcome of transfection with a DNA-based replicon may limit direct presentation of antigen by transfected APCs to CD8⁺ T cells, also contributing to lower vaccine potency.

This induction of apoptosis also raises concerns about potential tissue damage of the administration of such a vaccine. However, here, microscopic examination of the vital organs of E7/Hsp70-vaccinated mice did not reveal any significant histopathological changes.

Another risk is the presence of E7 protein in host cells (since E7 is a viral oncoprotein that disrupts cell cycle regulation by binding to tumor suppressor pRB protein in nuclei) leading to potential incidence and accumulation of genetic aberrations and eventual malignant transformation. Use of the suicidal DNA vector eases the concern about oncogenicity of E7 protein since the transfected cells eventually undergo apoptosis. Oncogenicity of E7 can be further reduced by introducing mutations into E7 DNA that eliminate binding of the E7 protein to pRB (Heck, DV *et al.*, *Proc Natl Acad Sci USA* 1992; 89:4442-6) while the cells still maintain most of their antigenicity.

In summary, the results revealed that fusion of DNA encoding *Mtb* Hsp70 to DNA encoding HPV-16 E7 in a suicidal DNA vaccine resulted in a vaccine that induced marked antigen (E7)-specific

CD8⁺ T cell-responses manifest as a state of anti-tumor immunity against tumors expressing the antigen. Since a majority of cervical cancers express HPV E7, the present is useful for controlling of HPV-associated tumors. These findings are applicable to other tumors and types of cancer where tumor-specific antigens can be identified. Further, these findings are directly applicable to vaccines against organisms responsible for infectious diseases such as viruses, protozoa, fungi and bacteria. Because the DNA-based RNA replicon vaccines are stable and easy to prepare in mass quantities, such vaccines are particularly desirable in developing countries which have high prevalence of HPV-associated cervical malignancy while lacking facilities for storing biological agents

EXAMPLE III

Enhancement of DNA Vaccine Potency by Linking DNA Encoding Antigen to DNA Encoding the Extracellular Domain of Flt3-Ligand

Prior to the present invention, To date, FL had not been used as part of a chimeric DNA vaccine. The present inventors and their colleagues investigated whether linking a full-length E7 DNA to DNA encoding the ECD of FL would enhance the potency of a DNA vaccine. They chose HPV-16 E7 as a model antigen for vaccine development (see above).

Studies were done to compare DNA vaccines containing wild-type E7 with DNA vaccines containing full-length E7 fused to FL for their stimulation of immune responses and their ability to protect animals against growth or metastasis of E7-expressing tumors (Lin *et al.*, *supra*). The results presented below indicate that linking DNA encoding the ECD of FL to E7 dramatically increased the expansion and activation of E7-specific CD8⁺ T cells, completely bypassing the CD4 arm. This strategy led not only to enhanced E7-specific CD8⁺ T cell responses, but also to potent anti-tumor immunity against established metastatic tumors expressing E7.

MATERIALS AND METHODS

Plasmid DNA Constructs and Preparation

pcDNA3 was used as an expression vector (instead of a previously described pCMV-Neo-Bam vector (Chen *et al.*, *supra*). The pcDNA3 expression vector was selected since it was used effectively to investigate the correlation between the E7-specific T cell responses with the antitumor effects produced various DNA vaccines. The production of HPV-16 E7-expressing plasmid, pcDNA3-E7 has been described previously (Chen *et al.*, *supra*).

For making the plasmid encoding the ECD of mouse FL, pcDNA3-FL, the DNA fragment encoding the signal peptide and ECD of mouse FL was first amplified with PCR using conditions described previously (Chen, CH *et al.*, *Cancer Research.* 60:1035-1042., 2000) with a mouse FL DNA

template, sfHAV-EO410 (ATCC, Manassas, VA) and a set of primers:

5'-gggtctagaatgacagtgtgtggcgccagc-3' [SEQ ID NO:28] and 5'-gggggatccctgcctgggcccaggctctgg-3' [SEQ ID NO:29].

The amplified product was digested with XbaI and BamHI and further cloned into the XbaI and BamHI cloning sites of pcDNA3 vector (Invitrogen, Carlsbad, CA).

For making pcDNA3-FL-E7, the E7 DNA fragment was isolated from pcDNA3-E7 by digestion with BamHI and HindIII and gel-

recovered. The isolated fragment was cloned into the BamHI and HindIII cloning sites of pcDNA3-FL. For making pcDNA3-GFP, a DNA fragment encoding the green fluorescent protein (GFP) was first

amplified in PCR using pEGFPN1 DNA (Clontech, Palo Alto, CA) and a set of primers: 5'-atcggatccatggtgagcaagggcgaggag-3' [SEQ ID NO:30] and 5'-gggaagctttactgtacagctctgcatg-3' [SEQ ID NO:31].

The amplified product was digested with BamHI and HindIII and cloned into the BamHI and HindIII cloning sites of pcDNA3 (Invitrogen). To make pDNA3-E7-GFP, the DNA fragment encoding E7 first amplified with PCR using pcDNA3-E7 as template and a set of primers: 5'-

ggggaattcatgcatggagatacaccta-3' [SEQ ID NO:32] and 5'-ggtggatccttgagaacagatgg-3' [SEQ ID NO:33]. The amplified product was digested with EcoRI and BamHI and cloned into the EcoRI and BamHI

cloning sites of pcDNA3-GFP. For making pcDNA3-FL-E7-GFP, the DNA encoding the signal peptide and ECD of FL was amplified with PCR using pcDNA3-FL as a DNA template and a set of primers:

5'-gggtctagaatgacagtgtgtggcgccagc-3' [SEQ ID NO:34] and 5'-cgagaattcctgcctgggcccaggctctg-3' [SEQ ID NO:35]. The amplified product was digested with XbaI and EcoRI and cloned into the XbaI and EcoRI cloning sites of pcDNA3-E7-GFP vector. The accuracy of these constructs was confirmed by

DNA sequencing. pcDNA3 DNA with FL, E7, FL-E7, E7-GFP or FL-E7-GFP inserts and the "empty" plasmid, pcDNA3 were transfected into subcloning-efficient DH5 α TM cells (Life Technologies, USA). The DNA was then amplified and purified (Chen *et al.*, *supra*). The integrity of plasmid DNA and the

absence of *E. coli* DNA or RNA was checked in each preparation using 1% agarose gel electrophoresis. DNA concentration was determined by the optical density measured at 260 nm. The presence of inserted

E7 fragment was confirmed by restriction enzyme digestion and gel electrophoresis.

The schematic domain structure of the Flt3-ligand protein and FL-E7 fusion peptide are shown in Figure 18A. The sequence of the FL-E7 construct, comprising the ECD of FL is shown in Figure 18B (SEQ ID NO:11 and 12). Residues 1-189 are FL-derived, residues 191-287 are E7-derived. The

remaining residues (*e.g.*, 288-302) are from the vector DNA.

Cell Lines

For description of TC-1 cells and their use, see above (and Lin *et al.*, *supra*). A human embryonic kidney cell line, 293, expressing MHC genes H-2D^b and H-2K^b (293 D^bK^b) (Bloom, MB *et al.*, *J Exp Med.* 185: 453-9, 1997) was a gift from Dr. JC Yang (National Cancer Institute, NIH,

Bethesda). These cells were grown in DMEM containing 10% heat-inactivated fetal calf serum, 0.3% glutamine, 0.01 M HEPES, 100 U/ml penicillin, 100 µg/ml streptomycin and 400 µg/ml G418.

The production and maintenance of TC-1 cells has been described previously (20). On the day of tumor challenge, TC-1 cells were harvested by trypsinization, washed twice with 1X Hanks buffered salt solution (HBSS) and resuspended in 1X HBSS to the designated concentration for injection. A human embryonic kidney cell line, 293, expressing the MHC genes H-2D^b and H-2K^b (293 D^bK^b) (24) was a gift from Dr. JC Yang (National Cancer Institute, NIH, Bethesda, MD). These cells were grown in DMEM containing 10% heat-inactivated fetal calf serum, 0.3% glutamine, 0.01 M HEPES, 100 U/ml penicillin, 100 µg/ml streptomycin and 400 µg/ml G418.

10 Confocal Fluorescence Microscopy

293 D^bK^b cells transfected with pcDNA E7-GFP and pcDNA FL-E7-GFP DNA were cultured for 24-36 hr, then cytocentrifuged onto glass slides. Cells were fixed with 4% paraformaldehyde in 1x PBS for 30 min at room temperature, permeabilized with 1x PBS containing 0.05% saponin and 1% BSA, and then incubated with mouse anti-calnexin mAb (Stressgen Biotechnologies, Victoria, Canada) at a concentration of 1 µg/ml for 30 min at room temperature. Unbound antibodies were removed by washing three times in 1X PBS. The cells were then incubated with Cy3-conjugated F(ab')₂ fragment of goat anti-mouse IgG (Jackson ImmunoResearch Laboratories) at a concentration of 10µg/ml for 30 min. The slides were washed with 1x PBS containing and 1% BSA. The glass slides were mounted with anti-fading medium, Mowiol 4-88 (Calbiochem Inc. La Jolla, CA) and covered with coverslips. Slides in which primary antibody was omitted were used as negative controls. Samples were examined on a confocal laser scanning microscope.

Mice

6- to 8-week-old female C57BL/6 mice from the National Cancer Institute (Frederick, MD) were purchased and kept in the oncology animal facility of the Johns Hopkins Hospital (Baltimore, MD). All animal procedures were performed according to approved protocols and in accordance with recommendations for the proper use and care of laboratory animals.

DNA Vaccination

Preparation of DNA-coated gold particles and gene gun particle-mediated DNA vaccination was performed using a helium-driven gene gun (Bio-rad, Hercules, CA) (see Chen *et al.*, *supra*).

30 Intracytoplasmic Cytokine Staining and Flow Cytometry Analysis

Pooled splenocytes from groups of naïve or vaccinated mice (5 mice per group) were incubated either with

- (a) the E7 peptide (aa 49-57) including an MHC class I epitope (Fettkamp *et al.*, *supra*) for detecting E7-specific CD8⁺ T cell precursors,, or
- (b) the E7 peptide (aa 30-67) containing the MHC class II peptide (Tindle *et al.*, *supra*) for detecting E7-specific CD4⁺ T helper cell precursors.

5 E7 peptide was added at 2µg/ml for 20 hours. Golgistop (PharMingen, San Diego, CA) was added 6 hours before harvesting cells from the culture. Cells were washed once in FACScan buffer and stained with phycoerythrin (PE)-conjugated monoclonal rat anti-mouse CD8 or CD4 antibody (PharMingen, San Diego, CA). Cells were subjected to intracellular cytokine staining using the Cytotfix/Cytoperm kit according to the manufacturer's instructions (PharMingen). FITC-conjugated anti-
10 IFN-γ antibody and the immunoglobulin isotype control antibody (rat IgG1) were all purchased from PharMingen. Analysis was done on a Becton Dickinson FACScan with CELLQuest software (Becton Dickinson Immunocytometry System, Mountain View, CA).

ELISA for anti-E7 antibody (see EXAMPLE I)

In vivo Tumor Protection

15 These studies were generally as Example I and II: gene gun with 2 µg of FL DNA, E7 DNA, FL-E7 DNA, FL mixed with E7 (FL+E7), or unvaccinated. One week later, mice were boosted with the same regimen and challenged subcutaneously on day 14 with 10⁴ TC-1 tumor cells in the right leg. Statistical analysis was performed using SAS version 6.12 (SAS Institute Inc., Cary, NC, USA). Percent of tumor free mice was analyzed using the Kaplan-Meier analysis. Statistical significance was tested
20 using log-rank statistics.

Tumor Therapy

The tumor cells and DNA vaccines were prepared as above. See Example II for description of tumor challenge. Three days later, mice were given 2 µg of a vaccine preparation ((FL DNA, E7 DNA, FL-E7 DNA via gene gun or were left unvaccinated)). One week later, animals were boosted using the
25 same regimen and were sacrificed on day 25. Lung tumor nodules were evaluated as in Example II. Statistical significance was tested using one-way ANOVA.

In vivo Antibody Depletion Experiments

The procedure was done as in Example II

Generation of DCs

30 DCs were generated by culturing bone marrow cells in the presence of GM-CSF as described previously (Fernandez, NC *et al.*, *Nat Med.* 5: 405-11, 1999). Briefly, bone marrow was collected from mouse femurs and tibias. Erythrocytes were lysed, and the remaining cells were passed through a

nylon mesh to remove small pieces of bone and debris. The cells were collected and 10^6 cells /ml were placed in 24-well plates in RPMI 1640 medium supplemented with 5% FCS, 2mM β -mercaptoethanol, 1% nonessential amino acids, 100 U/ml penicillin and 100 μ g/ml streptomycin (Life Technologies, Rockville, MD) and 100 U/ml GM-CSF (PharMingen, San Diego, CA). Two-thirds of the medium was replaced every 2 days. Non-adherent cells were harvested on day 7 and characterized by flow cytometry for DC markers as previously described (Wang, TL *et al.*, *Gene Therapy*. 7: 726-733., 2000).

Generation of E7-Specific CD8⁺ T Cell Lines

See Wang *et al.*, *supra*. Briefly, female C57BL/6 (H-2^b) mice were immunized intraperitoneally with vaccinia-Sig/E7/LAMP-1. Splenocytes were harvested on day 8. The cells were incubated with IL-2 (20 U/ml) and E7 peptide (aa 49-57) (1 μ M) for 6 days. Cells of the E7-specific CTL cell line were propagated in 24-well plates by mixing, in a final volume of 2 ml, (a) 10^6 splenocytes that included the E7-specific CTLs (b) 3×10^6 irradiated splenocytes (c) IL-2 (20 U/ml) and (d) E7 peptide (aa 49-57) at 1 μ M. This procedure was repeated every 6 days. The target-cell specificity of the E7 CTL line was characterized in a CTL assay. Flow cytometry was used demonstrate CD8 expression.

CTL Assay using Transfected 293 D^bK^b Cells as Target Cells

CTL assays were performed in 96-well round-bottom plates as described by Corr *et al.*, (Corr, M *et al.*, *J Immunol*. 163: 4721-7, 1999) and in Examples I and II. Transfected 293 D^bK^b cells were used as target cells while E7-specific CD8⁺ T cells served as effectors. 5×10^6 293 D^bK^b cells were transfected with 20 μ g of pcDNA3 (empty plasmid), E7, FL, or FL-E7 DNA vaccines with lipofectamine 2000 (Life Technologies, Rockville, MD) according to manufacturer's instructions. Cells were collected 40-44 hr after transfection. Levels of E7 protein expression, determined by ELISA, were similar in E7 and FL-E7-transfected cells. Cells were incubated and lysis measured as above.

CTL Assay Using DCs Pulsed with Lysates of Transfected 293 D^bK^b Cells as Target Cells

CTL assays using as targets DCs pulsed with cell lysates were generally in accordance with Uger, RA *et al.*, *J Immunol*. 160: 1598-605, 1998. Briefly, 293 D^bK^b cells were transfected as above and subjected to three freeze-thaw cycles. Protein concentrations were determined using the BioRad protein assay (Bio-Rad, Hercules, CA) using the vendor's protocol. The quantity of E7 protein was determined by ELISA. Cell lysates from E7 or FL-E7 DNA transfected 293 D^bK^b cells were standardized for E7 protein concentration.

DCs were prepared for use as target cells by pulsing 10^6 DCs with different concentrations of cell lysates (50, 10, 2 and 0.4 μ g/ml) in a final volume of 2 ml for 16-20 hrs. E7-specific CD8⁺ T cells were effector cells. CTL assays were performed at a fixed E/T ratio of 9 using 9×10^4 T cells and 10^4

prepared DC targets in a final volume of 200 μ l. Results were determined by measurements of LDH as above.

RESULTS

Linkage of the Extracellular Domain of FL to E7 Protein Re-routes E7 into the Endoplasmic Reticulum

To determine the expression and localization of wild-type E7 and E7 fusion proteins, DNA encoding the green fluorescent protein (GFP) was linked to the 3' end of E7 DNA and chimeric FL-E7 DNA as a tag. Transfection and subsequent examination by fluorescence microscopy was used to determine the expression and localization of wild-type and modified E7 protein. Levels of protein expression was quite similar between cells transfected with E7-GFP or FL-E7-GFP. As expected, cells transfected with the E7-GFP showed cytoplasmic/ nuclear distribution. In comparison, cells transfected with the chimeric FL-E7-GFP displayed a network pattern consistent with endoplasmic reticulum (ER) localization. To test whether the FL-E7-GFP chimera had in fact been distributed to the ER, cells were further stained with an antibody to calnexin and examined by immunofluorescence. Calnexin is a well-characterized marker for the ER. Co-localization of E7-GFP and calnexin was only observed in cells transfected with FL-E7-GFP but not E7-GFP, indicating that at least some of the FL-E7 fusion product but not E7-GFP was targeted to ER compartments. These results indicated that the addition of the ECD of FL to E7 facilitates the entry into ER compartments.

Vaccination with FL-E7 Fusion DNA Significantly Enhanced E7-Specific CD8⁺ T Cell Responses

CD8⁺ T lymphocytes are important effectors of anti-tumor immunity. As a measure of the E7-specific CD8⁺ T cell response generated by the FL-E7 DNA vaccine, intracellular IFN γ cytokine staining was evaluated in splenocytes from vaccinated mice. This is a sensitive functional assay for measuring IFN- γ production at the single-cell level (Murali-Kristna, K *et al.*, *Immunity*. 8: 177-87).

As shown in Figure 19A and B, vaccination of mice with FL-E7 DNA generated the highest number of E7-specific IFN- γ ⁺ CD8⁺ T cell precursors (940 per 3x10⁵ splenocytes) compared to vaccination with E7 DNA (12 per 3x10⁵ splenocytes) (p<0.01). Thus, FL-E7 DNA immunization led to a nearly 8-fold increase in the number of E7-specific CD8⁺ T cell precursors. These results also indicated that fusion of E7 to FL was required for this enhancement because vaccination with a mixture of FL-DNA and E7-DNA did not generate such enhanced CD8⁺ T cell activity.

Vaccination with FL-E7 Fusion DNA Did not Induce Significant E7-Specific CD4⁺ T Cell Responses or Anti-E7 Antibodies

E7-specific CD4⁺ T precursor cells secreting the cytokine IFN- γ or IL-4) were assessed by double staining for surface CD4 and intracellular IFN- γ or IL-4 using flow cytometry. Splenocytes were from immunized mice. Figure 20A shows that mice vaccinated with FL-E7 DNA developed no significant increase in

CD4⁺ IFN- γ ⁺ double positive cells compared to mice vaccinated with FL DNA, wild-type E7 DNA, plasmid DNA or unvaccinated naïve mice. Positive control splenocytes were from Sig/E7/LAMP-1 DNA vaccinated mice (Ji *et al.*, *supra*). Similarly, no significant increases in CD4⁺ IL-4⁺ double-positive cells were observed (FL-E7 vaccinated compared with FL DNA, wild-type E7 DNA, plasmid DNA or unvaccinated naïve mice (Figure 20B). IL-4-secreting activated mouse splenocytes (MiCK-2, PharMingen) were positive controls to assure successful intracellular IL-4 staining.

To determine the levels of E7-specific antibodies in the sera of the vaccinated mice, ELISA was performed 2 weeks after the last vaccination. No significant E7-specific antibody responses were detected in mice. Sera of the mice vaccinated with chimeric FL-E7 DNA did not have higher titers of E7-specific antibodies compared to mice vaccinated with FL, empty plasmids, or unvaccinated naïve mice.

Vaccination with Chimeric FL-E7 DNA Vaccine Protects Mice Better Against the Growth of E7-Expressing TC-1 Tumors

Results of an *in vivo* tumor protection study is shown in Figure 21. 100% of mice vaccinated with FL-E7 DNA remained tumor-free 70 days after challenge (log-rank, $p < 0.001$). In contrast, only 20% of mice receiving wild-type E7 remained tumor free after day 32 and all unvaccinated mice, or mice given FL DNA developed tumors within 20 days of challenge. Fusion of E7 to FL was required for generating protective immunity, since only 20% of mice vaccinated with a mixture E7 DNA and FL DNA) remained tumor free after 32 days. Therefore, FL-E7 fusion DNA significantly enhanced anti-tumor immunity.

Treatment with FL-E7 Fusion DNA Eradicates Established E7-expressing Tumors in the Lungs

To determine the therapeutic potential of a chimeric FL-E7 DNA construct in treating TC-1 lung metastases, each mouse was challenged with tumor cells i.v. Results are shown in Figure 22A as the mean number of pulmonary metastatic tumor nodules \pm SEM. Mice vaccinated with FL-E7 DNA had the lowest mean number of pulmonary nodules (5.8 ± 3.6) compared to mice vaccinated with wild-type E7 DNA (67.5 ± 3.5), FL DNA mixed with E7 DNA (68 ± 15), FL DNA (65.0 ± 5.0) or unvaccinated mice (50.7 ± 7.3) (one-way ANOVA, $p < 0.001$). Figure 22B shows lung weights (mean \pm SEM in grams). Mice vaccinated with FL-E7 DNA had the lowest lung weight (0.158 ± 0.025) compared to mice vaccinated with wild-type E7 DNA (0.462 ± 0.02), FL DNA plus E7 DNA (0.469 ± 0.08), or FL DNA (0.6 ± 0.03), or unvaccinated mice (0.645 ± 0.08) (one-way ANOVA, $p < 0.001$). Representative photographs of the lung tumors are shown in Figure 23.

CD8⁺ T Cells But Not CD4⁺ T cells are Essential for Anti-tumor Effects

To determine the types of lymphocytes required for protection against E7-expressing tumors, *in vivo* antibody depletion experiments (Lin *et al.*, *supra*; Wu *et al.*, 1995, *supra*) were done. Depletion of lymphocyte subsets was assessed on the day of tumor injection, and weekly thereafter by flow cytometric analysis of spleen cells. More than 99% of cells of the appropriate subset were depleted in all cases without effect on the other subsets. As shown in **Figure 24**, 100% of unvaccinated mice and mice depleted of CD8⁺ T cells grew tumors within 14 days after challenge. In contrast, all non-depleted mice and all mice depleted of CD4⁺ T cells remained tumor-free 60 days after challenge. 40% of mice in which NK1.1⁺ cells were depleted grew tumors 6 weeks after challenge. Even though there appeared to have been an effect of NK cell depletion, the difference from controls is result was not statistically significant (log-rank, p=.13). It was concluded that CD8⁺ T cells are essential for E7-specific antitumor immunity induced by the FL-E7 DNA vaccine.

Enhanced Presentation of E7 via the MHC Class I Pathway in Cells Transfected with FL-E7 DNA

As noted earlier, mice vaccinated with FL-E7 generated the highest number of E7-specific CD8⁺ T cell precursors. To understand the mechanism underlying this effect, the inventors tested whether enhanced MHC class I presentation of E7 occurred in cells expressing FL-E7 (in this case, human embryonic kidney 293 cells D^bK^b transfected with FL-E7). CTL assays employing D^b-restricted E7-specific CD8⁺ effector T cells were used to determine if target 293 D^bK^b cells transfected with FL-E7 were killed more efficiently than 293 D^bK^b cells transfected with wild-type E7. 293 D^bK^b cells were selected because of their stable and high transfection efficiency (Bloom *et al.*, *supra*). In addition, levels of E7 expression in 293 D^bK^b cells transfected with FL-E7 DNA or E7 DNA were similar.

In the CTL assays, targets were 293 D^bK^b cells that had been transfected with either empty plasmid, FL DNA, E7 DNA, or FL-E7 DNA, or that were not transfected.. Effector cells were added to achieve various E/T ratios (1, 3, 9, 27). As shown in **Figure 25**, 293 D^bK^b cells transfected with FL-E7 DNA were lysed at a higher level than targets cells transfected with wild-type E7 DNA. Transfection with FL-E7 DNA thus resulted in more efficient presentation of E7 antigen via the MHC class I pathway.

Enhanced Presentation of E7 Through the MHC Class I Pathway in DCs Pulsed With Chimeric FL-E7 Protein

Enhanced E7-specific CD8⁺ T cell responses *in vivo* may occur as a result of presentation of E7 via the MHC class I pathway resulting from uptake of lysed cellular material expressing various E7 constructs by host APCs ("cross-priming").

A cross priming experiment was performed to characterize the MHC class I presentation of E7 of DCs pulsed with cell lysates of 293 D^bK^b cells transfected with empty plasmid, FL, E7, or FL-E7 DNA.

Lysates of transfected 293 D^bK^b cells were obtained by repeated cycles of freeze-thaw. 10⁶ bone marrow-derived DCs were pulsed with serial dilutions (50, 10, 2 or 0.4 µg) of lysate derived from 293 D^bK^b cells transfected with different constructs. These DCs were used as target cells for lysis by D^b-restricted E7-specific CD8⁺ CTL. CTL assays were performed at a fixed E/T ratio of 9. As shown in **Figure 26**, DCs pulsed with lysates from 293 D^bK^b cells transfected with FL-E7 DNA were lysed at a higher percentage compared to DCs pulsed with lysates from 293 D^bK^b cells transfected with the other DNA constructs and non-transfected DCs. It was concluded that DCs pulsed with FL-E7 fusion protein presented E7 antigen through the MHC class I pathway more efficiently than DCs pulsed with wild-type E7 protein. Thus, the fusion of FL to E7 enhanced E7-specific CD8⁺ T cell immune responses via cross priming effects.

DISCUSSION

The foregoing study demonstrated that linking the ECD of FL to E7 significantly enhanced the potency of E7-expressing DNA vaccines to induce potent CD8⁺ T cell-immune responses that were protective and therapeutic against E7-expressing tumors, such that mice were protected from both primary tumor growth and development of lethal pulmonary metastases.

The incorporation of FL into the vaccine preferentially enhanced CD8⁺ T cell responses vs. E7-specific CD4⁺ T cell responses that were not significantly changed. Linking FL to E7 in the vaccine directly enhanced MHC class I presentation of E7 (compared to a wild-type E7 vaccine) in transfected cells *in vitro*. Since biolistic DNA delivery introduces DNA directly into dermal professional APCs, FL-E7 DNA-transfected APCs may act by directly presenting E7 via the MHC class I pathway to CD8⁺ T cells *in vivo*.

Although it is not clear how this linkage directly enhances MHC class I presentation, one mechanism involves a chaperone effect of FL. When expressed in cells, FL may be distributed to the ER (Chklovoskaia, E *et al.*, *Blood*. 93: 2595-604, 1999). Fluorescence microscopic examination revealed that in cells transfected with FL-E7-GFP, most of the FL-E7-GFP fusion protein co-localized with calnexin in the ER, suggesting that this linkage facilitates entry of E7 into the ER. Several studies demonstrated that ER targeting can enhance antigen-specific MHC class I-restricted CTL activity (Shiver, JW *et al.*, *J Pharm Sci* 85: 1317-24, 1996; Hsu, SC *et al.*, *Int Immunol*. 10: 1441-7, 1998).

Another mechanism that may contribute to the present observations is "cross-priming," whereby lysis of cells expressing FL-E7 releases protein that is taken up and processed by other APCs via the

MHC class I-restricted pathway. The present results show that DCs pulsed with FL-E7 fusion protein are capable of presenting E7 antigen through the MHC class I pathway in a more efficient manner than DCs pulsed with wild-type E7 protein. (Figure 26). However, the “cross-priming” of chimeric FL-E7 probably does not play a major role in gene gun-mediated FL-E7 DNA vaccination. Direct priming, but not cross-priming, of CD8⁺ T cells by DNA-transfected DCs is the key event in gene gun-mediated DNA immunization (Porgador, A *et al.*, *J Exp Med.* 188: 1075-82, 1998; Akbari, O *et al.*, *J Exp Med.* 189: 169-78, 1999). However, the possibility of cross-priming cannot be ruled out, because FL-E7 might be released from other cell types, such as keratinocytes (which are also transfected by gene gun vaccination), and then enter DCs via a cross-priming mechanism.

No significant increases in the numbers of DCs or NK cells were detected in the spleens of mice vaccinated with FL-E7 DNA vaccines even though FL is known to expand these cell populations (Peron *et al.*, *supra*; Williams, NS *et al.*, *J Exp Med.* 186: 1609-14, 1997, Shaw, SG *et al.*, *J Immunol.* 161: 2817-24, 1998). This may be related to small amounts of FL-E7 present in the circulation after DNA vaccination. FL-E7 protein could not be detected in sera of mice vaccinated with FL-E7 DNA, which also raises a question about the source of FL-E7 protein for cross-priming. FL-E7 protein from the lysis of transfected keratinocytes may be taken up by Langerhans’ cells and further processed in the draining lymph nodes without entering the circulation in detectable quantities.

The E7 DNA vaccine described above had weaker antitumor effects compared to an E7 DNA vaccine using a different mammalian expression vector (Chen *et al.*, 1999, *supra* Ji *et al.*, *supra*). In these previous studies, a pCMV-Neo-Bam expression vector that includes the HCMV promoter was used. E7 DNA vaccine using that vector generated a very impressive antitumor effect in the relative absence of E7-specific CD8⁺ T cell responses. In the current study, a relatively weak E7-specific CD8⁺ T cell response and a relatively weak anti-tumor response were observed in mice vaccinated with E7 DNA in the form of the pcDNA3 vector. The discrepancy in the anti-tumor response evoked by the same DNA in a different vector may be explained simply by different levels of expression. Furthermore, bacterial DNA can contain immunostimulatory elements such as CpG islands (Sato, Y *et al.*, *Science.* 273: 352-4, 1996; Klinman, DM *et al.*, *J Immunol.* 158: 3635-9, 1997), which can cause simultaneous maturation and activation of DCs (Sparwasser, T *et al.*, *Eur J Immunol.* 28: 2045-54, 1998) thereby acting as an adjuvant for tumor immunization (Weiner, GJ *et al.*, *Proc Natl Acad Sci USA.* 94: 10833-7, 1997).

The FL-E7 DNA vaccine may raise certain safety concerns because DNA could integrate into the host genome, though it is estimated that the frequency of integration is much lower than that of spontaneous mutation and should not pose any real risk (Nichols, WW *et al.*, *Annals of NY Academy of Science.* 772: 30-39., 1995). The risks of HPV-16 E7 protein was discussed above. There is a concern about possible autoimmune effects resulting from excessive expansion of DCs *in vivo*. However, here,

no significant increase in the numbers of DCs was observed in the spleen and lymph nodes of mice vaccinated with FL or FL-E7 DNA vaccines. Examination of vital organs in all of the FL-E7-vaccinated mice did not reveal any significant gross or microscopic pathology. Therefore, FL-E7 can be used as a potent DNA vaccine without observable detrimental side effects.

5 In summary, fusion of DNA encoding the ECD of the FL protein to E7 DNA generated potent E7-specific CD8⁺ T cell responses and anti-tumor effects against E7-expressing tumors. Linkage of DNA encoding FL to DNA encoding an antigen enhances the potency of DNA vaccines and are applicable to other tumors and types of cancer where tumor-specific antigens can be identified. Further, these findings are directly applicable to vaccines against organisms responsible for infectious diseases
10 such as viruses, protozoa, fungi and bacteria.

The references cited above are all incorporated by reference herein, whether specifically incorporated or not. All publications, patents, patent applications, GenBank sequences and ATCC deposits, cited herein are hereby expressly incorporated by reference for all purposes.

5 Citation of the documents herein is not intended as an admission that any of them is pertinent prior art. All statements as to the date or representation as to the contents of these documents is based on the information available to the applicant and does not constitute any admission as to the correctness of the dates or contents of these documents.

WHAT IS CLAIMED IS:

1. A nucleic acid molecule encoding a fusion polypeptide useful as a vaccine composition, which molecule comprises:

- 5 (a) a first nucleic acid sequence encoding a first polypeptide or peptide that (i) promotes processing via the MHC class I pathway (MHC-I-PP) and/or (ii) promotes development or activity of an antigen presenting cell (APC);
- (b) optionally, fused in frame with the first nucleic acid sequence, a linker nucleic acid sequence encoding a linker peptide; and
- 10 (c) a second nucleic acid sequence that is linked in frame to said first nucleic acid sequence or to said linker nucleic acid sequence and that encodes an antigenic polypeptide or peptide.

2. The nucleic acid molecule of claim 1, wherein the antigenic peptide comprises an epitope that binds to a MHC class I protein.

15 3. The nucleic acid molecule of claim 2, wherein said epitope is between about 8 amino acid residues and about 11 amino acid residues in length.

4. The nucleic acid molecule of any of claims 1-3 wherein the first polypeptide or peptide is Hsp70, an active C-terminal domain thereof, or a functional derivative of Hsp70 or of said C-terminal domain.

20 5. The nucleic acid molecule of any of claims 1-3 wherein the first polypeptide or peptide is the Flt3 ligand (FL), the extracellular domain thereof, or a functional derivative of FL or of said extracellular domain.

6. The nucleic acid molecule of claim 4, wherein the first polypeptide is SEQ ID NO:4 or the full length sequence of Hsp70 as set forth in GENBANK Z95324 AL123456 and encoded by nucleotides 10633-12510 of the *Mycobacterium tuberculosis* genome.

25 7. The nucleic acid molecule of claim 4 wherein the first polypeptide consists essentially of said C-terminal domain having a sequence from about residue 517 to about the C-terminal amino acid residue of:

- (i) SEQ ID NO:4, or
- (ii) the full length native sequence of Hsp70 as set forth in GENBANK Z95324 AL123456
- 30 and encoded by nucleotides 10633-12510 of the *Mycobacterium tuberculosis* genome.

8. The nucleic acid molecule of claim 5, wherein the first polypeptide is FL.

9. The nucleic acid molecule of claim 5, wherein the first polypeptide polypeptide consists essentially of the sequence SEQ ID NO:10.

10. The nucleic acid molecule of any of claims 1-9 wherein the antigen is one which is present on, or/ cross-reactive with an epitope of, a pathogenic organism, cell, or virus.

11. The nucleic acid molecule of claim 10, wherein the virus is a human papilloma virus.

12. The nucleic acid molecule of claim 11, wherein the antigen is the E7 polypeptide of HPV-16 or an antigenic fragment thereof.

13. The nucleic acid molecule of claim 12, wherein the HPV-16 E7 polypeptide is non-oncogenic.

14. The nucleic acid molecule of claim 10, wherein the pathogenic organism is a bacterium.

15. The nucleic acid molecule of claim 10, wherein the pathogenic cell is a tumor cell.

16. The nucleic acid molecule of claim 15, wherein the antigen is a tumor-specific or tumor-associated antigen, or any antigenic epitope thereof.

17. The nucleic acid molecule of claim 16, wherein the antigen comprises the HER-2/neu protein or a peptide thereof, mutant p53 or a melanoma-associated antigens selected from the group consisting of MAGE-1, MAGE-3, MART-1/Melan-A, tyrosinase, gp75, gp100, BAGE, GAGE-1, GAGE-2, GnT-V, and p15.

18. The nucleic acid molecule of any of claims 1-17 operatively linked to a promoter.

19. The nucleic acid molecule of claim 18, wherein the promoter is one which is expressed in an APC.

20. The nucleic acid molecule of claim 21, wherein the APC is a dendritic cell.

21. An isolated nucleic acid molecule that, under stringent hybridization conditions, hybridizes simultaneously with:

- (i) at least part of said first nucleic acid sequence and at least part of said second nucleic acid sequence,
- (ii) at least part of said first nucleic acid sequence and part of said linker nucleic acid sequence,
- (iii) at least part of said second nucleic acid sequence and part of said linker nucleic acid sequence, or

(iv) at least part of said first nucleic acid sequence, at least part of said first nucleic acid sequence and said linker nucleic acid sequence,
said sequences in accordance with any of claims 1-20.

22. An expression vector comprising the nucleic acid molecule of any of claims 1-20
operatively linked to

(a) a promoter; and

(b) optionally, additional regulatory sequences that regulate expression of said nucleic acid in a eukaryotic cell.

23. The expression vector of claim 22 which is a viral vector or a plasmid.

24. The expression vector of claim 22 wherein said plasmid is pcDNA3 which is derived from the sequence, prior to insertion of the nucleic acid sequences encoding said first or said second polypeptides, of SEQ ID NO:15.

25. The expression vector of claim 22 which is a self-replicating RNA replicon.

26. The expression vector of claim 25, wherein the self-replicating RNA replicon is a Sindbis virus self-replicating RNA replicon.

27. The expression vector of any of claims 25 or 26, wherein the replicon is SINrep5 which is derived from the sequence, prior to insertion of the nucleic acid sequences encoding said first or said second polypeptides, of SEQ ID NO:16.

28. The expression vector of claim 22 which is a suicidal DNA vector.

29. The expression vector of claim 28 wherein said suicidal DNA vector is an alphavirus DNA vector.

30. The expression vector of claim 29 wherein said alphavirus is Semliki Forest virus (SFV).

31. The expression vector of claim 30 wherein said SFV vector is pSCA1.

32. The expression vector of any of claims 28-30 wherein the suicidal DNA is derived from the sequence, prior to insertion of the nucleic acid sequences encoding said first or said second polypeptides, of SEQ ID NO:17.

33. The expression vector of any of claims 22-32 wherein the first encoded polypeptide or peptide is Hsp70, an active C-terminal domain thereof, or a functional derivative of Hsp70 or of said C-terminal domain.

34. The expression vector of claim 33 wherein the first encoded polypeptide consists essentially of the sequence SEQ ID NO:4 or the full length sequence of Hsp70 a set forth in GENBANK Z95324 AL123456 and encoded by nucleotides 10633-12510 of the *Mycobacterium tuberculosis* genome.

5 35. The expression vector of claim 32 wherein the first polypeptide consists essentially of said C-terminal domain having a sequence from about residue 517 to about the C-terminal amino acid residue of:

(i) SEQ ID NO:4, or

10 (ii) the full length native sequence of Hsp70 a set forth in GENBANK Z95324 AL123456 and encoded by nucleotides 10633-12510 of the *Mycobacterium tuberculosis* genome.

36. The expression vector of an of claims 34, that is a self-replicating RNA replicon that also encodes HPV protein E7, and has the nucleotide sequence SEQ ID NO:19.

37. The expression vector of claim 34, that is a suicidal DNA vector which also encodes HPV protein E7, and has the nucleotide sequence of SEQ ID NO:20.

15 38. The expression vector of any of claims 22-32 wherein the first encoded polypeptide or peptide is the Flt3 ligand (FL), the extracellular domain thereof, or or a functional derivative of FL or of said extracellular domain.

39. The expression vector of claim 38 wherein the first encoded polypeptide consists essentially of the extracellular domain of FL having a sequence SEQ ID NO:10.

20 40. The expression vector of claim 39 that comprises a naked DNA plasmid pcDNA3 that includes the coding sequence for HPV protein E7 and the FL extracellular domain, and has the nucleotide sequence SEQ ID NO:21.

41. A cell which has been modified to comprise the nucleic acid or expression vector of any of claims 1-40.

25 42. The cell of claim 41 which expresses said nucleic acid molecule and said fusion polypeptide.

43. The cell of claim 41 or 42 which is an APC.

44. The cell of claim 43, wherein the APC is a dendritic cell, a keratinocyte, a macrophage, a monocyte, a B lymphocyte, a microglial cell, an astrocyte, or an activated endothelial cell.

45. A particle comprising the nucleic acid or expression vector of any of claims 1-40.

46. The particle of claim 45 which comprises a material is suitable for introduction into a cell or an animal by particle bombardment.

47. The particle of claim 46, wherein the material is gold.

48. A fusion or chimeric polypeptide comprising

(a) a first polypeptide or peptide that (i) promotes processing via the MHC class I pathway and/or (ii) promotes development or activity of an APC; and

(b) a second polypeptide comprising an antigenic peptide or polypeptide.

49. The fusion or chimeric polypeptide of claim 48, wherein the antigenic peptide or polypeptide comprises an epitope that binds to, and is presented on the cell surface by, MHC class I proteins.

50. The fusion or chimeric polypeptide of claim 48 or 49 wherein the first polypeptide and the antigenic polypeptide or peptide are linked by a chemical linker.

51. The fusion polypeptide of any of claims 48-50 wherein the first polypeptide is N-terminal to the second polypeptide.

52. The fusion polypeptide of any of claims 48-50 wherein the second polypeptide is N-terminal to the first polypeptide.

53. The fusion or chimeric polypeptide of any of claims 48-52 wherein the first polypeptide or peptide is Hsp70, an active C-terminal domain thereof, or a functional derivative of Hsp70 or of said C-terminal domain.

54. The fusion or chimeric polypeptide of any of claims 48-52 wherein the first polypeptide or peptide is the Flt3 ligand (FL), the extracellular domain thereof, or a functional derivative of FL or of said extracellular domain.

55. The fusion polypeptide of claim 53, wherein the first polypeptide is SEQ ID NO:4 or the full length sequence of Hsp70 as set forth in GENBANK Z95324 AL123456 and encoded by nucleotides 10633-12510 of the *Mycobacterium tuberculosis* genome.

56. The fusion polypeptide of claim 8, wherein the first polypeptide polypeptide consists essentially of the sequence SEQ ID NO:10.

57. The fusion polypeptide encoded by the nucleic acid molecule of any of claims 1-20.

58. The fusion polypeptide encoded by the expression vector of any of claims 22-40.

59. A pharmaceutical composition capable of inducing or enhancing an antigen-specific immune response, comprising:

- (a) pharmaceutically and immunologically acceptable excipient in combination with;
- (b) a composition selected from:
 - (i) the nucleic acid molecule or expression vector of any of claims 1-40;
 - (ii) the cell of any of claims 41-44;
 - (iii) the particle of any of claims 45-47;
 - (iv) the fusion or chimeric polypeptide of any of claims 48-58; or
 - (v) any combination of (i)-(iv).

60. A method of inducing or enhancing an antigen specific immune response in a subject comprising administering to the subject an effective amount of the pharmaceutical composition of claims 59, thereby inducing or enhancing said response.

61. The method of claim 60, wherein the response is mediated at least in part by CD8⁺ cytotoxic T lymphocytes (CTL).

62. The method of claim 60, wherein the response is mediated at least in part by antibodies.

63. The method of any of claims 60-62 wherein said subject is a human.

64. A method of inducing or enhancing an antigen specific immune response in cells or in a subject comprising contacting said cells with, or administering to said subject, an effective amount of the pharmaceutical composition of claim 59, thereby inducing or enhancing said response.

65. The method of claim 64, comprising contacting said cells *ex vivo* with said composition.

66. The method of claim 65 wherein said cells comprise APCs.

67. The method of claim 66, wherein said APCs are dendritic cells.

68. The method of claim 66 or 67, wherein the APCs or said dendritic cells are of human origin..

68. The method of any of claims 66-68, wherein the APCs are isolated from a living subject.

69. The method of any of claims 65-69, further comprising a step of administering said cells to which were contacted with the composition *ex vivo* to (i) a histocompatible subject or (ii) the subject from which said cells were obtained.

70. The method of any of claims 64-69 wherein said cells are human cells and said subject is a human.

71. The method of any of claims 60-64 and 69-70 wherein said administering is by a intramuscular, intradermal, or subcutaneous route.

5 72. The method of any of claims 60-64 and 69-70, wherein the composition comprises said nucleic acid molecule, said expression vector or said particle, and said administering is by biolistic injection.

10 73. The method of any of claims 60-64, 69-72 wherein the administering is intratumoral or peritumoral.

74. A method of increasing the numbers or lytic activity of CD8⁺ CTLs specific for a selected antigen in a subject, comprising administering to said subject an effective amount of the pharmaceutical composition of claim 59, wherein

15 (i) said nucleic acid molecule, said expressio vector, said cell, said particle or said fusion or chimeric polypeptide comprises said selected antigen, and
(ii) said selected antigen comprises an epitope that binds to, and is presented on the cell surface by, MHC class I proteins,
thereby increasing the numbers or activity of said CTLs.

20 75. A method of inhibiting growth or preventing re-growth of a tumor in a subject, comprising administering to said subject an effective amount of a pharmaceutical composition of claim 59, wherein said nucleic acid molecule, said expression vector, said cell, said particle or said fusion or chimeric polypeptide comprises one or more tumor-associated or tumor-specific epitopes present on said tumor in said subject, thereby inhibiting said growth or preventing said re-growth.

76. The method of claim 75, wherein said administering is intratumoral or peritumoral.

25 77. The method of any of claims 63 or 64, further comprising treating said subject with radiotherapy or chemotherapy.

1/28

RNA TRANSCRIPTS




FIG. 1

2/28




FIG. 2

3/28




FIG. 3

4/28




FIG. 4

5/28





FIG. 5


6/28



NO INSERT
FIG. 6A




HSP 70
FIG. 6B




E7
FIG. 6C



E7/HSP 70 (0.1 μg)
FIG. 6D



E7/HSP 70 (1 μg)
FIG. 6E



E7/HSP 70 (10 μg)
FIG. 6F

7/28




FIG. 7

8/28




FIG. 8A




FIG. 8B

9/28




FIG. 9

10/28




FIG. 10

11/28




FIG. 11

12/28




FIG. 12A




FIG. 12B

13/28




FIG. 13A




FIG. 13B

14/28




FIG. 14

15/28




FIG. 15

16/28




FIG. 16A

17/28



FIG. 16B-1

FIG. 16B-2

FIG. 16B-3



FIG. 16B-4

FIG. 16B-5

18/28




FIG. 17




FIG. 18A

1/1 31/11
 atg aca gtg ctg gcg cca gcc tgg agc cca aat tcc tcc ctg ttg ctg ctg ttg ctg ctg
 Met thr val leu ala pro ala trp ser pro asn ser ser leu leu leu leu leu leu
 61/21 91/31
 ctg agt cct tgc ctg cgg ggg aca cct gac tgt tac ttc agc cac agt ccc atc tcc tcc
 leu ser pro cys leu arg gly thr pro asp cys tyr phe ser his ser pro ile ser ser
 121/41 151/51
 aac ttc aaa gtg aag ttt aga gag ttg act gac cac ctg ctt aaa gat tac cca gtc act
 asn phe lys val lys phe arg glu leu thr asp his leu leu lys asp tyr pro val thr
 181/61 211/71
 gtg gcc gtc aat ctt cag gac gag aag cac tgc aag gcc ttg tgg agc ctc ttc cta gcc
 val ala val asn leu gln asp glu lys his cys lys ala leu trp ser leu phe leu ala
 241/81 271/91
 cag cgc tgg ata gag caa ctg aag act gtg gca ggg tct aag atg caa acg ctt ctg gag
 gln arg trp ile glu gln leu lys thr val ala gly ser lys met gln thr leu leu glu
 301/101 331/111
 gac gtc aac acc gag ata cat ttt gtc acc tca tgt acc ttc cag ccc cta cca gaa tgt
 asp val asn thr glu ile his phe val thr ser cys thr phe gln pro leu pro glu cys
 361/121 391/131
 ctg cga ttc gtc cag acc aac atc tcc cac ctc ctg aag gac acc tgc aca cag ctg ctt
 leu arg phe val gln thr asn ile ser his leu leu lys asp thr cys thr gln leu leu
 421/141 451/151
 gct ctg aag ccc tgt atc ggg aag gcc tgc cag aat ttc tct cgg tgc ctg gag gtg cag
 ala leu lys pro cys ile gly lys ala cys gln asn phe ser arg cys leu glu val gln
 481/161 511/171
 tgc cag ccg gac tcc tcc acc ctg ctg ccc cca agg agt ccc ata gcc cta gaa gcc acg
 cys gln pro asp ser ser thr leu leu pro pro arg ser pro ile ala leu glu ala thr
 541/181 571/191
 gag ctc cca gag cct cgg ccc agg cag gga tcc atg cat gga gat aca cct aca ttg cat
 glu leu pro glu pro arg pro arg gln gly ser met his gly asp thr pro thr leu his
 601/201 631/211
 gaa tat atg tta gat ttg caa cca gag aca act gat ctc tac tgt tat gag caa tta aat
 glu tyr met leu asp leu gln pro glu thr thr asp leu tyr cys tyr glu gln leu asn
 661/221 691/231
 gac agc tca gag gag gag gat gaa ata gat ggt cca gct gga caa gca gaa ccg gac aga
 asp ser ser glu glu glu asp glu ile asp gly pro ala gly gln ala glu pro asp arg
 721/241 751/251
 gcc cat tac aat att gta acc ttt tgt tgc aag tgt gac tct acg ctt cgg ttg tgc gta
 ala his tyr asn ile val thr phe cys cys lys cys asp ser thr leu arg leu cys val
 781/261 811/271
 caa agc aca cac gta gac att cgt act ttg gaa gac ctg tta atg ggc aca cta gga att
 gln ser thr his val asp ile arg thr leu glu asp leu leu met gly thr leu gly ile
 841/281 871/291
 gtg tgc ccc atc tgt tct cag gat aag ctt aag ttt aaa ccg ctg atc agc ctc gac tgt
 val cys pro ile cys ser gln asp lys leu lys phe lys pro leu ile ser leu asp cys
 901/301
 gcc ttc tag
 ala phe AMB

FIG. 18B




FIG. 19A

21/28




FIG. 19B

22/28




FIG. 20A




FIG. 20B

23/28




FIG. 21

24/28




FIG. 22A




FIG. 22B




1cm FL-E7

FIG. 23A



1cm CONTROL

FIG. 23B



1cm FL+E7

FIG. 23C



1cm E7

FIG. 23D



1cm FL

FIG. 23E

26/28




FIG. 24

27/28




FIG. 25

28/28




FIG. 26

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US02/02598

A. CLASSIFICATION OF SUBJECT MATTER

IPC(7) : A61K 39/00; C07H 21/04; C12N 15/00, 15/63, 15/85

US CL : 424/192.1; 435/69.7, 320.1, 325; 536/23.1

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 424/192.1; 435/69.7, 320.1, 325; 536/23.1

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)
Please See Continuation Sheet

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	SUZUE, K. et al. Heat shock fusion proteins as vehicles for antigen delivery into the major histocompatibility complex class I presentation pathway. Proc. Natl. Acad. Sci., USA. November 1997, Vol. 94, pages 1316-13151, entire document.	1-9, 48-50, 56, 60-63 and 75-76
Y	CHU, N. R. et al. Immunotherapy of a human papillomavirus (HPV) type 16 E7-expressing tumour by administration of a fusion protein comprising Mycobacterium bovis bacille Calmette-Guerin (BCG) hsp65 and HPV16 E7. Clin. Exp. Immunol. 2000, Vol. 121, pages 216-225, entire document.	1-9, 48-50, 56, 60-63, 75-76
Y	MORE, S. et al. Activation of cytotoxic T cells in vitro by recombinant gp96 fusion proteins irrespective of the 'fused' antigenic peptide sequence. Immunology Letters. 1991, Vol. 69, pages 275-282, entire document.	1-9, 48-50, 56, 60-63, 75-76
A	CHENG et al. Enhancement of Sindbis Virus Self-Replicating RNA Vaccine Potency by Linkage of Mycobacterium tuberculosis Heat Shock Protein 70 Gene to an Antigen Gene. J. Immunology. 2001, Vol. 166, pages 6218-6226.	1-9, 48-50, 56, 60-63, 75-76



Further documents are listed in the continuation of Box C.



See patent family annex.

* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier application or patent published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T"

later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X"

document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y"

document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&"

document member of the same patent family

Date of the actual completion of the international search

06 September 2002 (06.09.2002)

Date of mailing of the international search report

20 SEP 2002

Name and mailing address of the ISA/US

Commissioner of Patents and Trademarks
Box PCT
Washington, D.C. 20231

Facsimile No. (703)305-3230

Authorized officer

Amy DeCloux

Telephone No. 703 308-0196

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US02/02598

C. (Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A,P	HUNG, C.-F. et al. Enhancement of DNA Vaccine Potency by Linkage of Antigen Gene to a Gene Encoding the Extracellular Domain of Fms-like Tyrosine Kinase 3-Ligand. Cancer Research. February 2001, Vol. 61, pages 1080-1088.	1-9, 48-50, 56, 60-63, 75-76

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US02/02598

Box I Observations where certain claims were found unsearchable (Continuation of Item 1 of first sheet)

This international report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claim Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
2. ☐ Claim Nos.:
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:
3. ☒ Claim Nos.: 10-47, 51-55, 57-59, 67-74 and 77
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of Item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

☐
☐

The additional search fees were accompanied by the applicant's protest.

No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US02/02598

Continuation of Item 4 of the first sheet:

Title is longer than 17 words, PCT Rule 4,3. Suggested new title follows:

"Nucleic Acid derived vaccine that encodes an antigen linked to a polypeptide that promotes antigen presentation".

Continuation of B. FIELDS SEARCHED Item 3:

STN/CAS:Medline, CAPLUS, Embase, Biosis, WEST